

# Numerical methods in magnetic resonance imaging: the Bloch-Torrey equation



Jonathan Doucette

University of British Columbia


April 3, 2018



- 1 Animation of a typical “spin echo” MRI sequence


## ① Spins (water molecules) do not truly refocus:

- ① Precession rate depends on position
- ② Diffusion plays an important role




① Spins (water molecules) do not truly refocus:

- ① Precession rate depends on position
- ② Diffusion plays an important role



- ① Spins (water molecules) do not truly refocus:
  - ① Precession rate depends on position
  - ② Diffusion plays an important role






Figure: Simulated imaging voxel




Figure: Cross section of precession rate

- 1 Evolution of the magnetization is modelled by the **Bloch-Torrey equation**

$$\frac{\partial \mathcal{M}}{\partial t} = D \Delta \mathcal{M} - \Gamma \mathcal{M}$$

where:

$$\begin{aligned}\mathcal{M} &= M_x + i M_y \\ \Gamma(\mathbf{x}) &= R(\mathbf{x}) + i \omega(\mathbf{x})\end{aligned}$$

- 2 IC:  $\mathcal{M}(\mathbf{x}, 0) = \mathcal{M}_0(\mathbf{x})$  is given
- 3 BC: zero Neumann or periodic
- 4 Note:

$$D = 0 \Rightarrow \mathcal{M}(\mathbf{x}, t) = \mathcal{M}_0(\mathbf{x}) e^{-\Gamma(\mathbf{x})t}$$

- 1 Evolution of the magnetization is modelled by the **Bloch-Torrey equation**

$$\frac{\partial \mathcal{M}}{\partial t} = D \Delta \mathcal{M} - \Gamma \mathcal{M}$$

where:

$$\begin{aligned}\mathcal{M} &= M_x + i M_y \\ \Gamma(\mathbf{x}) &= R(\mathbf{x}) + i \omega(\mathbf{x})\end{aligned}$$

- 2 IC:  $\mathcal{M}(\mathbf{x}, 0) = \mathcal{M}_0(\mathbf{x})$  is given
- 3 BC: zero Neumann or periodic
- 4 Note:

$$D = 0 \Rightarrow \mathcal{M}(\mathbf{x}, t) = \mathcal{M}_0(\mathbf{x}) e^{-\Gamma(\mathbf{x})t}$$



- 1 Evolution of the magnetization is modelled by the **Bloch-Torrey equation**

$$\frac{\partial \mathcal{M}}{\partial t} = D \Delta \mathcal{M} - \Gamma \mathcal{M}$$

where:

$$\begin{aligned}\mathcal{M} &= M_x + i M_y \\ \Gamma(\mathbf{x}) &= R(\mathbf{x}) + i \omega(\mathbf{x})\end{aligned}$$

- 2 IC:  $\mathcal{M}(\mathbf{x}, 0) = \mathcal{M}_0(\mathbf{x})$  is given
- 3 BC: zero Neumann or periodic
- 4 Note:

$$D = 0 \Rightarrow \mathcal{M}(\mathbf{x}, t) = \mathcal{M}_0(\mathbf{x}) e^{-\Gamma(\mathbf{x})t}$$

- 1 Evolution of the magnetization is modelled by the **Bloch-Torrey equation**

$$\frac{\partial \mathcal{M}}{\partial t} = D \Delta \mathcal{M} - \Gamma \mathcal{M}$$

where:

$$\begin{aligned}\mathcal{M} &= M_x + i M_y \\ \Gamma(\mathbf{x}) &= R(\mathbf{x}) + i \omega(\mathbf{x})\end{aligned}$$

- 2 IC:  $\mathcal{M}(\mathbf{x}, 0) = \mathcal{M}_0(\mathbf{x})$  is given
- 3 BC: zero Neumann or periodic
- 4 Note:

$$D = 0 \Rightarrow \mathcal{M}(\mathbf{x}, t) = \mathcal{M}_0(\mathbf{x}) e^{-\Gamma(\mathbf{x})t}$$

- 1 First, the BT PDE is rewritten in the more suggestive form

$$\frac{\partial \mathcal{M}}{\partial t} = H\mathcal{M}$$

where

$$H = -D\Delta + \Gamma.$$

- 2 Then, the general solution  $\mathcal{M}$  may then be written as

$$\mathcal{M} = e^{-Ht} \mathcal{M}_0$$

where  $e^{-Ht}$  is the *evolution operator*

- 1 First, the BT PDE is rewritten in the more suggestive form

$$\frac{\partial \mathcal{M}}{\partial t} = H\mathcal{M}$$

where

$$H = -D\Delta + \Gamma.$$

- 2 Then, the general solution  $\mathcal{M}$  may then be written as

$$\mathcal{M} = e^{-Ht}\mathcal{M}_0$$

where  $e^{-Ht}$  is the *evolution operator*

- ① Now, the evolution operator may be *split* using the approximation

$$\begin{aligned}e^{-Ht} &= e^{D\Delta t - \Gamma t} \\ &\approx e^{-\Gamma t/2} e^{D\Delta t} e^{-\Gamma t/2} + \mathcal{O}(t^3)\end{aligned}$$

- ② Although  $e^{-Ht}$  has no closed form, the split operators do:

$$\begin{aligned}e^{-\Gamma t/2} \mathcal{M} &= e^{-\Gamma(x)t/2} \odot \mathcal{M} \\ e^{D\Delta t} \mathcal{M} &= \Phi * \mathcal{M}\end{aligned}$$

- ① Now, the evolution operator may be *split* using the approximation

$$\begin{aligned} e^{-Ht} &= e^{D\Delta t - \Gamma t} \\ &\approx e^{-\Gamma t/2} e^{D\Delta t} e^{-\Gamma t/2} + \mathcal{O}(t^3) \end{aligned}$$

- ② Although  $e^{-Ht}$  has no closed form, the split operators do:

$$\begin{aligned} e^{-\Gamma t/2} \mathcal{M} &= e^{-\Gamma(\mathbf{x})t/2} \odot \mathcal{M} \\ e^{D\Delta t} \mathcal{M} &= \Phi * \mathcal{M} \end{aligned}$$

- 1 The BT equation can also be solved using FEM
- 2 First, let  $u = M_x$  and  $v = M_y$  and write:

$$\begin{cases} \frac{\partial u}{\partial t} = D\Delta u - Ru + \omega v \\ \frac{\partial v}{\partial t} = D\Delta v - Rv - \omega u \end{cases}$$

- 1 The BT equation can also be solved using FEM
- 2 First, let  $u = M_x$  and  $v = M_y$  and write:

$$\begin{cases} \frac{\partial u}{\partial t} = D\Delta u - Ru + \omega v \\ \frac{\partial v}{\partial t} = D\Delta v - Rv - \omega u \end{cases}$$



- 1 Applying the method of lines:

$$M^h \mathbf{u}_t = -(DK^h + R^h) \mathbf{u} + W^h \mathbf{v}$$

$$M^h \mathbf{v}_t = -(DK^h + R^h) \mathbf{v} - W^h \mathbf{u}$$

where:

$$R_{ij}^h := \int R \phi_i \phi_j dx$$

$$W_{ij}^h := \int \omega \phi_i \phi_j dx$$

$$M^h \mathbf{u}_t = -(DK^h + R^h) \mathbf{u} + W^h \mathbf{v}$$

$$M^h \mathbf{v}_t = -(DK^h + R^h) \mathbf{v} - W^h \mathbf{u}$$


- 1 The Bloch-Torrey smooths high frequency modes
- 2 Strongly A-stable time stepping methods are required

$$M^h \mathbf{u}_t = -(DK^h + R^h) \mathbf{u} + W^h \mathbf{v}$$

$$M^h \mathbf{v}_t = -(DK^h + R^h) \mathbf{v} - W^h \mathbf{u}$$

- 1 The Bloch-Torrey smooths high frequency modes
- 2 Strongly A-stable time stepping methods are required

# Operator Splitting versus FEM



- ① Splitting methods:
  - + Fast
  - Resolution limitations
  - ± Periodic boundary conditions
  
- ② Finite element methods:
  - Slow
  - + Flexible boundary conditions
  - + Adaptivity?

- ① Splitting methods:
  - + Fast
  - Resolution limitations
  - ± Periodic boundary conditions
  
- ② Finite element methods:
  - Slow
  - + Flexible boundary conditions
  - + Adaptivity?