

Optimization of the Western Canada Power Grid

Incorporation of Renewable Energy

Jun Duan

University of Victoria

10th April 2017



University
of Victoria

Frame Title

Frame Subtitle

- Lorem ipsum dolor sit amet.
- Populo laoreet indoctum ne nec, putent mnesarchum at duo.



Frame Title

Frame Subtitle

- Lorem ipsum dolor sit amet.
- Populo laoreet indoctum ne nec, putent mnesarchum at duo.



Frame Title

Frame Subtitle

- Lorem ipsum dolor sit amet.
- Populo laoreet indoctum ne nec, putent mnesarchum at duo.



Frame Title

Frame Subtitle

- Lorem ipsum dolor sit amet.
- Populo laoreet indoctum ne nec, putent mnesarchum at duo.



Frame Title

Frame Subtitle

- Lorem ipsum dolor sit amet.
- Populo laoreet indoctum ne nec, putent mnesarchum at duo.



Frame Title

Frame Subtitle

- Lorem ipsum dolor sit amet.
- Populo laoreet indoctum ne nec, putent mnesarchum at duo.



Frame Title

Frame Subtitle

- Lorem ipsum dolor sit amet.
- Populo laoreet indoctum ne nec, putent mnesarchum at duo.



Forecasting functional data

Some notation

Let $y_t(x_i)$ be the observed data in period t at location x_i , $i = 1, \dots, p$, $t = 1, \dots, n$.



Forecasting functional data

Some notation

Let $y_t(x_i)$ be the observed data in period t at location x_i , $i = 1, \dots, p$, $t = 1, \dots, n$.

$$y_t(x_i) = f_t(x_i) + \sigma_t(x_i)\varepsilon_{t,i}$$

where $\varepsilon_{t,i}$ is iid $N(0,1)$ and $\sigma_t(x_i)$ allows the amount of noise to vary with x .



Forecasting functional data

Some notation

Let $y_t(x_i)$ be the observed data in period t at location x_i , $i = 1, \dots, p$, $t = 1, \dots, n$.

$$y_t(x_i) = f_t(x_i) + \sigma_t(x_i)\varepsilon_{t,i}$$

where $\varepsilon_{t,i}$ is iid $N(0,1)$ and $\sigma_t(x_i)$ allows the amount of noise to vary with x .

- 1 We assume $f_t(x)$ is a smooth function of x .



Forecasting functional data

Some notation

Let $y_t(x_i)$ be the observed data in period t at location x_i , $i = 1, \dots, p$, $t = 1, \dots, n$.

$$y_t(x_i) = f_t(x_i) + \sigma_t(x_i)\varepsilon_{t,i}$$

where $\varepsilon_{t,i}$ is iid $N(0,1)$ and $\sigma_t(x_i)$ allows the amount of noise to vary with x .

- 1 We assume $f_t(x)$ is a smooth function of x .
- 2 We need to estimate $f_t(x)$ from the data for $x_1 < x < x_p$.

