

# MAT 67 Homework 5

Hardy Jones

999397426

Professor Bandyopadhyay

Fall 2013

1. Let  $V$  be a vector space over  $\mathbb{F}$ , and suppose that the list  $(v_1, v_2, \dots, v_n)$  of vectors spans  $V$ , where each  $v_i \in V$ . Prove that the list

$$(v_1 - v_2, v_2 - v_3, v_3 - v_4, \dots, v_{n-2} - v_{n-1}, v_{n-1} - v_n, v_n)$$

also spans  $V$ .

*Proof.* Each  $v_j \in (v_1, v_2, \dots, v_n)$  can be constructed from our new list.

$$\begin{aligned} v_1 &= (v_1 - v_2) + (v_2 - v_3) + (v_3 - v_4) + \dots + (v_{n-2} - v_{n-1}) + (v_{n-1} - v_n) + v_n \\ &= (v_1 - \cancel{v_2}) + (\cancel{v_2} - \cancel{v_3}) + (\cancel{v_3} - \cancel{v_4}) + \dots + (\cancel{v_{n-2}} - \cancel{v_{n-1}}) + (\cancel{v_{n-1}} - \cancel{v_n}) + \cancel{v_n} \\ &= v_1 \\ v_2 &= 0(v_1 - v_2) + (v_2 - v_3) + (v_3 - v_4) + \dots + (v_{n-2} - v_{n-1}) + (v_{n-1} - v_n) + v_n \\ &= (v_2 - \cancel{v_3}) + (\cancel{v_3} - \cancel{v_4}) + \dots + (\cancel{v_{n-2}} - \cancel{v_{n-1}}) + (\cancel{v_{n-1}} - \cancel{v_n}) + \cancel{v_n} \\ &= v_2 \\ &\vdots \\ v_n &= 0(v_1 - v_2) + 0(v_2 - v_3) + 0(v_3 - v_4) + \dots + 0(v_{n-2} - v_{n-1}) + 0(v_{n-1} - v_n) + v_n \\ &= v_n \end{aligned}$$

Since we see that we can generate each one of these, we can generate the entire list  $(v_1, v_2, \dots, v_n)$ , which spans  $V$ . So,  $\text{span}(v_1 - v_2, v_2 - v_3, \dots, v_{n-1} - v_n, v_n) = V$   $\square$