

MAT 108 HW 7

Hardy Jones

999397426

Professor Bandyopadhyay

Spring 2015

§3.2 5 (b) *Proof.* We need to show reflexivity, symmetry, and transitivity.

- Choose any $m \in \mathbb{N}$.

Then m has a digit $p \in \{0, 1, \dots, 9\}$ in its tens place. Since m has the same digit p in its tens place as m , we have mRm .

So R is reflexive on m

Since the choice of m was arbitrary, this result holds for all $m \in \mathbb{N}$.

Thus, R is reflexive on \mathbb{N}

- Choose any $m, n \in \mathbb{N}$ with mRn .

Then m has the same digit in the tens place as n . Then n has the same digit in the tens place as m . Then nRm . So R is symmetric on m, n .

Since the choice of m, n was arbitrary, this result holds for all $m, n \in \mathbb{N}$.

Thus, R is symmetric on \mathbb{N}

- Choose any $m, n, p \in \mathbb{N}$ with mRn and nRp .

Then m has the same digit in the tens place as n . Call this digit q . Then n has the same digit in the tens place as p . Call this digit r .

Since q and r are the tens digit of n , we have $q = r$. So we know that the tens place of p is q .

This means that m has the same digit in the tens place as p . So mRp .

And R is transitive on m, n, p .

Since the choice of m, n, p was arbitrary, this result holds for all $m, n, p \in \mathbb{N}$.

Thus, R is transitive on \mathbb{N}

□

An element of $106/R$ less than 50 is 3, since the tens place of 106 is 0, the tens place of 3 is 0, and $3 < 50$.

An element of $106/R$ between 150 and 300 is 203, since the tens place of 106 is 0, the tens place of 203 is 0, and $150 < 203 < 300$.

An element of $106/R$ greater than 1000 is 2003, since the tens place of 106 is 0, the tens place of 2003 is 0, and $1000 < 2003$.

Three elements of $635/R$ are 30, 31, and 32, since the tens place of 635 is 3, the tens place of 30 is 3, the tens place of 31 is 3, and the tens place of 32 is 3.

(c) *Proof.* We need to show reflexivity, symmetry, and transitivity.

- Choose any $m \in \mathbb{R}$.
Then
So V is reflexive on m
Since the choice of m was arbitrary, this result holds for all $m \in \mathbb{R}$.
Thus, V is reflexive on \mathbb{R}
- Choose any $m, n \in \mathbb{R}$ with mVn .
Then m has the same digit in the tens place as n . Then n has the same digit in the tens place as m . Then nVm . So V is symmetric on m, n .
Since the choice of m, n was arbitrary, this result holds for all $m, n \in \mathbb{R}$.
Thus, V is symmetric on \mathbb{R}
- Choose any $m, n, p \in \mathbb{R}$ with mVn and nVp .
Then m has the same digit in the tens place as n . Call this digit q . Then n has the same digit in the tens place as p . Call this digit r .
Since q and r are the tens digit of n , we have $q = r$. So we know that the tens place of p is q .
This means that m has the same digit in the tens place as p . So mVp .
And V is transitive on m, n, p .
Since the choice of m, n, p was arbitrary, this result holds for all $m, n, p \in \mathbb{R}$.
Thus, V is transitive on \mathbb{R}

□

An element of $106/R$ less than 50 is 3, since the tens place of 106 is 0, the tens place of 3 is 0, and $3 < 50$.

An element of $106/R$ between 150 and 300 is 203, since the tens place of 106 is 0, the tens place of 203 is 0, and $150 < 203 < 300$.

An element of $106/R$ greater than 1000 is 2003, since the tens place of 106 is 0, the tens place of 2003 is 0, and $1000 < 2003$.

Three elements of $635/R$ are 30, 31, and 32, since the tens place of 635 is 3, the tens place of 30 is 3, the tens place of 31 is 3, and the tens place of 32 is 3.

(d)

(e)

(h)

8 (d) The equivalence classes for the relation congruence modulo 7 are:

$$\begin{aligned}\bar{0} &= \{\dots, -14, -7, 0, 7, 14, \dots\} \\ \bar{1} &= \{\dots, -13, -6, 1, 8, 15, \dots\} \\ \bar{2} &= \{\dots, -12, -5, 2, 9, 16, \dots\} \\ \bar{3} &= \{\dots, -11, -4, 3, 10, 17, \dots\} \\ \bar{4} &= \{\dots, -10, -3, 4, 11, 18, \dots\} \\ \bar{5} &= \{\dots, -9, -2, 5, 12, 19, \dots\} \\ \bar{6} &= \{\dots, -8, -1, 6, 13, 20, \dots\}\end{aligned}$$

9 We use the equivalent definition that $a \equiv b \pmod{m}$ is the same as $a - b = km$ for some $k \in \mathbb{Z}$.

(d) We want to find an a such that $a - 3 = 4k$ and $a - 3 = 5l$.
We can solve this algebraically by substitution.

$$\begin{aligned}a - 3 &= 4k \\ (5l + 3) - 3 &= 4k && \text{since } a = 5l + 3 \\ 5l &= 4k \\ l &= \frac{4}{5}k\end{aligned}$$

Since $k, l \in \mathbb{Z}$ we know that k must be a multiple of 5.
Choose $k = 5$. then we have $l = \frac{4}{5}5 = 4$, and we end up with

$$\begin{aligned}a - 3 &= 5l \\ &= 5(4) \\ &= 20 \\ a &= 23\end{aligned}$$

Choose $k = -5$. then we have $l = \frac{4}{5}(-5) = -4$, and we end up with

$$\begin{aligned}a - 3 &= 5l \\ &= 5(-4) \\ &= -20 \\ a &= -17\end{aligned}$$

So a positive integer that is congruent to 3 (mod 4) and congruent to 3 (mod 5) is 23.

And a negative integer that is congruent to 3 (mod 4) and congruent to 3 (mod 5) is -17.

- (e) We want to find an a such that $a - 1 = 3k$ and $a - 1 = 7l$.
We can solve this algebraically by substitution.

$$\begin{aligned} a - 1 &= 3k \\ (7l + 1) - 1 &= 3k && \text{since } a = 7l + 1 \\ 7l &= 3k \\ l &= \frac{3}{7}k \end{aligned}$$

Since $k, l \in \mathbb{Z}$ we know that k must be a multiple of 7.
Choose $k = 7$. then we have $l = \frac{3}{7}7 = 3$, and we end up with

$$\begin{aligned} a - 1 &= 7l \\ &= 7(3) \\ &= 21 \\ a &= 22 \end{aligned}$$

Choose $k = -7$. then we have $l = \frac{3}{7}(-7) = -3$, and we end up with

$$\begin{aligned} a - 1 &= 7l \\ &= 7(-3) \\ &= -21 \\ a &= -20 \end{aligned}$$

So a positive integer that is congruent to 1 (mod 3) and congruent to 1 (mod 7) is 22.

And a negative integer that is congruent to 1 (mod 3) and congruent to 1 (mod 7) is -20.

- 11 *Proof.* We can show that S is not an equivalence relation by showing that S is not transitive.

Choose $x = 1, y = 2, z = 4$. Then $x, y, z, w \in \mathbb{N}$ and $1 + 2 = 3 = 3(1), 2 + 4 = 6 = 3(2)$, so xSy and ySz .

Now, if S were transitive, we'd have xSz . However, $1 + 4 = 5$ and 3 does not divide 5.

So S is not transitive.

Thus, S is not an equivalence relation. □

- §3.3 3 (a) The relation R creates equivalence classes where each x in each class has the same fractional part.

The following are examples of these equivalence classes:

$$\mathbb{Z}, \{\dots, -1.5, 0.5, 1.5, \dots\}, \{\dots, e-3, e-2, e-1, e, e+1, \dots\}$$

Then we have the partition is the family of sets of the form:

$$A_r = \{x \in \mathbb{R} : x - \lfloor x \rfloor = r\}, \text{ where } r \in [0, 1)$$

(d) For any $x, y \in \mathbb{R}$, x and y are in the same equivalence class iff $x = \pm y$.

The following are examples of these equivalence classes:

$$\{0\}, \{-1, 1\}, \left\{-\frac{3}{7}, \frac{3}{7}\right\}, \{-e, e\}$$

Then we have the partition as the family of sets of the form:

$$A_r = \{r, -r\}, \text{ where } r \in \mathbb{R}$$

- 4
- 6 (a)
(e)
- §4.1 1 (e)
(f)
- 3 (b)
(e)
(f)
- 4 (e)
(f)
- 5 (b)
(c)
- 8 (b)
(c)
- 9 (b)
(d)
- 10 (b)
(d)
- 11 (c)
(d)
- 17 (a)
(b)