

# MAT 25 Homework 5

Hardy Jones  
999397426  
Professor Bae  
Fall 2013

## 1. 2.2.1

(a)  $\lim_{n \rightarrow \infty} \frac{1}{6n^2+1} = 0$

We need to show

$$\begin{aligned}\frac{1}{6n^2+1} &< \epsilon \\ \frac{1}{\epsilon} &< 6n^2+1 \\ \frac{1}{\epsilon} - 1 &< 6n^2 \\ \frac{1-\epsilon}{\epsilon} &< 6n^2 \\ \frac{1-\epsilon}{6\epsilon} &< n^2 \\ \sqrt{\frac{1-\epsilon}{6\epsilon}} &< n\end{aligned}$$

Let  $\epsilon > 0$ . Choose  $N \in \mathbb{N} | N > \sqrt{\frac{1-\epsilon}{6\epsilon}}$ .

Let  $n \geq N$ . So,  $n \geq N > \sqrt{\frac{1-\epsilon}{6\epsilon}} \implies \frac{1}{6n^2+1} < \epsilon$

Thus  $|a_n - 0| < \epsilon$ .

(b)  $\lim_{n \rightarrow \infty} \frac{3n+1}{2n+5} = \frac{3}{2}$

We need to show

$$\begin{aligned}\frac{3n+1}{2n+5} &< \epsilon \\ 3n+1 &< 2n\epsilon+5\epsilon \\ 1-5\epsilon &< (2\epsilon-3)n \\ \frac{1-5\epsilon}{2\epsilon-3} &< n\end{aligned}$$

Let  $\epsilon > 0$ . Choose  $N \in \mathbb{N} | N > \frac{1-5\epsilon}{2\epsilon-3}$ .

Let  $n \geq N$ . So,  $n \geq N > \frac{1-5\epsilon}{2\epsilon-3} \implies \frac{3n+1}{2n+5} < \epsilon$ .

Thus  $|a_n - \frac{3}{2}| < \epsilon$ .

(c)  $\lim \frac{2}{\sqrt{n+3}} = 0$

We need to show

$$\begin{aligned}\frac{2}{\sqrt{n+3}} &< \epsilon \\ \frac{2}{\epsilon} &< \sqrt{n+3} \\ \frac{4}{\epsilon^2} &< n+3 \\ \frac{4}{\epsilon^2} - 3 &< n\end{aligned}$$

Let  $\epsilon > 0$ . Choose  $N \in \mathbb{N} | N > \frac{4}{\epsilon^2} - 3$ .

Let  $n \geq N$ . So,  $n \geq N > \frac{4}{\epsilon^2} - 3 \implies \frac{2}{\sqrt{n+3}} < \epsilon$ .

Thus  $|a_n - 0| < \epsilon$ .

## 2. 2.2.5

(a)  $a_n = \lfloor \frac{1}{n} \rfloor$

It is easy to see that after the first element in the sequence, all values are 0.

$$\lim a_n = 0$$

*Proof.* Let  $\epsilon > 0$ . Choose  $N > 1$ .

Let  $n \geq N$ . So,  $n \geq N > 1 \implies \lfloor \frac{1}{n} \rfloor = 0 < \epsilon$ .

Thus,  $|a_n - 0| < \epsilon$ . □

(b)  $a_n = \lfloor \frac{10+n}{2n} \rfloor$

Again we see that after some elements all values are 0.

$$\lim a_n = 0$$

*Proof.* Let  $\epsilon > 0$ . Choose  $N > 10$ .

Let  $n \geq N$ . So,  $n \geq N > 10 \implies \lfloor \frac{10+n}{2n} \rfloor = 0 < \epsilon$ .

Thus,  $|a_n - 0| < \epsilon$ . □

## 3. 2.2.7

(a) A sequence  $(a_n)$  diverges to  $\infty$  if, for every positive number  $\epsilon$ , there exists an  $N \in \mathbb{N}$  such that whenever  $n \geq N$  it follows that  $|a_n| > \epsilon$

$$\lim \sqrt{n} = \infty$$

*Proof.* Let  $\epsilon > 0$ .

We want to show

$$\begin{aligned}\sqrt{n} &> \epsilon \\ n &> \epsilon^2\end{aligned}$$

Choose  $N > \epsilon^2$ .

Let  $n \geq N$ . So,  $n \geq N > \epsilon^2 \implies \sqrt{n} > \epsilon$ .

Thus,  $|a_n| > \epsilon$ . □

- (b) It states that this particular sequence does not diverge to  $\infty$ . The reason being, if you choose some  $\epsilon > 0$ , and any  $N \in \mathbb{N}$ , then  $\exists n \geq N$  either  $n = 0$  or  $n + 1 = 0$ . So,  $|a_n| \not\geq \epsilon, \forall n$ .

4. 2.3.4 Using the Algebraic Limit Theorem, if  $\lim a_n = l_1$  and  $\lim a_n = l_2$ , then

$$\lim(a_n - a_n) = 0$$

$$l_1 - l_2 = 0$$

$$l_1 = l_2$$

5. 2.3.7

- (a) Since  $(a_n)$  is bounded,  $\exists M > 0$  such that  $|a_n| \leq M, \forall n \in \mathbb{N}$ .

Also, since  $\lim b_n = 0$  we can choose a special epsilon,  $\forall \epsilon > 0, \exists N \in \mathbb{N}$  such that  $n \geq N \implies |b_n| < \frac{\epsilon}{M}$

So,  $|a_n||b_n| < M \frac{\epsilon}{M} \implies |a_n||b_n| < \epsilon$ .

Which is equivalent to  $|a_n b_n - 0| < \epsilon$ .

By Definition 2.2.3, this sequence  $(a_n b_n)$  goes to 0.

We were not allowed to use the Algebraic Limit Theorem because  $(a_n)$  is not necessarily convergent.

- (b) The only thing we can conclude when  $\lim b_n = b$  is that  $(a_n b_n)$  is bounded by  $b$  times the bounds of  $(a_n)$ .

6. 2.3.11 Since  $(x_n)$  converges to some number  $X$  we know  $\forall \epsilon > 0, \exists N \in \mathbb{N}$  such that  $n \geq N \implies |x_n| < \epsilon$

$$\begin{aligned}
|y_n - X| &= \left| \frac{a_1 + a_2 + \cdots + a_n}{n} - X \right| \\
&= \left| \frac{a_1 + a_2 + \cdots + a_n}{n} - \frac{nX}{n} \right| \\
&= \left| \frac{a_1 + a_2 + \cdots + a_n - nX}{n} \right| \\
&= \left| \frac{(a_1 - X) + (a_2 - X) + \cdots + (a_n - X)}{n} \right| \\
&= \left| \frac{a_1 - X}{n} + \frac{a_2 - X}{n} + \cdots + \frac{a_n - X}{n} \right| \\
&= \left| \frac{a_1 - X}{n} \right| + \left| \frac{a_2 - X}{n} \right| + \cdots + \left| \frac{a_n - X}{n} \right| \\
&= \frac{|a_1 - X|}{n} + \frac{|a_2 - X|}{n} + \cdots + \frac{|a_n - X|}{n} \\
&< \frac{\epsilon}{n} + \frac{\epsilon}{n} + \cdots + \frac{\epsilon}{n} \\
&< \epsilon
\end{aligned}$$

So,  $\lim y_n = X$ .

An example of a sequence  $(x_n)$  that doesn't converge, but  $(y_n)$  does is  $x_n = (-1)^n$

We can see this by looking at some terms

$$\begin{aligned}
x_1 &= -1 \\
x_2 &= 1 \\
x_3 &= -1 \\
x_4 &= 1 \\
x_5 &= -1 \\
x_6 &= 1 \\
x_7 &= -1 \\
&\vdots
\end{aligned}$$

Which clearly doesn't converge

$$\begin{aligned}
y_1 &= \frac{-1}{1} &= -1 \\
y_2 &= \frac{-1+1}{2} &= 0 \\
y_3 &= \frac{-1+1-1}{3} &= -\frac{1}{3} \\
y_4 &= \frac{-1+1-1+1}{4} &= 0 \\
y_5 &= \frac{-1+1-1+1-1}{5} &= -\frac{1}{5} \\
y_6 &= \frac{-1+1-1+1-1+1}{6} &= 0 \\
y_7 &= \frac{-1+1-1+1-1+1-1}{7} &= -\frac{1}{7} \\
&\vdots
\end{aligned}$$

It's easy to see that this sequence converges to 0.

(2)

Suppose  $(a_n)$  is a sequence such that  $a_n \in \mathbb{Z}$ . Suppose  $(a_n)$  converges to  $a$ .  
Prove that  $a \in \mathbb{Z}$ .

*Proof.*

In order for  $(a_n)$  to converge to  $a$  the follow must hold:  $|a_n - a| < \epsilon$  for any  $\epsilon > 0$

Now, the only way this statement is true, is if  $a_n - a = 0 \implies a_n = a$ .

Since all  $a_n \in \mathbb{Z}$  it follows that  $a \in \mathbb{Z}$

□