

ECS 122A Homework 2

Hardy Jones

999397426

Professor Bai

Spring 2014

1. (a) We can expand some of the polynomial

$$(n+2)^{10} = n^{10} + 20n^9 + \dots + 1024$$

Looking at everything except the first term, we see that the greatest power is n^9 . We can set a *little-oh* bound of n^{10} , because $\forall c > 0, \exists n_0 > 0$ such that $0 \leq 20n^9 + \dots + 1024 < cn^{10}, \forall n \geq n_0$. That is:

$$(n+2)^{10} = n^{10} + o(n^{10})$$

Now, this is $\Theta(n^{10})$ because

$\exists c_1 > 0, c_2 > 0, n_0 > 0$ such that $0 \leq c_1 n^{10} \leq n^{10} + o(n^{10}) \leq c_2 n^{10}, \forall n \geq n_0$

This simplifies to $0 \leq c_1 \leq 1 + \frac{o(n^{10})}{n^{10}} \leq c_2$.

Now, since $o(n^{10})$ is an upper bound we know the largest the term $\frac{o(n^{10})}{n^{10}}$ can be is 1. So, this further simplifies to $0 \leq c_1 \leq 2 \leq c_2$.

So this is $\Theta(n^{10})$ if we choose appropriate constants. For instance we can choose $c_1 = c_2 = 2$.

Thus, $(n+2)^{10} = \Theta(n^{10})$

- (b) We can apply reasoning similar to the previous problem.

We can expand some of the polynomial

$$(n+a)^b = n^b + \binom{b}{1} n^{b-1} a + \dots + a^b$$

Looking at everything except the first term, we see that the greatest power is n^{b-1} . We can set a *little-oh* bound of n^b , because $\forall c > 0, \exists n_0 > 0$ such that $0 \leq \binom{b}{1} n^{b-1} a + \dots + a^b < cn^b, \forall n \geq n_0$. That is:

$$(n+a)^b = n^b + o(n^b)$$

Now, this is $\Theta(n^b)$ because

$\exists c_1 > 0, c_2 > 0, n_0 > 0$ such that $0 \leq c_1 n^b \leq n^b + o(n^b) \leq c_2 n^b, \forall n \geq n_0$

This simplifies to $0 \leq c_1 \leq 1 + \frac{o(n^b)}{n^b} \leq c_2$.

Now, since $o(n^b)$ is an upper bound we know the largest the term $\frac{o(n^b)}{n^b}$ can be is 1. So, this further simplifies to $0 \leq c_1 \leq 2 \leq c_2$.

So this is $\Theta(n^b)$ if we choose appropriate constants. For instance we can choose $c_1 = c_2 = 2$.

Thus, $(n+a)^b = \Theta(n^b)$