

# Virtual homological torsion in low dimensions

University of Haifa, Dec. 14 2025

---

Jonathan Fruchter

University of Bonn

# Overview

---

# Overview

---

- Virtual homological torsion: an introduction

# Overview

---

- Virtual homological torsion: an introduction
- A 2-dimensional setting

# Overview

---

- Virtual homological torsion: an introduction
- A 2-dimensional setting
-  Detour: an application

# Overview

---

- Virtual homological torsion: an introduction
- A 2-dimensional setting
-  Detour: an application
- Further results

# Overview

---

- Virtual homological torsion: an introduction
- A 2-dimensional setting
-  Detour: an application
- Further results
- What's next?

# A mystery in 3-dimensions

---

# A mystery in 3-dimensions

In hyperbolic 3-manifolds:

**topological complexity of finite covers grows at a rate reflecting geometry**



# A mystery in 3-dimensions

In hyperbolic 3-manifolds:

**topological complexity of finite covers grows at a rate reflecting geometry**

**Conjecture (Bergeron–Venkatesh)**

Let  $M$  be a closed, hyperbolic 3-manifold and let

$$M = M_0 \leftarrow M_1 \leftarrow M_2 \leftarrow \dots$$

be a cofinal tower of finite-sheeted normal covers of  $M$ , that is,

$$\bigcap_n \pi_1(M_n) = 1.$$

Then

$$\lim_{n \rightarrow \infty} \frac{\log(|\text{Tor}(H_1(M_n; \mathbb{Z}))|)}{[\pi_1(M) : \pi_1(M_n)]} = \frac{\text{vol}(M)}{6\pi}.$$



# At the crossroads

---

## At the crossroads

---

- **Geometry  $\rightsquigarrow$  Topology:** “Gauss-Bonnet on steroids” –

$$\chi(\Sigma) = \frac{K \cdot \text{Area}(\Sigma)}{2\pi} \quad (\Sigma \text{ closed, orientable, constant curvature } K).$$

## At the crossroads

---

- **Geometry  $\rightsquigarrow$  Topology:** “Gauss-Bonnet on steroids” –

$$\chi(\Sigma) = \frac{K \cdot \text{Area}(\Sigma)}{2\pi} \quad (\Sigma \text{ closed, orientable, constant curvature } K).$$

- **Topology  $\rightsquigarrow$  Analysis:** Prediction  $\text{vol}(M)/6\pi$  obtained using analytic methods on the universal cover  $\tilde{M} \cong \mathbb{H}^3$  of  $M$  ( $L^2$ -torsion, Lück–Schick ’99).

## At the crossroads

---

- **Geometry  $\rightsquigarrow$  Topology:** “Gauss-Bonnet on steroids” –

$$\chi(\Sigma) = \frac{K \cdot \text{Area}(\Sigma)}{2\pi} \quad (\Sigma \text{ closed, orientable, constant curvature } K).$$

- **Topology  $\rightsquigarrow$  Analysis:** Prediction  $\text{vol}(M)/6\pi$  obtained using analytic methods on the universal cover  $\tilde{M} \cong \mathbb{H}^3$  of  $M$  ( $L^2$ -torsion, Lück–Schick ’99).
- **Geometry  $\rightsquigarrow$  Number theory:** Roughly, in arithmetic groups, homological torsion parametrizes field extensions  $K/\mathbb{Q}$ .

# At the crossroads

- **Geometry  $\rightsquigarrow$  Topology:** “Gauss-Bonnet on steroids” –

$$\chi(\Sigma) = \frac{K \cdot \text{Area}(\Sigma)}{2\pi} \quad (\Sigma \text{ closed, orientable, constant curvature } K).$$

- **Topology  $\rightsquigarrow$  Analysis:** Prediction  $\text{vol}(M)/6\pi$  obtained using analytic methods on the universal cover  $\tilde{M} \cong \mathbb{H}^3$  of  $M$  ( $L^2$ -torsion, Lück–Schick ’99).
- **Geometry  $\rightsquigarrow$  Number theory:** Roughly, in arithmetic groups, homological torsion parametrizes field extensions  $K/\mathbb{Q}$ .

BV Conjecture:  $\forall p,$  
$$\lim_{n \rightarrow \infty} \frac{\log(|\text{Tor}_p(H_1(M_n; \mathbb{Z}))|)}{[\pi_1(M): \pi_1(M_n)]} = 0,$$

so torsion must involve larger and larger sporadic primes.

# Beyond 3-manifolds

## Open problem

Is there a finitely presented residually finite group  $G$ , and a residual normal chain

$$G \triangleright G_1 \triangleright G_2 \triangleright \dots$$

(i.e.  $\bigcap_n G_n = \{1\}$ ), such that

$$\lim_{n \rightarrow \infty} \frac{\log(|\text{Tor}(G_n^{\text{ab}})|)}{[G : G_n]} > 0 ?$$

## Relaxations: examples in every direction

---

## Relaxations: examples in every direction

---

- **Finite presentation:** (Kar–Kropholler–Nikolov '17) For every  $f : \mathbb{N} \longrightarrow \mathbb{N}$  there is a fg  $G$  and an exhausting normal chain  $G \triangleright G_1 \triangleright G_2 \triangleright \dots$  such that  $|\text{Tor}(G^{\text{ab}})| > f([G : G_n])$ .

## Relaxations: examples in every direction

---

- **Finite presentation:** (Kar–Kropholler–Nikolov '17) For every  $f : \mathbb{N} \longrightarrow \mathbb{N}$  there is a fg  $G$  and an exhausting normal chain  $G \triangleright G_1 \triangleright G_2 \triangleright \dots$  such that  $|\text{Tor}(G^{ab})| > f([G : G_n])$ .
- **Trivial intersection:** (Silver–Williams '02) Cyclic covers of knot complements in  $\mathbb{S}^3$ .



## Relaxations: examples in every direction

---

- **Finite presentation:** (Kar–Kropholler–Nikolov '17) For every  $f : \mathbb{N} \longrightarrow \mathbb{N}$  there is a fg  $G$  and an exhausting normal chain  $G \triangleright G_1 \triangleright G_2 \triangleright \dots$  such that  $|\text{Tor}(G^{ab})| > f([G : G_n])$ .
- **Trivial intersection:** (Silver–Williams '02) Cyclic covers of knot complements in  $\mathbb{S}^3$ .
- **Dimension:** (Avramidi–Okun–Schreve '21) Right-angled Artin groups where  $\log |\text{Tor}(H_2(–; \mathbb{Z}))|$  grows exponentially in the index.



## Relaxations: examples in every direction

- **Finite presentation:** (Kar–Kropholler–Nikolov '17) For every  $f : \mathbb{N} \longrightarrow \mathbb{N}$  there is a fg  $G$  and an exhausting normal chain  $G \triangleright G_1 \triangleright G_2 \triangleright \dots$  such that  $|\text{Tor}(G^{\text{ab}})| > f([G : G_n])$ .
- **Trivial intersection:** (Silver–Williams '02) Cyclic covers of knot complements in  $\mathbb{S}^3$ . 
- **Dimension:** (Avramidi–Okun–Schreve '21) Right-angled Artin groups where  $\log |\text{Tor}(H_2(–; \mathbb{Z}))|$  grows exponentially in the index.
- **Normality:** (Liu '19) Cofinal towers of finite-sheeted *non-normal* covers of closed, hyperbolic  $M^3$  with exponential growth.

## From above

---

## From above

---

Subexponential upper bounds (usually) come in two flavours:

## From above

---

Subexponential upper bounds (usually) come in two flavours:

- **Amenability.** More generally, infinite amenable normal subgroups (+ mild assumptions) (Lück '13).

## From above

---

Subexponential upper bounds (usually) come in two flavours:

- **Amenability**. More generally, infinite amenable normal subgroups (+ mild assumptions) (Lück '13).
- **Cheap rebuilding** (Abert–Bergeron–Frączyk–Gaboriau '24).




## From above

---

Subexponential upper bounds (usually) come in two flavours:

- **Amenability**. More generally, infinite amenable normal subgroups (+ mild assumptions) (Lück '13).
- **Cheap rebuilding** (Abert–Bergeron–Frączyk–Gaboriau '24).




## From above

---

Subexponential upper bounds (usually) come in two flavours:

- **Amenability**. More generally, infinite amenable normal subgroups (+ mild assumptions) (Lück '13).
- **Cheap rebuilding** (Abert–Bergeron–Frączyk–Gaboriau '24).



## Back to 3-manifolds

---

## Back to 3-manifolds

---

### Theorem (Sun '15)

Let  $M$  be a closed hyperbolic 3-manifold and let  $A$  be a finite abelian group. There exists a finite-sheeted cover  $\hat{M} \rightarrow M$  such that

$$H_1(\hat{M}; \mathbb{Z}) = A \oplus \cdots.$$

## Back to 3-manifolds

### Theorem (Sun '15)

Let  $M$  be a closed hyperbolic 3-manifold and let  $A$  be a finite abelian group. There exists a finite-sheeted cover  $\hat{M} \rightarrow M$  such that

$$H_1(\hat{M}; \mathbb{Z}) = A \oplus \cdots.$$

← “Abundance of virtual homological torsion”

## Back to 3-manifolds

### Theorem (Sun '15)

Let  $M$  be a closed hyperbolic 3-manifold and let  $A$  be a finite abelian group. There exists a finite-sheeted cover  $\hat{M} \rightarrow M$  such that

$$H_1(\hat{M}; \mathbb{Z}) = A \oplus \dots .$$

“Abundance of virtual homological torsion”

Similar proof works for all 3-manifolds with empty or toroidal boundary which are not graph manifolds (Chu–Groves '22).

## Back to 3-manifolds

### Theorem (Sun '15)

Let  $M$  be a closed hyperbolic 3-manifold and let  $A$  be a finite abelian group. There exists a finite-sheeted cover  $\widehat{M} \rightarrow M$  such that

$$\rightarrow H_1(\widehat{M}; \mathbb{Z}) = A \oplus \dots .$$

“Abundance of virtual homological torsion”

Similar proof works for all 3-manifolds with empty or toroidal boundary which are not graph manifolds (Chu–Groves '22).

In graph manifolds:


- Solv or Seifert-fibred – restrictions on torsion.
- Non-trivial JSJ – any number can divide  $|\text{Tor}(H_1(\widehat{M}; \mathbb{Z}))|$  (F–Hughes–Valiuunas).

# Virtual homological torsion in 3-manifolds: a recipe


---

# Virtual homological torsion in 3-manifolds: a recipe


---




# Virtual homological torsion in 3-manifolds: a recipe




# Virtual homological torsion in 3-manifolds: a recipe




# Virtual homological torsion in 3-manifolds: a recipe




# Virtual homological torsion in 3-manifolds: a recipe




# Virtual homological torsion in 3-manifolds: a recipe



# Virtual homological torsion in 3-manifolds: a recipe



# Virtual homological torsion in 3-manifolds: a recipe




## (Hyperbolic) Graphs of free groups with cyclic edges

---

(Fundamental groups of) graphs joined together by cylinders.


## (Hyperbolic) Graphs of free groups with cyclic edges

(Fundamental groups of) graphs joined together by cylinders.




# (Hyperbolic) Graphs of free groups with cyclic edges

(Fundamental groups of) graphs joined together by cylinders.



# (Hyperbolic) Graphs of free groups with cyclic edges


(Fundamental groups of) graphs joined together by cylinders.



# (Hyperbolic) Graphs of free groups with cyclic edges


---

(Fundamental groups of) graphs joined together by cylinders.




# (Hyperbolic) Graphs of free groups with cyclic edges

(Fundamental groups of) graphs joined together by cylinders.




## (Hyperbolic) Graphs of free groups with cyclic edges

(Fundamental groups of) graphs joined together by cylinders.





## (Hyperbolic) Graphs of free groups with cyclic edges

(Fundamental groups of) graphs joined together by cylinders.





## (Hyperbolic) Graphs of free groups with cyclic edges

(Fundamental groups of) graphs joined together by cylinders.






# (Hyperbolic) Graphs of free groups with cyclic edges

(Fundamental groups of) graphs joined together by cylinders.





# (Hyperbolic) Graphs of free groups with cyclic edges

(Fundamental groups of) graphs joined together by cylinders.






# (Hyperbolic) Graphs of free groups with cyclic edges

(Fundamental groups of) graphs joined together by cylinders.






# (Hyperbolic) Graphs of free groups with cyclic edges

(Fundamental groups of) graphs joined together by cylinders.






# (Hyperbolic) Graphs of free groups with cyclic edges

(Fundamental groups of) graphs joined together by cylinders.






# (Hyperbolic) Graphs of free groups with cyclic edges

(Fundamental groups of) graphs joined together by cylinders.







# (Hyperbolic) Graphs of free groups with cyclic edges

(Fundamental groups of) graphs joined together by cylinders.







# (Hyperbolic) Graphs of free groups with cyclic edges

(Fundamental groups of) graphs joined together by cylinders.






# (Hyperbolic) Graphs of free groups with cyclic edges

(Fundamental groups of) graphs joined together by cylinders.







# (Hyperbolic) Graphs of free groups with cyclic edges

(Fundamental groups of) graphs joined together by cylinders.







# (Hyperbolic) Graphs of free groups with cyclic edges

(Fundamental groups of) graphs joined together by cylinders.








# (Hyperbolic) Graphs of free groups with cyclic edges

(Fundamental groups of) graphs joined together by cylinders.








# (Hyperbolic) Graphs of free groups with cyclic edges

(Fundamental groups of) graphs joined together by cylinders.







# (Hyperbolic) Graphs of free groups with cyclic edges

(Fundamental groups of) graphs joined together by cylinders.







# (Hyperbolic) Graphs of free groups with cyclic edges

(Fundamental groups of) graphs joined together by cylinders.




# (Hyperbolic) Graphs of free groups with cyclic edges

(Fundamental groups of) graphs joined together by cylinders.






# (Hyperbolic) Graphs of free groups with cyclic edges

(Fundamental groups of) graphs joined together by cylinders.




Free and surface groups  
(and their free products)






# (Hyperbolic) Graphs of free groups with cyclic edges

(Fundamental groups of) graphs joined together by cylinders.




Free and surface groups  
(and their free products)





# (Hyperbolic) Graphs of free groups with cyclic edges


(Fundamental groups of) graphs joined together by cylinders.



Free and surface groups  
(and their free products)




“More than a surface”





# (Hyperbolic) Graphs of free groups with cyclic edges


(Fundamental groups of) graphs joined together by cylinders.



Free and surface groups  
(and their free products)



“More than a surface”



“Has a rigid vertex”

## Homological torsion: Existence

---

# Homological torsion: Existence

## Theorem (Ascari-F '25)

Let  $G$  be a hyperbolic graph of free groups with  $\mathbb{Z}$  edges, which is not a free product of free and surface groups. Then  $G$  has abundant virtual homological torsion, that is,

for every finite abelian group  $A$ , there is a finite-index subgroup  $G_0 \leq G$  such that



$$G_0^{\text{ab}} = A \oplus \cdots.$$

## Homological torsion: By the book

---



## Homological torsion: By the book

---





## Homological torsion: By the book

---




## Homological torsion: By the book


---




## Homological torsion: By the book



## Homological torsion: By the book



## Homological torsion: By the book




## Homological torsion: Systems of surfaces

---


## Homological torsion: Systems of surfaces

---




## Homological torsion: Systems of surfaces

---




## Homological torsion: Systems of surfaces

---




## Homological torsion: Systems of surfaces


---




## Homological torsion: Systems of surfaces



## Homological torsion: Systems of surfaces




## Homological torsion: Systems of surfaces



In first homology:

## Homological torsion: Systems of surfaces




In first homology:

$$\sigma_1 + \sigma_4 = 0 \quad \sigma_2 + \sigma_3 = 0$$

$$\sigma_1 + \sigma_2 = 0 \quad \sigma_3 + \sigma_4 = 0$$

$$\sigma_1 + \sigma_3 = 0 \quad \sigma_2 + \sigma_4 = 0$$

# Homological torsion: Systems of surfaces



In first homology:

$$\sigma_1 + \sigma_4 = 0 \quad \sigma_2 + \sigma_3 = 0$$

$$\sigma_1 + \sigma_2 = 0 \quad \sigma_3 + \sigma_4 = 0$$

$$\sigma_1 + \sigma_3 = 0 \quad \sigma_2 + \sigma_4 = 0$$

$$\Rightarrow$$


$$\sigma_1 = \sigma_2 = \sigma_3 = \sigma_4$$

$$2 \cdot \sigma_1 = 0$$

## Homological torsion: Two and three


---

## Homological torsion: Two and three



Enter a rigid vertex


---



# Enter a rigid vertex

Theorem (Wilton '18)

If  $G$  is not free then  $G$  has a surface subgroup.






# Enter a rigid vertex

Theorem (Wilton '18)

If  $G$  is not free then  $G$  has a surface subgroup.

B




X

# Enter a rigid vertex



## Theorem (Wilton '18)

If  $G$  is not free then  $G$  has a surface subgroup.

B



Just a map (not  $\pi_1$ -injective)



X

# Enter a rigid vertex


## Theorem (Wilton '18)

If  $G$  is not free then  $G$  has a surface subgroup.



## Theorem (Calegari '09)

Let  $F$  be a free group and let  $g \in [F, F]$ . Recall  
 $\text{cl}(g) = \min\{\ell \mid g = \text{product of } \ell \text{ commutators}\}$ .

B



Just a map (not  $\pi_1$ -injective)



X

# Enter a rigid vertex

## Theorem (Wilton '18)


If  $G$  is not free then  $G$  has a surface subgroup.

## Theorem (Calegari '09)

Let  $F$  be a free group and let  $g \in [F, F]$ . Recall  
 $\text{cl}(g) = \min\{\ell \mid g = \text{product of } \ell \text{ commutators}\}$ .

Then

$$\text{scl}(g) = \lim_{n \rightarrow \infty} \frac{\text{cl}(g^n)}{n} \in \mathbb{Q}.$$



# Enter a rigid vertex

## Theorem (Wilton '18)

If  $G$  is not free then  $G$  has a surface subgroup.


## Theorem (Calegari '09)

Let  $F$  be a free group and let  $g \in [F, F]$ . Recall  
 $\text{cl}(g) = \min\{\ell \mid g = \text{product of } \ell \text{ commutators}\}$ .  
Then



$$\text{scl}(g) = \lim_{n \rightarrow \infty} \frac{\text{cl}(g^n)}{n} \in \mathbb{Q}.$$

$\rightsquigarrow$  has a topological interpretation in terms of surfaces

B



Just a map (not  $\pi_1$ -injective)



X

# Enter a rigid vertex

## Theorem (Wilton '18)

If  $G$  is not free then  $G$  has a surface subgroup.




## Theorem (Calegari '09)

Let  $F$  be a free group and let  $g \in [F, F]$ . Recall  $\text{cl}(g) = \min\{\ell \mid g = \text{product of } \ell \text{ commutators}\}$ . Then

$$\text{scl}(g) = \lim_{n \rightarrow \infty} \frac{\text{cl}(g^n)}{n} \in \mathbb{Q}.$$

↷ has a topological interpretation in terms of surfaces

B



X

# Enter a rigid vertex

## Theorem (Wilton '18)

If  $G$  is not free then  $G$  has a surface subgroup.

## Theorem (Calegari '09)


Let  $F$  be a free group and let  $g \in [F, F]$ . Recall  $\text{cl}(g) = \min\{\ell \mid g = \text{product of } \ell \text{ commutators}\}$ . Then

$$\text{scl}(g) = \lim_{n \rightarrow \infty} \frac{\text{cl}(g^n)}{n} \in \mathbb{Q}.$$

⚠  $\text{Tor}(H_1)$



injective

B



↷ has a topological interpretation in terms of surfaces

$\widehat{X}$




X


# Enter a rigid vertex

Applicable to other settings?

e.g. graphs of graphs.



B



## Profinite rigidity 101


---

Let  $G$  be a finitely generated *residually finite* group (every  $g \in G$  survives in some finite quotient  $q : G \twoheadrightarrow Q$ ).

# Profinite rigidity 101

Let  $G$  be a finitely generated *residually finite* group (every  $g \in G$  survives in some finite quotient  $q : G \twoheadrightarrow Q$ ).


## Lattice/poset of finite-index normal subgroups of $G$



# Profinite rigidity 101

Let  $G$  be a finitely generated *residually finite* group (every  $g \in G$  survives in some finite quotient  $q : G \twoheadrightarrow Q$ ).


## Lattice/poset of finite-index normal subgroups of $G$



# Profinite rigidity 101

Let  $G$  be a finitely generated *residually finite* group (every  $g \in G$  survives in some finite quotient  $q : G \twoheadrightarrow Q$ ).


## Lattice/poset of finite-index normal subgroups of $G$



# Profinite rigidity 101

Let  $G$  be a finitely generated *residually finite* group (every  $g \in G$  survives in some finite quotient  $q : G \twoheadrightarrow Q$ ).

## Lattice/poset of finite-index normal subgroups of $G$



### Question

Which fg residually finite groups  $H$  can fill the empty  $\square$ 's?

# Profinite rigidity 101

---

## The basics

- Full lattice structure encoded in the set of finite quotients of  $G$  (“Dixon’s theorem”, DFPR ’82).

# Profinite rigidity 101

---

## The basics

- Full lattice structure encoded in the set of finite quotients of  $G$  (“Dixon’s theorem”, DFPR ’82).
- Finitely generated free abelian groups are *profinitely rigid*.

# Profinite rigidity 101

## The basics

- Full lattice structure encoded in the set of finite quotients of  $G$  (“Dixon’s theorem”, DFPR ’82).
- Finitely generated free abelian groups are *profinately rigid*.
- The lattice determines the abelianization of each  $\square$ :

If  $G$  and  $H$  have the same lattice, then

$$G_n^{\text{ab}} \cong H_n^{\text{ab}}.$$

# Profinite rigidity 101

## The basics

- Full lattice structure encoded in the set of finite quotients of  $G$  (“Dixon’s theorem”, DFPR ’82).
- Finitely generated free abelian groups are *profinately rigid*.
- The lattice determines the abelianization of each  $\square$ :

If  $G$  and  $H$  have the same lattice, then

$$G_n^{\text{ab}} \cong H_n^{\text{ab}}.$$

- Normality above is not required!

# Profinite rigidity 101

## The basics

- Full lattice structure encoded in the set of finite quotients of  $G$  (“Dixon’s theorem”, DFPR ’82).
- Finitely generated free abelian groups are *profinitely rigid*.
- The lattice determines the abelianization of each  $\square$ :

If  $G$  and  $H$  have the same lattice, then

$$G_n^{\text{ab}} \cong H_n^{\text{ab}}.$$

- Normality above is not required!

## Question (Remeslennikov)

Are finitely generated (non-abelian) free groups profinitely rigid?

# Profinite rigidity 201

---

# Profinite rigidity 201

---

The largest natural class of groups where Remeslennikov's question has an answer:

The largest natural class of groups where Remeslennikov's question has an answer:

## Theorem (Wilton, '18)

*Free groups are profinitely rigid amongst hyperbolic graphs of free groups with  $\mathbb{Z}$  edges.*

The largest natural class of groups where Remeslennikov's question has an answer:

## Theorem (Wilton, '18)

*Free groups are profinitely rigid amongst hyperbolic graphs of free groups with  $\mathbb{Z}$  edges.*

## Theorem (Ascari-F '25)

*Free products of free and surface groups are profinitely rigid amongst hyperbolic graphs of free groups with  $\mathbb{Z}$  edges.*

## Growth: non-normal towers

---

## Growth: non-normal towers

---

### Theorem (Ascari-F '25)

*There exists a tower of finite-index non-normal subgroups*

$$G \geq G_1 \geq G_2 \geq \cdots, \quad \text{with} \quad \bigcap_n G_n = \{1\},$$

*such that simultaneously for every prime  $p$ ,*

$$\lim_{n \rightarrow \infty} \frac{\log(|\mathrm{Tor}_p(G_n^{\mathrm{ab}}))|)}{[G : G_n]} > 0.$$

## Fractal-like towers

---


## Fractal-like towers

---




## Fractal-like towers

---




## Fractal-like towers

---




## Fractal-like towers


---




## Fractal-like towers



## Fractal-like towers



## Fractal-like towers



## Growth: normal towers

---

## Growth: normal towers

---

### Theorem (F '25)

*In books of  $\mathbb{Z}$ -bundles, homological torsion grows subexponentially along any exhausting tower of finite-index normal subgroups.*

## Growth: normal towers

---

No infinite amenable normal subgroups

**Theorem (F '25)**



*In books of I-bundles, homological torsion grows subexponentially along any exhausting tower of finite-index normal subgroups.*

## Growth: normal towers

No infinite amenable normal subgroups

No cheap rebuilding

**Theorem (F '25)**

*In books of  $\mathbb{I}$ -bundles, homological torsion grows subexponentially along any exhausting tower of finite-index normal subgroups.*

## Growth: normal towers

No infinite amenable normal subgroups

No cheap rebuilding

**Theorem (F '25)**

*In books of  $I$ -bundles, homological torsion grows subexponentially along any exhausting tower of finite-index normal subgroups.*

- **Tension:** Books of  $I$ -bundles can have abundant virtual homological torsion.

## Growth: normal towers

No infinite amenable normal subgroups

No cheap rebuilding

**Theorem (F '25)**

*In books of  $I$ -bundles, homological torsion grows subexponentially along any exhausting tower of finite-index normal subgroups.*

- **Tension:** Books of  $I$ -bundles can have abundant virtual homological torsion.
- Bergeron-Venkatesh: such behaviour expected in dimension  $\geq 4$ .


## A computation: unwrapping cylinders

---


## A computation: unwrapping cylinders

---


$$G_i = \langle a, b \rangle_{[a,b]^i = [c,d]^i} \langle c, d \rangle$$




# A computation: unwrapping cylinders




$$G_i = \langle a, b \rangle_{[a,b]^i = [c,d]^i} \langle c, d \rangle$$




# A computation: unwrapping cylinders



$$G_i = \langle a, b \rangle_{[a,b]^i = [c,d]^i} \langle c, d \rangle$$




# A computation: unwrapping cylinders



\*  $C_n$  measures the wrapping degree of the cylinder in the  $n^{\text{th}}$  cover.

$$G_i = \langle a, b \rangle_{[a,b]^i = [c,d]^i} \langle c, d \rangle$$



## Enter a rigid vertex: revisited

---

## Enter a rigid vertex: revisited

---

-   **Restrictions lifted:** Torsion no longer confined to cylinders.



## Enter a rigid vertex: revisited

---

-   **Restrictions lifted:** Torsion no longer confined to cylinders.
- Refining the construction ↗ torsion elements **inside** a rigid vertex. Multiply fast, difficult to control.

## Enter a rigid vertex: revisited

-  **Restrictions lifted:** Torsion no longer confined to cylinders.
- Refining the construction ↗ torsion elements **inside** a rigid vertex. Multiply fast, difficult to control.
- **Computations:**



$G_{i+1}$  obtained from  $G_i$  by adding a single letter to its presentation

# The landscape

---

# The landscape

$\text{Tor}(H_1)$	(Hyp) 3-manifolds	Surface groups	Books of $I$ -bundles	Rigid vertex
Abundance	(Sun '15)		(Ascari-F '25)	(Ascari-F '25)
Non-normal exponential growth	(Liu '19)		(Ascari-F '25)	(Ascari-F '25)
Normal growth rate	(Exp.)		Subexp. (F '25)	

# The landscape

$\text{Tor}(H_1)$	(Hyp) 3-manifolds	Surface groups	Books of $I$ -bundles	Rigid vertex
Abundance	(Sun '15)		(Ascari-F '25)	(Ascari-F '25)
Non-normal exponential growth	(Liu '19)		(Ascari-F '25)	(Ascari-F '25)
Normal growth rate	(Exp.)		Subexp. (F '25)	Subexp. (F, recently)

# The landscape

$\text{Tor}(H_1)$	(Hyp) 3-manifolds	Surface groups	Books of $I$ -bundles	Rigid vertex
Abundance	(Sun '15)		(Ascari-F '25)	(Ascari-F '25)
Non-normal exponential growth	(Liu '19)		(Ascari-F '25)	(Ascari-F '25)
Normal growth rate	(Exp.)		Subexp. (F '25)	Subexp. (F, recently)
"Cylinder decay" rate			$O(1/c_n)$	$O(\log(c_n)/c_n)$

## What's next?

---

## What's next?

---

- “Cylinder decay” rate:

matching lower bounds  $\rightsquigarrow$  applications to profinite rigidity

## What's next?

---

- “Cylinder decay” rate:

matching lower bounds  $\rightsquigarrow$  applications to profinite rigidity

- Graphs of graphs:

unwrapping cylinders  $\rightsquigarrow$  more topological complexity

# What's next?

---

- “Cylinder decay” rate:

matching lower bounds  $\rightsquigarrow$  applications to profinite rigidity

- Graphs of graphs:

unwrapping cylinders  $\rightsquigarrow$  more topological complexity

$\rightsquigarrow$  cross the threshold?



Thank you for listening!

