South Australian Paediatric Clinical Guidelines

Pruritic urticarial papules and plaques of pregnancy (PUPPP)

© Department of Health, Government of South Australia. All rights reserved.

Pruritic urticarial papules and plaques of pregnancy (PUPPP)

- Pruritic urticarial papules and plaques of pregnancy (PUPPP) is also known as polymorphic eruption of pregnancy, Bourne's toxaemic rash of pregnancy, toxaemic erythema of pregnancy, and late onset prurigo of pregnancy (Kroumpouzos and Cohen 2003)
- > PUPPP is the most common pregnancy related dermatoses and the rash usually resolves before or within several weeks of birth (Rapini 2004).
- > PUPPP is poorly defined because of its variable clinical presentation, lack of pathognomonic diagnostic features and lack of laboratory abnormalities.
- The main hypothesis is that rapid abdominal wall distention may cause damage to connective tissue in the striae with subsequent conversion of nonantigenic molecules to antigenic ones, thus triggering an inflammatory process (Kroumpouzos and Cohen 2003)
- Recurrence in subsequent pregnancies, with menses or oral contraceptives is uncommon

Clinical features

- Occurs predominantly in primigravidas late in the third trimester (exceptionally postpartum)
- Associated with multiple gestation (possibly due to excessive abdominal distention and higher progesterone levels)
- Lesions (pruritic papules) commonly begin in the abdominal striae with periumbilical sparing
- Erythematous urticarial papules and plaques may spread to the proximal thighs, buttocks and proximal arms
- > Small vesicles may occasionally be noted
- Itching is severe in 80 % of cases
- Usually does not affect the face, palms, or soles.
- Excoriations are rarely found

Incidence

- Between 1 in 130 pregnancies and 1 in 300 pregnancies
- More common in multiple pregnancy (7 in 200) (Elling et al. 2000)

Pathogenesis

- No immunologic or hormonal abnormalities have been found
- One recent study reported a decrease in serum cortisol (Vaughan Jones et al. 1999)
- > Fetal DNA has been identified in the skin of women with PUPPP, suggesting that microchimerism might be relevant in the pathogenesis of PUPPP



South Australia

UNKNOWN
South Australian Paediatric Clinical Guidelines Reference Committee. South
Australian Child Health Clinical Network
South Australian Paediatric Clinical Guidelines Reference Committee:
cywhs.paediatricclinicalguidelines@health.sa.gov.au

South Australian Paediatric Clinical Guidelines

Pruritic urticarial papules and plaques of pregnancy (PUPPP)

© Department of Health, Government of South Australia. All rights reserved.

- There is no diagnostic test for PUPPP
- Consider referral to infectious diseases consultant or dermatologist

Exclude the following:

- Obstetric cholestasis
- Contact dermatitis (allergic or irritant)
- Drug eruptions
- Erythema multiforme
- Herpes gestationis (immunofluorescence or viral culture)
- > Insect bites
- Scabies (skin scrapings)
- > Sea bather's eruption
- Urticaria (acute or chronic)

Treatment

- Symptomatic: Oral cetirizine [Zyrtec®] and topically betamethasone 0.02 % [Celestone M® cream] or similar
- Rarely, a short course of oral prednisolone may be necessary

References

- 1. Kroumpouzos G, Cohen LM. Specific dermatoses of pregnancy: An evidence-based systematic review. Am J Obstet Gynecol 2003; 188: 1083 92 (Level I)
- 2. Rapini RP. The skin and pregnancy. In: Creasy RK, Resnik R, Iams JD, editors. Maternal-fetal medicine. Principles and practice, 5th ed. Philadelphia: Saunders; 2004. p. 1204
- 3. Elling SV, McKenna P, Powell FC. Pruritic urticarial papules and plaques of pregnancy in twin and triplet pregnancies. J Eur Acad Dermatol Venereol 2000: 14: 378 81 (Level III-2)
- Vaughan Jones SA, Hern S, Nelson-Piercy C, Seed PT, Black MM. A prospective study of 200 women with dermatoses of pregnancy correlating clinical findings with hormonal and immunopathological profiles. Br J Dermatol 1999; 141: 71-81(Level III-2)
- 5. Ahmadi S, Powell FC. Pruritic urticarial papules and plaques of pregnancy: Current status. Australasian journal of dermatology 2005; 46: 53-60 (Level IV)

Version control and change history

PDS reference: OCE use only

Version	Date from	Date to	Amendment
1.0	19 June 04	21 April 09	Original version
2.0	21 April 09	Current	



Contact:

UNKNOWN

South Australian Paediatric Clinical Guidelines Reference Committee. South Australian Child Health Clinical Network

South Australian Paediatric Clinical Guidelines Reference Committee: cywhs.paediatricclinicalguidelines@health.sa.gov.au

