## South Australian Perinatal Practice Guidelines

# Cryptosporidiosis

© Department of Health, Government of South Australia. All rights reserved.

## Introduction

- Cryptosporidiosis (commonly called 'crypto') is a diarrhoeal disease caused by the microscopic parasite Cryptosporidium parvum
- > Cryptosporidium has a worldwide distribution
- > The parasite is protected by an outer shell that allows it to survive outside the body for long periods of time (it is very resistant to chlorine disinfection)

## Clinical symptoms

- > May be asymptomatic
- Cryptosporidium parvum gastroenteritis (in immunocompetent patients is usually self-limiting within 14 days and requires no treatment)) involves the following symptoms:
  - > Watery diarrhoea
  - >Abdominal cramps
  - >Slight fever
  - > Dehydration
- > In immunocompromised patients watery diarrhoea may be severe and prolonged

## Incubation period

> 1 – 12 days

## Route of transmission

- Cryptosporidium parvum can live in the intestine of animals or humans and is passed in the stool of an infected person or animal
- Direct contact (ingestion) with food, water or soil contaminated with cryptosporidium parasites e.g. swallowing recreational water in swimming pools, fountains, lakes, rivers or streams
- > Swimming in contaminated water
- Direct contact with surfaces contaminated with faeces of infected humans or animals

# Infection precautions

> Standard precautions

# Diagnosis

> Stool specimen for microscopy and sensitivity



ISBN number: Endorsed by: Contact: UNKNOWN
SA Maternal & Neonatal Clinical Network
South Australian Perinatal Practice Guidelines workgroup at:
cywhs.perinatalprotocol@health.sa.gov.au

## South Australian Perinatal Practice Guidelines

# Cryptosporidiosis

© Department of Health, Government of South Australia. All rights reserved.

### **Treatment**

- > There is no consistently effective treatment
- > Encourage fluids to avoid dehydration
- Anti-diarrhoeal medication should be avoided in pregnancy. However, codeine and dihydrocodeine are ADEC Category A medications and are therefore regarded as safe
- Nitazoxamide 500mg orally, every 12 hours for 3 days. ADEC Category B medication. (This drug is not registered in Australia; available via Special Access Scheme: Tel: 02-6232 8111)

### References

- Communicable Disease Control Branch. You've got what specific conditions cryptosporidiosis. Department of Health, Government of South Australia. Available from: URL: http://www.dh.sa.gov.au/pehs/Youve-got-what/ygw-cryptosporidiosis-feb07.pdf
- Centers for Disease Control and Prevention (CDC). Parasitic disease information.
   Cryptosporidiosis. Division of Parasitic Diseases. [cited 2008 December 16] [4 screens].
   Available from: URL: http://www.cdc.gov/ncidod/dpd/parasites

## Version control and change history

PDS reference: OCE use only

Version	Date from	Date to	Amendment
1.0	19 May 04	25 Jan 10	Original version
2.0	25 Jan 10	Current	Reviewed



ISBN number: Endorsed by: Contact: UNKNOWN
SA Maternal & Neonatal Clinical Network
South Australian Perinatal Practice Guidelines workgroup at:
cywhs.perinatalprotocol@health.sa.gov.au