

# spironolactone

## 5mg/mL oral mixture\*

© Department of Health, Government of South Australia. All rights reserved

### Note

This guideline provides advice of a general nature. This statewide guideline has been prepared to promote and facilitate standardisation and consistency of practice, using a multidisciplinary approach. The guideline is based on a review of published evidence and expert opinion.

Information in this statewide guideline is current at the time of publication.

SA Health does not accept responsibility for the quality or accuracy of material on websites linked from this site and does not sponsor, approve or endorse materials on such links.

Health practitioners in the South Australian public health sector are expected to review specific details of each patient and professionally assess the applicability of the relevant guideline to that clinical situation.

If for good clinical reasons, a decision is made to depart from the guideline, the responsible clinician must document in the patient's medical record, the decision made, by whom, and detailed reasons for the departure from the guideline.

This statewide guideline does not address all the elements of clinical practice and assumes that the individual clinicians are responsible for discussing care with consumers in an environment that is culturally appropriate and which enables respectful confidential discussion. This includes:

- The use of interpreter services where necessary,
- Advising consumers of their choice and ensuring informed consent is obtained,
- Providing care within scope of practice, meeting all legislative requirements and maintaining standards of professional conduct, and
- Documenting all care in accordance with mandatory and local requirements

## Dose and Indications

**Treatment of fluid overload states associated with hyperaldosteronism, including congestive heart failure and chronic lung disease (BPD); in combination with other diuretics**

### Oral

1mg/kg every twelve hours

Dose may be increased up to 2mg/kg if required

## Preparation and Administration

### Oral

The 5mg/mL solution contains:

Dose	2mg	4mg	6mg	8mg	10mg	12mg
Volume	0.4mL	0.8mL	1.2mL	1.6mL	2mL	2.4mL

\* 5mg/mL oral mixture is not commercially available however is manufactured at Women's & Children's Health Network Pharmacy and Flinders Medical Centre Pharmacy.

Give with feeds to minimise gastrointestinal irritation

**ISBN number:**  
**Endorsed by:**  
**Last Revised:**  
**Contact:**

000-0-00000-000-0  
South Australian Maternal & Neonatal Clinical Network  
8/11/2012  
South Australian Neonatal Medication Guidelines Workgroup at:  
NeoMed@health.sa.gov.au

# spironolactone

5mg/mL oral mixture\*

## Adverse Effects

### Common

Hyperkalaemia, hyponatraemia, hypochloraemia (especially when combined with thiazide diuretics), vomiting

### Infrequent

Gastrointestinal cramps, diarrhoea, drowsiness, gynaecomastia, mild acidosis, renal impairment

### Rare

Agranulocytosis, hepatotoxicity, rash, lichen planus, lupus-like syndrome, cutaneous vasculitis, urticaria, alopecia, chloasma, osteomalacia

## Monitoring

- > Electrolytes particularly potassium
- > Renal function
- > Periodic liver function tests and full blood counts if used long term

## Practice Points

- > As there is a high risk of hyperkalaemia in renal impairment, avoid if rapidly deteriorating or severe renal impairment

## Version control and change history

**PDS reference:** OCE use only

Version	Date from	Date to	Amendment
1.0	November 2012	current	Original version

**ISBN number:**  
**Endorsed by:**  
**Last Revised:**  
**Contact:**

978-1-74243-447-6  
South Australian Maternal & Neonatal Clinical Network  
8/11/2012  
South Australian Neonatal Medication Guidelines Workgroup at:  
NeoMed@health.sa.gov.au