### South Australian Perinatal Practice Guidelines

# Worms

© Department of Health, Government of South Australia. All rights reserved.

#### Introduction

- > Worldwide, several worm varieties exist that can infest the intestinal tract of people
- > The most common worm in Australia is Enterobius vermicularis (pinworm) and is more commonly seen in children
- Infection with other worms such as roundworms, hookworms and tapeworms is uncommon in Australia

## Enterobius vermicularis (pinworm)

- > Pinworms are about 10 mm by 0.7 mm and inhabit the terminal ileum, caecum, appendix, and proximal ascending colon of humans
- The female worm migrates to the rectum after copulation, and if not expelled during defecation, migrates to the perineum (usually at night), where an average of 11,000 eggs are released

# Clinical symptoms

- > Anal and vulval pruritus
- Restless sleeping due to pruritus
- Evidence of worm in faeces

## Route of transmission

- > Faecal oral contamination
- > Reinfection may occur by hand to oral transfer of eggs (after scratching) from the faeces / perineal region of an infected person
- Person to person transmission can also occur through handling of contaminated clothes or bed linen
- > Contact with surfaces that are contaminated with pinworm eggs (e.g. curtains, carpets)

# Incubation period

- After ingestion, eggs usually hatch in the duodenum within 6 hours. Worms mature at around two weeks and have a life span of approximately 2 months
- > Symptoms and infection are associated with high worm numbers, resulting from repeated infections

# Infection precautions

Standard precautions



ISBN number: Endorsed by: Contact:

# South Australian Perinatal Practice Guidelines

# Worms

© Department of Health, Government of South Australia. All rights reserved.

#### **Treatment**

#### Preventative measures

- > Encourage good hand washing
- > Change bed linen and underwear daily for several days after treatment. Normal hot washing of clothes and bed linen will kill pinworm eggs

#### Drug treatment

First line treatment

- > Pyrantel oral tablets in a single dose depending on weight (ADEC Category B2) Second line treatment
  - Mebendazole (ADEC Category B3)

### Reference

 Center for Disease Control (CDC). Pinworm infection. Fact sheet. Division of Parasitic Diseases. 1999 August [cited 2004 May 20]; [3 screens]. Available from: URL: http://www.cdc.gov/ncidod/dpd/parasites/pinworm/factsht\_pinworm.htm

#### Useful websites

E-Medicine – Enterobiasis. Available from URL: http://www.emedicine.com/ped/topic684.htm

South Australian Department of Health. You've got what – Worms. Available from URL: http://www.dh.sa.gov.au/pehs/Youve-got-what/ygw-worms.pdf

# Version control and change history

PDS reference: OCE use only

Version	Date from	Date to	Amendment
1.0	19 May 04	25 Jan 10	Original version
2.0	25 Jan 10	Current	



ISBN number: Endorsed by: Contact: UNKNOWN
SA Maternal & Neonatal Clinical Network
South Australian Perinatal Practice Guidelines workgroup at:
cywhs.perinatalprotocol@health.sa.gov.au