# Dermatophyosis Dematomycosis tinea (Ringworm)

© Department of Health, Government of South Australia. All rights reserved.

# Ringworm

- > Ringworm is a contagious fungus infection that can affect the scalp, the body, the feet (athlete's foot), or the nails
- > The fungi that causes ringworm of the scalp lives in humans and animals
- > The fungi that causes ringworm of the body lives in humans, animals and soil
- > The fungi that causes ringworm of the foot and nails live only in humans
- > The name ringworm comes from the characteristic red ring that can appear on an infected person's skin

## Clinical features

#### Scalp

- Begins as a small pimple that becomes larger, leaving scaly patches of temporary baldness
- > Infected hairs may become brittle and break off easily

#### **Body**

- > Flat, round patch anywhere on the skin except the for the scalp and feet
- > The groin is a common area of infection
- As the rash gradually expands, its centre clears to produce a ring
- > May be itchy

#### **Feet**

> Appears as scaling or cracking of the skin, especially between the toes

#### **Nails**

> The nails become thicker, discoloured and brittle, or become chalky and disintegrate

## Diagnosis

- Microscopic examination of scrapings taken from skin lesions
- > Examination of scalp lesions under ultraviolet light

#### Mode of transmission

- > Direct skin to skin contact with an infected person or pet
- Indirect contact with an object or surface that an infected person or pet has contacted



# Dermatophyosis Dematomycosis tinea (Ringworm)

© Department of Health, Government of South Australia. All rights reserved.

## Incubation period

- > One to three weeks
- Varies with the site of infection

## Infection precautions

- > Additional precautions
- Minimize close contact with others until treated
- > Ensure that the person or animal that was the source of infection is treated

## **Treatment**

- Nail hair and widespread dermatophytosis infections are best treated with antifungal drugs
- Serise of ulvin should be avoided during pregnancy
- Use topical treatments e.g. clotrimazole (Canesten®)

#### Reference

1. Ellis D, Watson A. Systemic antifungal agents for cutaneous fungal infections. Australian prescriber [serial online] 1996 [cited 2004 Jun 8]; 19: [5 screens]. Available from URL:

http://www.australianprescriber.com/magazine/19/3/72/5/

#### Other useful web sites

http://www.health.sa.gov.au/PEHS/Youve-got-what/ygw-fungal-skin-nails.pdf

## Version control and change history

PDS reference: OCE use only

Version	Date from	Date to	Amendment
1.0	19 June 04	21 April 09	Original version
2.0	21 April 09	current	

