

South Australian Neonatal Medication Guidelines

phenoxymethylpenicillin

oral mixture

© Department of Health, Government of South Australia. All rights reserved

Note

This guideline provides advice of a general nature. This statewide guideline has been prepared to promote and facilitate standardisation and consistency of practice, using a multidisciplinary approach. The guideline is based on a review of published evidence and expert opinion.

Information in this statewide guideline is current at the time of publication.

SA Health does not accept responsibility for the quality or accuracy of material on websites linked from this site and does not sponsor, approve or endorse materials on such links.

Health practitioners in the South Australian public health sector are expected to review specific details of each patient and professionally assess the applicability of the relevant guideline to that clinical situation.

If for good clinical reasons, a decision is made to depart from the guideline, the responsible clinician must document in the patient's medical record, the decision made, by whom, and detailed reasons for the departure from the guideline.

This statewide guideline does not address all the elements of clinical practice and assumes that the individual clinicians are responsible for discussing care with consumers in an environment that is culturally appropriate and which enables respectful confidential discussion. This includes:

- The use of interpreter services where necessary,
- Advising consumers of their choice and ensuring informed consent is obtained,
- Providing care within scope of practice, meeting all legislative requirements and maintaining standards of professional conduct, and
- Documenting all care in accordance with mandatory and local requirements

Synonyms

Penicillin V

Dose and Indications

Infection due to susceptible organisms

Oral

10mg/kg/dose given four times a day

Length of treatment should be guided by pathology and clinical picture.

Preparation and Administration

Oral

There are various strengths available, refer to local guidelines for the specific strength available at your institution or unit and product information for storage conditions.

Give preferably on an empty stomach to optimise absorption

ISBN number:
Endorsed by:
Last Revised:
Contact:

978-1-74243-423-0
South Australian Maternal & Neonatal Clinical Network
8/11/2012
South Australian Neonatal Medication Guidelines Workgroup at:
NeoMed@health.sa.gov.au

Adverse Effects

Common

Diarrhoea

Infrequent

Vomiting, [Clostridium difficile-associated disease](#)

Rare

Black tongue, electrolyte disturbances, neurotoxicity, bleeding, blood dyscrasias

Anaphylactic shock is not commonly seen in the neonates.

Version control and change history

PDS reference: OCE use only

Version	Date from	Date to	Amendment
1.0	November 2012	current	Original version

ISBN number:
Endorsed by:
Last Revised:
Contact:

978-1-74243-423-0
South Australian Maternal & Neonatal Clinical Network
8/11/2012
South Australian Neonatal Medication Guidelines Workgroup at:
NeoMed@health.sa.gov.au