


Chapter 6 Confidence Intervals

STA2023 SECTION 6.1 Confidence Intervals for the Mean (σ Known) NOTES

Learning Outcomes:

- 1) Find a point estimate and a margin of error
- 2) Construct and interpret confidence intervals for the population mean when σ is known
- 3) Determine the minimum sample size required when estimating a population mean



Inferential statistics are then used to generalize and predictions about the parameter based on the statistic gained from the sample.

Point Estimate – a point or single value used to approximate a population parameter (population mean μ or population proportion p). The sample mean \bar{x} is the best point estimate for the population mean μ and the sample proportion \hat{p} is the best point estimate for the population proportion p .


*Point estimates (sample values) are a good approximation to population values, but are not exact (sampling error), so we give a range or interval of values instead.

Confidence Interval or Interval Estimate – a range or interval of values used to estimate the true value of a population parameter.

Confidence Level – a percentage (0%-100%) that measures the success rate of the method used to construct the confidence interval. If we draw many samples and use each one to construct a confidence interval, in the long run, the percentage of confidence intervals that cover the true value of the population would be equal to the confidence level.

Let $c = \text{confidence level}$ and let $\alpha = 1 - c$. Therefore c and α are complements. Confidence levels that are most commonly used are $c = 90\%$ or $c = 95\%$ or $c = 99\%$.

Let $c = 90\%$.




Let $c = 95\%$.

Let $c = 99\%$.

$$Z_{0.05} = \text{invNorm}(0.05, 0, 1) = -1.6449$$

$$Z_{0.95} = 1.6449$$

$$C = 99\%$$



$$Z_{0.995} = \text{inv Norm} \left(2.5758 \right)$$

$$= 2.5758$$

$$Z_{0.005} = -2.5758$$

$$Z_{\frac{\alpha}{2}}$$

$$Z_{1 - \frac{\alpha}{2}}$$

critical value

$CI \rightarrow 90\%, 95\%, 99\% \Rightarrow C$
 \downarrow for pop. parameter $\alpha = 1 - C$

6.1

CI for pop mean

Z interval (σ is known)

6.2

CI for pop mean

T interval (σ is unknown)

6.3

CI for pop proportion

"1-prop. Z interval"
 [Sample prop. will be given as % or x and n format.]

point estimate (PE)
 $= \bar{x}$

PE is \bar{x}

PE is $\hat{p} = \frac{x}{n}$

Critical value is
 $Z_{\frac{\alpha}{2}}$ (lower critical value)
 and $Z_{1-\frac{\alpha}{2}}$ (upper)

Critical value is
 $t_{\frac{\alpha}{2}}$ (lower),

$t_{1-\frac{\alpha}{2}}$ (upper)

$t_{\frac{\alpha}{2}} = \text{invT}(\frac{\alpha}{2}, df)$

$df = n - 1$

Critical value
 $Z_{\frac{\alpha}{2}}$ (lower)
 and $Z_{1-\frac{\alpha}{2}}$ (upper)

$Z_{\frac{\alpha}{2}} = \text{invNorm}(\frac{\alpha}{2}, 0, 1)$

6.1

Standard error (SE)

$$SE = \frac{\sigma}{\sqrt{n}}$$

6.2

SE is

$$SE = \frac{s}{\sqrt{n}}$$

6.3

SE is

$$SE = \sqrt{\frac{\hat{p}\hat{q}}{n}}$$

$$\hat{p} = \frac{x}{n}$$

$$\hat{q} = 1 - \hat{p}$$

n = Sample size

Margin of error (ME)

$$ME = SE \times \text{critical value}$$

$$= \frac{\sigma}{\sqrt{n}} \times Z_{1-\frac{\alpha}{2}}$$

$$ME = SE \times \text{critical value}$$

$$ME = \frac{s}{\sqrt{n}} \times t_{1-\frac{\alpha}{2}}$$

$$ME = SE \times$$

critical value

$$ME = \sqrt{\frac{\hat{p}\hat{q}}{n}}$$

$$\times Z_{1-\frac{\alpha}{2}}$$

CI is

$$(PE - ME, PE + ME)$$

CI is

$$(PE - ME, PE + ME)$$

CI is

$$(PE - ME, PE + ME)$$

Chapter 6 Confidence Intervals

$z_{\frac{\alpha}{2}}$

is called the critical value of z . It represents –

Margin of Error – the maximum likely difference between the observed sample value and the true population parameter. (maximum likely difference between \bar{x} and μ or between \hat{p} and p).

CONSTRUCTING A CONFIDENCE INTERVAL ESTIMATE FOR THE POPULATION MEAN μ

For means, we are using sample means from a sample of size n – The Central Limit Theorem says we can use the normal distribution if _____ and _____. The standard error of the sampling distribution will be :


And therefore, the Margin of Error will be:

To construct a confidence interval estimate: *point estimate* \pm *margin of error*

Example #1: In a simple random sample of 150 Fort Myers households, the sample mean number of personal computers was $\bar{x} = 1.32$. Assume that the population standard deviation is $\sigma = 0.41$.

- a. Construct a 95% confidence interval estimate for the mean number of personal computers for all households in Fort Myers.

$$n = 150, \quad \sigma = 0.41, \quad \bar{x} = 1.32$$
$$C = 95\% = 0.95, \quad \alpha = 1 - C = 0.05$$



$$\checkmark SE = \frac{\sigma}{\sqrt{n}} = \frac{.41}{\sqrt{150}} = .0335$$

$$\checkmark ME = SE \times \text{critical value} = 0.0335 \times 1.9599 \\ = 0.0657$$

\checkmark 95% CI is

$$\begin{aligned} & (PE - ME, PE + ME) \\ & = (1.32 - 0.0657, 1.32 + 0.0657) \\ & = (1.2543, 1.3857) \quad \checkmark \end{aligned}$$

We are 95% confident that
our mean # of PC in all
households in Fort Myers is between
(1.2543, 1.3857) \checkmark

σ known \rightarrow Z-interval

$$C = 90\% \quad , \quad \alpha = 1 - C = 10\% = 0.10$$


$$= 0.90$$

critical value $\boxed{Z_{1-\frac{\alpha}{2}}} > Z_{\frac{\alpha}{2}}$

$$Z_{1-\frac{\alpha}{2}} = \text{invNorm}\left(1-\frac{\alpha}{2}, 0, 1\right)$$

$$SE = \frac{\sigma}{\sqrt{n}}$$

$$ME = Z_{1-\frac{\alpha}{2}} \times \sqrt{n}$$



$$\Rightarrow 25 = 1.9599 \times \frac{210}{\sqrt{n}}$$

$$\Rightarrow \frac{25}{1.9599 \times 210} = \frac{1}{\sqrt{n}}$$

$$\Rightarrow 0.0607 = \frac{1}{\sqrt{n}}$$

$$\Rightarrow (\sqrt{n})^2 = \frac{1}{0.0607} = (16.46316)^2$$

$$\Rightarrow n = 271.0356 \approx \boxed{272}$$

$$\boxed{Z_{0.975}}$$

$$= \text{invNorm}(0.975, 0, 1)$$

$$= 1.9599$$

Chapter 6 Confidence Intervals

Example 2: You wish to estimate the mean amount of time that internet users spend on Facebook each month. How many internet users must be surveyed in order to be 95% confident that your sample mean is within 25 minutes of the true population mean? From a prior *Nielson* survey, assume the standard deviation of the population is ~210 minutes.

$$\sigma = 210, \quad C = 0.95, \quad ME = 25$$

$$n = 272$$

Example 3: An admissions director wants to estimate the mean age of all students enrolled at a college. The estimate must be within 1.5 years of the population mean. Assume the population of ages is normally distributed and $\sigma = 2.7$ years. Determine the minimum sample size required to estimate the population mean with 95% confidence.

$$C = 0.95, \quad \sigma = 2.7, \quad ME = 1.5$$

$$ME = Z_{0.975} \cdot \frac{\sigma}{\sqrt{n}}$$

$$\Rightarrow 1.5 = 1.9599 \cdot \frac{2.7}{\sqrt{n}}$$

$$\Rightarrow \frac{1.5}{(1.9599) \cdot (2.7)} = \frac{1}{\sqrt{n}}$$

$$\Rightarrow 0.2835 = \frac{1}{\sqrt{n}}$$

$$\Rightarrow \sqrt{n} = \frac{1}{0.2835} = 3.5278$$


$$n = (3.5278)^2 = 12.45$$

Example 4: An admissions director wants to estimate the mean age of all students enrolled at a college. The estimate must be within 2 years of the population mean. Assume the population of ages is normally distributed and $\sigma = 1.3$ years. Determine the minimum sample size required to estimate the population mean with 90% confidence.

$$\approx 13$$

$$\alpha = 5\%$$


Normal dist



$$0.95 + 0.025 = 0.975$$

$$z_{0.975} = \text{invNorm}(0.975, 0, 1)$$

$$= 1.9599$$



AL = subscript

$$1 - 0.005$$

Chapter 6 Confidence Intervals

STA 2023 SECTION 6.2 Confidence Intervals for the Mean (σ Unknown) NOTES

T-interval

Learning Outcomes:


- Interpret the t -distribution
- Construct and interpret confidence intervals for a population mean when σ is not known

When trying to estimate the value of the population mean μ , if the value of the population standard deviation σ is known and $n > 30$, then we can use the standard normal distribution to build confidence intervals.

If a value of σ is not known (which is usually the case in research situations), then we must use the sample standard deviation (s), which does not always follow a normal distribution. So, instead of the normal distribution, we use another distribution called the t-distribution.

Properties of the t-distribution:

- The graph of the distribution **changes for different sample sizes**. As n gets larger, the shape of the graph approaches that of the normal distribution.
- The graph of the distribution is a somewhat bell-shaped symmetric curve with a mean of $t = 0$.
- The graph reflects greater variability that is expected with smaller samples, therefore the standard deviation is greater than 1 and **changes depending on sample size**.




Degrees of Freedom (df) – For a collection of sample data, the degrees of freedom is the number of sample values that can vary after a certain restriction has been imposed on all data values. The degrees of freedom is always one less than the sample size. Therefore **$df = n - 1$** .

Finding Critical Values of t:

*Note:

Suppose $n = 15$ and we want 95% confidence:

$$C = 0.95, \alpha = 0.05$$



Chapter 6 Confidence Intervals

Constructing a Confidence Interval Estimate for a Population Mean μ ; σ is NOT known

$$C = .95, .90, .99$$
$$t_{1-\frac{\alpha}{2}} =$$
$$\alpha = 1 - C$$
$$SE = \frac{s}{\sqrt{n}}$$

$$ME = t_{1-\frac{\alpha}{2}} \cdot \frac{s}{\sqrt{n}}$$

Example 1: The University of Arizona (Phoenix) conducted The Garbage Project. For 16 randomly chosen families, the mean weight of plastics discarded in the trash (that could be recycled) was $\bar{x} = 1.911$ pounds with standard deviation of $s = 0.892$ pounds. Construct a 90% confidence interval estimate for the true mean amount of plastics discarded in the trash for all families in Phoenix.

→ $C = 0.90, s = 0.892, \bar{x} = 1.911$

$$90\% \text{ CI} \Rightarrow (1.5201, 2.3019)$$

90% confident that μ is betⁿ (1.5201, 2.3019)

Example 2: In a random sample of 10 people, the mean commute time to work was 40 minutes with a standard deviation of 6.5 minutes. Find a point estimate for the population mean. Assume the population is normally distributed.

a) Find a point estimate for the population mean commute time.

b) Construct and interpret a 95% confidence interval to estimate the population mean. Do not round.