

THE EQUIVARIANT UNIFORM KAN FIBRATION MODEL OF CUBICAL HOMOTOPY TYPE THEORY (E. RIEHL)

Notes by Jon Sterling

August 12, 2019

(0.1) Joint work with Awodey, Caallo, Coquand, Sattler

(0.2) We want a $\mathbf{P}(\square)$ -based model of HoTT which is a Quillen model category equivalent to spaces (Kan complexes in simplicial sets). We may further ask that this equivalence be a *nice* functor (for instance, a triangulation), and that \square supports inductive constructions (in the sense of being an Eilenberg-Zilber category).

Team	Cubes	Equivalent to spaces?
BCH	symmetric monoidal cubes	No: consider \mathbb{I}^2/swap (Buchholtz)
ABCFHL, A	Cartesian cubes	No: analogous argument by Sattler
CCHM	De Morgan cubes	No: consider \mathbb{I}/rev (Buchholtz)
CCHM	Dedekind	open problem

(0.3) We will work with Cartesian cubes, but change the notion of fibration to rule out the counterexample. We need $* \rightarrow \mathbb{I}^2/\text{swap}$ to be a trivial cofibration. The idea is to add an equivariance condition $j(x, y)\sigma = j(x, y\sigma)$ relative to symmetries σ :

$$\begin{array}{ccccccc}
 * & \longrightarrow & * & \longrightarrow & * & \xrightarrow{x} & X \\
 \downarrow & & \downarrow & & \downarrow & \nearrow j(x, y\sigma) & \nearrow \\
 \mathbb{I}^2 & \xrightarrow{\sigma} & \mathbb{I}^2 & \xrightarrow{e} & \mathbb{I}^2/\text{swap} & \xrightarrow{y} & Y \\
 & & & & \downarrow j(x, y) & & \\
 & & & & & &
 \end{array}$$

We get the desired lift using the universal property of the quotient map e .