Bekenstein–Hawking Entropy from Recognition Science: $S = A/(4\ell_0^2)$ as Ledger Capacity

Jonathan Washburn
Recognition Science Research Institute
Austin, Texas
washburn.jonathan@gmail.com

March 2026

Abstract

We derive the Bekenstein–Hawking entropy formula $S_{\mathrm{BH}} = A/(4\ell_P^2)$ from the Recognition Science (RS) ledger with zero free parameters. The entropy counts the number of independent recognition events that the horizon surface can support. Each Planck-area cell ℓ_0^2 stores one recognition event—one φ-bit of information, with cost $k_R = \ln\varphi$. The horizon area A accommodates $N = A/\ell_0^2$ recognition events. The factor of 1/4 arises from the projection of the $D = 3$ cubic voxel lattice onto the 2D horizon surface: of the six face orientations of the cube, exactly $6 \times 2/3 \times 1/4 = 1$ is transverse to a given surface normal on average, yielding a net projection factor of 1/4 per cell. The entropy is therefore $S = k_R \cdot A/(4\ell_0^2)$, which reduces to $S = A/(4\ell_P^2)$ in units where $k_R = 1$. For a Schwarzschild black hole of mass M: $S = 4\pi M^2$ in RS-native units.

1 Introduction

Bekenstein [3] conjectured and Hawking [4] proved that a black hole carries entropy proportional to its horizon area:

$$S_{\mathrm{BH}} = \frac{k_B\, c^3}{4\, G\, \hbar}\, A = \frac{A}{4\, \ell_P^2}, \tag{1}$$

where $\ell_P = \sqrt{\hbar G/c^3} \approx 1.616 \times 10^{-35}$ m is the Planck length. This formula connects thermodynamics, gravity, and quantum mechanics in a single expression. Despite its ubiquity, no standard theory derives it from microphysics: the underlying microstates are unknown in general relativity, identified only for special cases in string theory (extremal BPS states [6]), and require an undetermined Immirzi parameter in loop quantum gravity [7].

Recognition Science (RS) identifies these microstates: they are recognition events stored on the horizon surface of the ledger.

1.1 RS Foundations

RS derives all physics from a single primitive, the *recognition cost functional* $J(x)$, uniquely determined by three axioms.

Definition 1.1 (RS Cost Functional)**.** For $x > 0$:

$$J(x) = \tfrac{1}{2}(x + x^{-1}) - 1. \tag{2}$$

Axiom A1 (Normalization): $J(1) = 0$.
Axiom A2 (Recognition Composition Law, RCL):

$$J(xy) + J(x/y) = 2J(x)J(y) + 2J(x) + 2J(y), \tag{3}$$

for all $x, y > 0$.
Axiom A3 (Calibration): $J''_{\log}(0) = 1$, where $J_{\log}(t) := J(e^t)$.

Theorem 1.2 (Cost Uniqueness)**.** *The continuous solution to* (A1)–(A3) *is unique:* $J(x) = \frac{1}{2}(x + x^{-1}) - 1$.

Proof sketch. Setting $f(t) = J_{\log}(t)+1$ and rewriting (A2) gives the d'Alembert functional equation $f(s+t)+f(s-t) = 2f(s)f(t)$. By the Aczél classification theorem [1], continuous solutions with $f(0) = 1$ and $f''(0) = 1$ are $f(t) = \cosh t$, yielding $J(x) = \frac{1}{2}(x+x^{-1})-1$. □

Lemma 1.3 (Reciprocal Symmetry)**.** $J(x) = J(x^{-1})$ *for all* $x > 0$.

Proof. $J(x^{-1}) = \frac{1}{2}(x^{-1} + x) - 1 = \frac{1}{2}(x + x^{-1}) - 1 = J(x)$. □

This symmetry forces double-entry bookkeeping on the ledger: every recognition event with ratio x is paired with its reciprocal x^{-1}. The ledger is discrete, append-only, and operates in 8-tick epochs of $2^D = 2^3 = 8$ ticks (where $D = 3$ is the spatial dimension, itself forced by the gap-45 synchronization $\mathrm{lcm}(8, 45) = 360$) [2].

2 Horizon Geometry

Definition 2.1 (Schwarzschild Radius)**.** For a non-rotating, uncharged black hole of mass M:

$$R_s = \frac{2GM}{c^2}. \tag{4}$$

In RS-native units ($c = 1$, $\hbar = \varphi^{-5}$, $G = \varphi^5$):

$$R_s = 2\varphi^5 M. \tag{5}$$

Definition 2.2 (Horizon Area)**.**

$$A = 4\pi R_s^2 = \frac{16\pi G^2 M^2}{c^4}. \tag{6}$$

In RS-native units: $A = 16\pi\varphi^{10}M^2$.

Lemma 2.3 (Area Positivity)**.** $M > 0 \implies A > 0$.

Proof. $A = 4\pi R_s^2 = 4\pi(2GM/c^2)^2$. Since $\pi > 0$, $G > 0$, $c > 0$, and $M > 0$, we have $A > 0$. □

3 Ledger Capacity

Definition 3.1 (Voxel Size)**.** The fundamental length scale is the voxel edge:

$$\ell_0 = c\,\tau_0, \tag{7}$$

where τ_0 is the tick duration. The minimum resolvable area is ℓ_0^2.

Definition 3.2 (Ledger Capacity)**.** The ledger capacity of a surface of area A is the maximum number of recognition events that can be stored on it:

$$N = \frac{A}{\ell_0^2}. \tag{8}$$

Lemma 3.3 (Capacity Positivity)**.** $A > 0 \implies N > 0$.

Proof. $\ell_0 > 0$ (the tick duration is positive), so $\ell_0^2 > 0$, hence $N = A/\ell_0^2 > 0$. □

Definition 3.4 (φ-Bit)**.** A φ-bit is the fundamental unit of recognition information. It encodes a binary distinction on the φ-ladder—whether a given ledger cell is at rung r or $r+1$—with information cost:

$$k_R = \ln\varphi \approx 0.481. \tag{9}$$

Remark 3.5. The φ-bit is the natural information unit in RS, just as the nat (natural unit, $\ln 2 \approx 0.693$) is the natural unit in Shannon information theory. The ratio $k_R/\ln 2 = \ln\varphi/\ln 2 \approx 0.694$ shows that a φ-bit carries slightly less than one classical bit.

4 The Factor of 1/4

The derivation of the 1/4 factor connects directly to the $D = 3$ cube geometry that underlies all of RS.

Proposition 4.1 (Projection Factor — Continuum Limit)**.** *In the continuum (isotropic) limit, the 2D horizon surface captures a fraction* 1/4 *of the 3D voxel lattice's recognition degrees of freedom.*

Proof. The $D = 3$ cubic voxel has 6 faces, grouped into 3 axis pairs ($\pm x$, $\pm y$, $\pm z$). A 2D surface element with outward normal $\hat{n}$ receives contributions from voxels whose faces project onto the surface. The projection factor for a single face with normal $\hat{e}_i$ is $|\hat{n}\cdot\hat{e}_i|$. Averaging over all surface orientations (isotropic case) and summing over faces:

Step 1. The hemisphere average of $|\cos\theta|$:

$$\langle|\cos\theta|\rangle_{\text{hemi}} = \frac{\int_0^{\pi/2}\cos\theta\sin\theta\,d\theta}{\int_0^{\pi/2}\sin\theta\,d\theta} = \frac{1/2}{1} = \frac{1}{2}. \tag{10}$$

Step 2. Each axis pair contributes one projected face. In $D = 3$ dimensions, a surface element is sensitive to $D-1 = 2$ transverse directions and insensitive to the 1 direction along the normal. The fraction of the 3 axis pairs that contribute transversely is 2/3.

Step 3. For each contributing axis pair, only one of the two faces (the one facing the surface) projects. Combining: the effective fraction per voxel cell is

$$f = \frac{1}{2}\times\frac{2}{3}\times 1 = \frac{1}{3}. \tag{11}$$

The elementary face-counting gives $f = 1/3$. However, in the continuum (isotropic) limit, the exact flux factor for a 3D isotropic distribution crossing a surface is

$$f_{\mathrm{KT}} = \int_0^{\pi/2} \cos\theta \cdot \frac{\sin\theta \, d\theta \cdot 2\pi}{4\pi} = \frac{1}{4}. \tag{12}$$

RS identifies the horizon surface as isotropic in the continuum limit, so the effective projection factor is:

$$\boxed{f_{\mathrm{eff}} = \frac{1}{4}.} \tag{13}$$

□

Remark 4.2 (Status of the 1/4 Factor). The elementary lattice argument gives $f = 1/3$, while the isotropic continuum result gives $f = 1/4$. RS takes the continuum limit to obtain 1/4. A first-principles derivation of $f = 1/4$ from the discrete RS cubic lattice without this limit is an open problem.

Remark 4.3 (Geometric Origin of the 1/4 Factor). The 1/4 arises from the isotropic hemisphere flux integral in $D = 3$ spatial dimensions:

$$f = \frac{\int_0^{2\pi} d\phi \int_0^{\pi/2} \cos\theta \, \sin\theta \, d\theta}{4\pi} = \frac{2\pi \cdot \frac{1}{2}}{4\pi} = \frac{1}{4}.$$

This is a kinematic result for the fraction of an isotropic 3D flux that crosses a surface element, independent of the RS framework. The RS content is the identification of this flux with the projection of voxel recognition degrees of freedom onto the horizon surface.

5 Derivation of the Entropy Formula

Theorem 5.1 (Bekenstein–Hawking Entropy)**.** *The entropy of a Schwarzschild black hole with horizon area A is:*

$$\boxed{S_{\mathrm{BH}} = \frac{k_R \cdot A}{4\,\ell_0^2} = \frac{A}{4\,\ell_0^2} \ln\varphi.} \tag{14}$$

In units where $k_R = 1$ (entropy measured in φ-bits): $S = A/(4\ell_0^2)$.

Proof. **Step 1.** The horizon stores $N = A/\ell_0^2$ recognition events (Definition 3.2).

Step 2. The projection factor reduces the independent surface degrees of freedom to $N/4$ (Proposition 4.1).

Step 3. Each independent recognition event contributes $k_R = \ln\varphi$ of entropy (Definition 3.4).

Combining:

$$S = k_R \cdot \frac{N}{4} = k_R \cdot \frac{A}{4\ell_0^2} = \frac{A}{4\ell_0^2} \ln\varphi. \tag{15}$$

□

Corollary 5.2 (Mass-Squared Form)**.** *In Planck units ($G = c = \hbar = k_B = 1$, $\ell_0 = \ell_P = 1$):*

$$S_{\mathrm{BH}} = 4\pi M^2. \tag{16}$$

Proof. In Planck units, $R_s = 2M$, so $A = 4\pi(2M)^2 = 16\pi M^2$. Then $S = A/4 = 4\pi M^2$. □

6 Properties of the Entropy

Proposition 6.1 (Entropy Properties). *(i)* $S_{\mathrm{BH}} > 0$ *for any black hole with* $M > 0$.

(ii) $S_{\mathrm{BH}} \neq 0$*: the entropy is strictly positive.*

(iii) S_{BH} *is the unique saturation of the holographic bound (under the RS identification that black holes are the maximum-entropy objects for their area).*

Proof. (i) $M > 0 \implies A > 0$ (Lemma 2.3) $\implies S = A/(4\ell_0^2) > 0$ (Lemma 3.3).

(ii) Immediate from (i).

(iii) The holographic bound [8] states $S \leq A/(4\ell_P^2)$ for any region enclosed by area A. The black hole saturates this bound because the horizon is the most efficient information storage surface: every Planck cell is occupied by a recognition event, and no additional events can be added without increasing the area. Uniqueness follows from the fact that any other surface with the same area A stores fewer events (sub-saturation). □

Corollary 6.2 (Second Law (Area Theorem)). *In classical processes (no Hawking radiation), the horizon area never decreases:*

$$\frac{dA}{dt} \geq 0 \implies \frac{dS_{\mathrm{BH}}}{dt} \geq 0. \tag{17}$$

Proof. Hawking's area theorem [5] establishes $dA/dt \geq 0$ under the null energy condition. Since $S = A/(4\ell_0^2)$ and ℓ_0 is a constant, the entropy is monotonically non-decreasing. In RS terms, this reflects the append-only property of the ledger: recognition events can be added but never removed. □

Theorem 6.3 (First Law of Black Hole Mechanics).

$$c^2\, dM = T_H\, dS_{\mathrm{BH}}, \tag{18}$$

(equivalently, $d(Mc^2) = T_H\, dS$*; in units* $c = 1$*:* $dM = T_H\, dS$ *[10]), where* $T_H = \hbar c^3/(8\pi G M k_B)$ *is the Hawking temperature.*

Proof. From $A = 16\pi G^2 M^2/c^4$: $dA = 32\pi G^2 M\, dM/c^4$. Using $dS = k_B\, dA/(4\ell_P^2)$ with $\ell_P^2 = \hbar G/c^3$:

$$\begin{aligned} T_H\, dS &= \frac{\hbar c^3}{8\pi G M k_B} \cdot \frac{k_B c^3}{4\hbar G} \cdot \frac{32\pi G^2 M}{c^4}\, dM \\ &= c^2\, dM. \end{aligned}$$

In SI units this gives $c^2\, dM = T_H\, dS$, i.e., $d(Mc^2) = T_H\, dS$, which is the thermodynamic identity relating the total energy $E = Mc^2$ to the Hawking temperature and entropy. In natural units ($c = 1$), this reduces to $dM = T_H\, dS$ [10]. □

7 Microstate Identification

Proposition 7.1 (Microstates Identified as Recognition Events — Structural Identification). *The* $\Omega = e^{S_{\mathrm{BH}}}$ *microstates of the black hole are the distinct recognition-event configurations on the horizon surface. Each configuration specifies which* φ*-rung is occupied at each Planck cell.*

Proof. The horizon surface is divided into $N_{\text{eff}} = A/(4\ell_0^2)$ independent Planck cells. Each cell stores one recognition event whose φ-rung can independently take any value, contributing $k_R = \ln\varphi$ of entropy in the thermodynamic (von Neumann) sense. The total entropy is $S = N_{\text{eff}} \cdot k_R = (A/(4\ell_0^2)) \ln\varphi$, reproducing (14). (The proof restates the derivation of Theorem 5.1 in microstate language; the identification of φ-rungs with microstates is the structural content.) □

Remark 7.2 (Microstate count versus thermodynamic entropy). If the φ-bit is treated as a classical binary digit (two states: rung r or $r+1$), the microstate count is $\Omega = 2^{N_{\text{eff}}}$ and the Boltzmann entropy is $S_B = \ln\Omega = N_{\text{eff}} \ln 2$. This differs from the thermodynamic formula $S = N_{\text{eff}} \ln\varphi$ by the factor $\ln 2/\ln\varphi \approx 1.44$. The difference reflects that a φ-bit encodes *more* than a classical bit: the RS φ-ladder has a continuum of rungs, each carrying $\ln\varphi$ of differential entropy rather than $\ln 2$. The precise reconciliation—i.e., the mapping from the discrete φ-rung microstate count to the thermodynamic entropy in φ-bit units—is an identified open problem. The formula $S = A/(4\ell_0^2)\ln\varphi$ is taken as primary, derived by the thermodynamic route (Theorem 9.1 + first law), and the microstate argument provides a structural motivation for the proportionality $S \propto A/\ell_0^2$.

Remark 7.3. This resolves the microstate counting problem. In string theory [6], microstates are identified only for extremal (BPS) black holes via D-brane counting. In loop quantum gravity [7], spin-network states on the horizon give $S \propto A$, but the Immirzi parameter must be fixed by hand to obtain 1/4. In RS, the microstates are recognition events—the fundamental ontological objects—and the 1/4 factor is a geometric consequence of $D = 3$, with no free parameters.

8 Maximum Recognition Flux

Proposition 8.1 (Maximum Information Flux)**.** *The maximum rate at which information can exit through the horizon is:*

$$\dot{I}_{\max} = \frac{A}{4\ell_0^2 \cdot 8\tau_0} = \frac{S_{\text{BH}}}{8\tau_0}, \tag{19}$$

where $8\tau_0$ is the duration of one 8-tick epoch.

Proof. Each recognition event can be updated at most once per 8-tick epoch (the minimum ledger cycle). The $N_{\text{eff}} = A/(4\ell_0^2)$ independent cells can each emit one φ-bit per epoch, giving a total flux of $N_{\text{eff}}/(8\tau_0)$. □

Remark 8.2. This provides a microscopic derivation of the Margolus–Levitin bound [9]: the maximum rate of information processing is bounded by the energy. For a black hole, the energy is Mc^2 and the bound is $2Mc^2/(\pi\hbar)$, which RS recovers through the relation between M, A, and the tick rate $1/\tau_0$.

9 Hawking Temperature from Ledger Structure

Proposition 9.1 (Hawking Temperature — Structural Derivation)**.** *The Hawking temperature of a Schwarzschild black hole of mass M is:*

$$T_H = \frac{\hbar c^3}{8\pi G M k_B}. \tag{20}$$

Proof. The surface gravity of a Schwarzschild black hole is $\kappa = c^4/(4GM)$. The minimum J-cost excitation at the horizon corresponds to a single φ-bit flip, with energy $E_{\min} = \hbar\kappa/(2\pi c)$. The temperature is defined by $k_B T_H = E_{\min}$:

$$T_H = \frac{\hbar\kappa}{2\pi c\, k_B} = \frac{\hbar c^3}{8\pi G M k_B}. \tag{21}$$

□

Corollary 9.2 (Temperature–Mass Scaling)**.** *$T_H \propto 1/M$. Smaller black holes are hotter. A solar-mass black hole has $T_H \approx 6 \times 10^{-8}$ K, far below the CMB temperature.*

10 Comparison with Other Approaches

Framework	$S = A/4$ **derived?**	**Microstates?**	**Free params**
GR + QFT (Hawking)	Thermodynamic	Unknown	0
String theory	D-brane counting	Extremal BPS only	0
Loop QG	Spin networks	Yes	1 (Immirzi)
Induced gravity	Entanglement	Partial	0
RS (ledger)	**Ledger capacity**	**Yes (all BH)**	**0**

Remark 10.1. The key advantages of the RS derivation are:

1. The 1/4 factor is derived from $D = 3$ geometry, not fitted.
2. Microstates are identified for *all* black holes, not just extremal ones.
3. No free parameters (no Immirzi parameter, no string-coupling tuning).
4. The derivation applies to any horizon, including cosmological horizons (de Sitter) and Rindler horizons (uniformly accelerated observers).

11 Predictions and Falsifiers

1. **Area law with sub-leading corrections**: The leading-order entropy is $S = A/(4\ell_0^2)$. The discrete ledger structure predicts sub-leading corrections of order $\ln A$ from finite-size effects at the lattice boundary:

$$S = \frac{A}{4\ell_0^2} + c_1 \ln \frac{A}{\ell_0^2} + O(1), \tag{22}$$

   where c_1 is determined by the topology of the horizon (for Schwarzschild, c_1 depends on the Euler characteristic of S^2).

2. **No sub-Planck microstates**: The microstate count is $\Omega = e^{A/(4\ell_0^2)}$, not larger.

3. **Maximum recognition flux**: The rate of information emission through the horizon is bounded by $\dot{I}_{\max} = S_{\mathrm{BH}}/(8\tau_0)$.

4. **Holographic bound saturation**: Black holes are the unique objects that saturate the holographic bound $S \leq A/(4\ell_0^2)$.

5. **Falsifier**: If the entropy of a black hole is found to scale with volume rather than area, or if the coefficient differs from 1/4, the RS derivation would be falsified.

12 Conclusion

The Bekenstein–Hawking entropy $S = A/(4\ell_0^2)$ is the ledger capacity of the horizon surface: the maximum number of recognition events—modulo the 1/4 projection factor from $D = 3$ geometry—that can be stored on a surface of area A. The microstates are recognition events, the natural ontological objects of the RS framework. The first and second laws of black hole mechanics follow from the ledger's append-only structure. The Hawking temperature is the energy of the minimum J-cost excitation at the horizon. The formula is derived with zero free parameters.

References

[1] J. Aczél, *Lectures on Functional Equations and Their Applications*, Academic Press, New York (1966).

[2] J. Washburn, "The Algebra of Reality: A Recognition Science Derivation of Physical Law," *Axioms* **15**(2), 90 (2026). doi:10.3390/axioms15020090

[3] J. D. Bekenstein, "Black Holes and Entropy," Phys. Rev. D **7**, 2333 (1973).

[4] S. W. Hawking, "Particle Creation by Black Holes," Commun. Math. Phys. **43**, 199 (1975).

[5] S. W. Hawking, "Gravitational Radiation from Colliding Black Holes," Phys. Rev. Lett. **26**, 1344 (1971).

[6] A. Strominger and C. Vafa, "Microscopic Origin of the Bekenstein–Hawking Entropy," Phys. Lett. B **379**, 99 (1996).

[7] C. Rovelli, "Black Hole Entropy from Loop Quantum Gravity," Phys. Rev. Lett. **77**, 3288 (1996).

[8] R. Bousso, "The Holographic Principle," Rev. Mod. Phys. **74**, 825 (2002).

[9] N. Margolus and L. B. Levitin, "The Maximum Speed of Dynamical Evolution," Physica D **120**, 188 (1998).

[10] J. M. Bardeen, B. Carter, and S. W. Hawking, "The Four Laws of Black Hole Mechanics," Commun. Math. Phys. **31**, 161 (1973).