Time, the Arrow, and the Past Hypothesis from a Single Cost Functional

Jonathan Washburn
Recognition Science Research Institute
Austin, Texas
washburn.jonathan@gmail.com

March 2026

Abstract

We show that a single cost functional $J(x) = \frac{1}{2}(x + x^{-1}) - 1$, uniquely determined by the Recognition Composition Law on $\mathbb{R}_{>0}$, forces three long-standing puzzles about time to become theorems rather than postulates. First, time is identified with the natural-number index of ledger updates, eliminating background temporality and yielding a minimal period of 8 ticks set by dimension forcing (2^D, $D = 3$). Second, the arrow of time is the direction of non-increasing defect along the tick sequence: recognition events reduce cost, and this reduction is provably irreversible. Third, the low-entropy initial condition (the "Past Hypothesis") is dissolved: the unique zero-cost configuration $\mathbf{1} = (1, \ldots, 1)$ is the sole global minimizer of total defect, and it serves as the asymptotic *future* attractor of RS dynamics; the Big Bang's low thermodynamic entropy corresponds to *high* RS defect, which is generic (not fine-tuned). No statistical reasoning, no fine-tuning, and no additional postulates beyond the cost axioms are required.

1 Introduction

The nature of time, its irreversible direction, and the special character of its boundary conditions constitute three of the deepest open problems in foundational physics.

The *nature of time* remains contested between the block-universe view, in which time is a static coordinate [3], and the process view, in which temporal passage is ontologically primitive [7]. Neither standpoint derives time from something simpler.

The *arrow of time*—the phrase coined by Eddington [8] for the empirical fact that macroscopic processes are irreversible—has been debated since Boltzmann [4]. Standard statistical mechanics accounts for it via the overwhelmingly large phase-space volume of high-entropy macrostates, but this argument presupposes a low-entropy initial condition [6, 2].

The *Past Hypothesis*—the postulate that the universe began in a state of extraordinarily low entropy—is widely regarded as the most puzzling boundary condition in cosmology. Penrose expressed the improbability as $1 : 10^{10^{123}}$ [6], and Carroll has emphasized that any fundamental theory should *derive* this condition rather than postulate it [5].

In this paper we show that all three problems have a common resolution within Recognition Science (RS) [1], a framework built on a single cost functional. The paper is organized as follows. Section 2 introduces the cost axioms and derives the unique cost functional J. Section 3 defines time as ledger tick count and establishes the 8-tick epoch structure. Section 4 proves the arrow of time via defect monotonicity and irreversibility of recognition. Section 5 proves the Past Theorem. Section 6 discusses implications and comparison with prior work.

2 The Cost Functional

We begin with three axioms that uniquely determine a cost functional on the positive reals. These axioms encode the minimal structure required for a self-consistent accounting framework: an identity element, a composition law, and a curvature normalization.

Axiom 1 (Normalization — A1)**.** $J(1) = 0$. The identity ratio has zero cost.

Axiom 2 (Recognition Composition Law — A2)**.** For all $x, y > 0$,

$$J(xy) + J(x/y) \;=\; 2\,J(x)\,J(y) + 2\,J(x) + 2\,J(y).$$

Axiom 3 (Calibration — A3)**.** $J''_{\log}(0) = 1$, where $J_{\log}(t) := J(e^t)$.

These axioms constitute a calibrated form of the classical d'Alembert (cosine) functional equation.

Theorem 2.1 (Cost Uniqueness — T5)**.** *The unique continuous function* $J : \mathbb{R}_{>0} \to \mathbb{R}$ *satisfying* A1–A3 *is*

$$J(x) \;=\; \frac{1}{2}\left(x + \frac{1}{x}\right) - 1. \tag{1}$$

Proof. Setting $y = 1$ in A2 and using A1 gives $J(x) + J(x) = 2\,J(x) \cdot 0 + 2\,J(x) + 0$, which is consistent for all J. The substitution $H(t) = 1 + J(e^t)$ transforms A2 into the d'Alembert equation $H(s+t) + H(s-t) = 2\,H(s)\,H(t)$, whose continuous solutions are $H(t) = \cosh(\lambda t)$. A3 fixes $\lambda = 1$, giving $J(e^t) = \cosh(t) - 1$. Substituting $x = e^t$ yields (1). For uniqueness, the d'Alembert equation on $\mathbb{R}$ with $H(0) = 1$ and $H''(0) = 1$ has $H = \cosh$ as its unique continuous even solution [9]. □

We record several immediate consequences.

Definition 2.2 (Defect)**.** For $x > 0$, the *defect* of x is $\mathrm{defect}(x) := J(x)$.

Lemma 2.3 (Non-negativity)**.** $J(x) \geq 0$ *for all* $x > 0$*, with equality if and only if* $x = 1$.

Proof. By the AM–GM inequality, $x + x^{-1} \geq 2$ for $x > 0$, so $J(x) = \frac{1}{2}(x + x^{-1}) - 1 \geq 0$. Equality holds iff $x = x^{-1}$, i.e., $x = 1$. □

Lemma 2.4 (Reciprocal Symmetry)**.** $J(x) = J(1/x)$ *for all* $x > 0$.

Proof. $\frac{1}{2}(x^{-1} + x) - 1 = \frac{1}{2}(x + x^{-1}) - 1$. □

Corollary 2.5 (Law of Existence)**.** *The unique zero-cost state is* $x = 1$. *We write* $\mathrm{RSExists}(x) \iff x > 0 \wedge J(x) = 0 \iff x = 1$.

Remark 2.6. The divergence $J(0^+) = +\infty$ is a theorem: as $x \to 0^+$, $x^{-1} \to \infty$, so $J(x) \to \infty$. This recovers the meta-principle "nothing has infinite cost" without postulating it.

3 Time as Tick Count

3.1 Discrete Temporality

In Recognition Science, time is not a background manifold parameter. Temporal progression is identified with the natural-number index of ledger updates.

Definition 3.1 (Tick)**.** A *tick* is a natural number $t \in \mathbb{N}$ labeling a ledger update event. The ordering of ticks is the standard ordering on $\mathbb{N}$.

Definition 3.2 (Ledger Snapshot)**.** A *ledger snapshot* at tick t is a pair (t, d) where $d \geq 0$ is the total defect of the ledger at that tick.

Theorem 3.3 (Discrete Time — F-004 Core)**.** *Time is discrete: the minimal temporal resolution is one tick. For any two ledger snapshots (t_1, d_1) and (t_2, d_2) with $t_1 < t_2$,*

$$t_2 - t_1 \;\geq\; 1.$$

No sub-tick dynamics exist.

Proof. Since $t_1, t_2 \in \mathbb{N}$ and $t_1 < t_2$, we have $t_2 \geq t_1 + 1$. □

3.2 The 8-Tick Epoch

The minimal period of a complete ledger update cycle is determined by dimension forcing.

Theorem 3.4 (Dimension Forcing — T8)**.** *The spatial dimension $D = 3$ is the unique RS-compatible dimension. Three independent arguments converge:*

(i) ***Linking:*** *Non-trivial topological linking of closed curves requires $D = 3$ (Alexander duality).*

(ii) ***Kepler stability:*** *Non-precessing near-circular orbits require $D = 3$ (Bertrand's theorem via the Binet equation).*

(iii) ***Gap synchronization:*** *The minimal period 2^D must synchronize with the gap-45 rung structure via $\mathrm{lcm}(2^D, 45)$; the minimum is achieved uniquely at $D = 3$, giving $\mathrm{lcm}(8, 45) = 360$.*

Definition 3.5 (Epoch)**.** An *epoch* is a complete cycle of $2^D = 2^3 = 8$ ticks. This is the minimal ledger-compatible walk on the D-dimensional hypercube (a Gray-code traversal of Q_3).

Theorem 3.6 (Epoch Length)**.** *The epoch length equals $8 = 2^3$ ticks.*

Remark 3.7. Continuous time, as it appears in classical and quantum mechanics, emerges as an effective approximation valid for large tick counts. The fundamental ontology is discrete.

4 The Arrow of Time

4.1 Defect Monotonicity

The direction of time is not imposed by fiat or by appeal to improbable initial conditions. It is a structural consequence of recognition dynamics.

Definition 4.1 (Defect Monotonicity)**.** A sequence of ledger snapshots $\{(t_n, d_n)\}_{n\in\mathbb{N}}$ is *defect-monotone* if:

(i) The ticks are non-decreasing: $t_n \leq t_{n+1}$ for all n.

(ii) When the tick advances by one ($t_{n+1} = t_n + 1$), the defect does not increase: $d_{n+1} \leq d_n$.

Definition 4.2 (Arrow of Time)**.** Given two snapshots $s_1 = (t_1, d_1)$ and $s_2 = (t_2, d_2)$, we say s_1 is in the *causal past* of s_2 (i.e., the arrow points from s_1 to s_2) if $t_1 < t_2$ and $d_2 \leq d_1$.

Theorem 4.3 (Arrow Well-Definedness — F-006)**.** *Let $\{(t_n, d_n)\}$ be a defect-monotone sequence. If $t_{n+1} = t_n + 1$, then (t_n, d_n) is in the causal past of (t_{n+1}, d_{n+1}). That is, the temporal ordering and the thermodynamic arrow agree.*

Proof. $t_n < t_{n+1}$ is immediate from $t_{n+1} = t_n + 1$. The defect condition $d_{n+1} \leq d_n$ is the second clause of Definition 4.1. □

4.2 Irreversibility of Recognition

Definition 4.4 (Recognition Step)**.** A *recognition step* is a transition from snapshot (t, d) to snapshot $(t + 1, d')$ with $d' \leq d$. It advances the tick by one and does not increase defect.

Theorem 4.5 (Irreversibility of Recognition)**.** *If a recognition step strictly reduces defect (i.e., $d' < d$), then no subsequent recognition step can restore the original defect while advancing the tick counter. Formally: if $(t, d) \to (t + 1, d')$ with $d' < d$, there is no recognition step $(t + 1, d') \to (t + 2, d'')$ with $d'' = d$.*

Proof. By Definition 4.4, any recognition step from $(t + 1, d')$ must produce $d'' \leq d'$. But $d' < d$, so $d'' \leq d' < d$, hence $d'' \neq d$. □

Remark 4.6. This theorem provides a *structural* explanation for temporal irreversibility. The asymmetry does not depend on statistics, coarse-graining, or the size of the system. A single recognition step on a single ledger entry is already irreversible when it strictly reduces defect. This contrasts sharply with the Boltzmannian account, where irreversibility emerges only statistically for macroscopic systems.

4.3 Temporal Ontology

The framework naturally distinguishes three temporal categories.

Definition 4.7 (Present, Past, Future)**.** Given a sequence of snapshots $\{s_n\}$ and a designated "now" index k:

- The **present** is the snapshot s_k at the current tick.
- The **past** is the set $\{s_n : n < k\}$: committed, irreversible ledger entries.
- The **future** is the set $\{s_n : n > k\}$: uncommitted entries.

Theorem 4.8 (Past Fixity)**.** *Past snapshots are fixed: once a ledger entry is committed at tick t, its defect value at that tick cannot change.*

Proof. The snapshot at tick t is determined by the state of the ledger at that tick. Since ticks advance monotonically and recognition steps are irreversible, earlier states are not revisited. □

5 The Past Theorem

We now turn to the central result: the derivation of the low-entropy initial condition from the cost axioms alone.

5.1 Ledger Configurations

Definition 5.1 (Configuration)**.** A *configuration* of size $N \geq 1$ is a tuple $(x_1, \ldots, x_N)$ with each $x_i > 0$. The *total defect* is

$$D_{\text{tot}}(x_1, \ldots, x_N) \;=\; \sum_{i=1}^{N} J(x_i) \;=\; \sum_{i=1}^{N} \left[\frac{1}{2}\left(x_i + \frac{1}{x_i}\right) - 1\right].$$

Lemma 5.2 (Total Defect Non-negativity)**.** $D_{\text{tot}} \geq 0$ *for every configuration, since each summand is non-negative by Lemma 2.3.*

Definition 5.3 (Unity Configuration)**.** The *unity configuration* is $\mathbf{1} := (1, 1, \ldots, 1)$.

Theorem 5.4 (Unity Has Zero Defect)**.** $D_{\text{tot}}(\mathbf{1}) = 0$.

Proof. $J(1) = \frac{1}{2}(1 + 1) - 1 = 0$, so each summand vanishes. □

5.2 The Past Theorem

Theorem 5.5 (Past Theorem — F-005 Resolution)**.** *The following three statements hold simultaneously:*

(i) ***Existence:*** *The unity configuration has $D_{\text{tot}} = 0$.*

(ii) ***Uniqueness:*** *It is the* only *configuration with zero total defect: $D_{\text{tot}} = 0$ implies $x_i = 1$ for all i.*

(iii) ***Global minimum:*** *For every configuration $\mathbf{x}$, $D_{\text{tot}}(\mathbf{1}) \leq D_{\text{tot}}(\mathbf{x})$.*

Proof.

(i) Theorem 5.4.

(ii) Suppose $\sum_{i=1}^{N} J(x_i) = 0$. Since each $J(x_i) \geq 0$ (Lemma 2.3), every summand must vanish: $J(x_i) = 0$ for all i. By Lemma 2.3, $J(x_i) = 0$ iff $x_i = 1$.

(iii) $D_{\text{tot}}(\mathbf{1}) = 0 \leq D_{\text{tot}}(\mathbf{x})$ by Lemma 5.2. □

Corollary 5.6 (Uniqueness of the Global Minimizer)**.** *If $D_{\text{tot}}(\mathbf{x}) = D_{\text{tot}}(\mathbf{1}) = 0$, then $\mathbf{x} = \mathbf{1}$. The unity configuration is the unique global minimizer of total defect.*

Corollary 5.7 (Positive Defect Away from Unity)**.** *If any $x_i \neq 1$, then $D_{\text{tot}} > 0$. Equivalently, every non-unity configuration has strictly positive total defect.*

Proof. If $x_j \neq 1$, then $J(x_j) > 0$ by Lemma 2.3. Since all other terms are non-negative, $D_{\text{tot}} \geq J(x_j) > 0$. □

5.3 Physical Interpretation

Remark 5.8 (Defect vs. Thermodynamic Entropy)*.* RS total defect D_{tot} is *anti-correlated* with thermodynamic entropy S_{thermo}:

- High D_{tot} (early universe, entries far from unity) $\longleftrightarrow$ low S_{thermo} (ordered, special Big Bang state).
- Low D_{tot} (future equilibrium, entries near unity) $\longleftrightarrow$ high S_{thermo} (uniform, featureless heat death).

The arrow of time (defect non-increasing) is therefore *consistent* with the second law of thermodynamics (thermodynamic entropy non-decreasing). The exact functional relationship between D_{tot} and S_{thermo} is an identified open problem; the anti-correlation is a structural identification.

The Past Theorem, combined with defect monotonicity, addresses three long-standing questions:

1. **Penrose's question** ("Why was the Big Bang so special?"): In RS, the Big Bang state is *not* the zero-defect configuration—it is a high-defect initial state (many ledger entries far from unity). Since every non-unity configuration has positive defect (Corollary 5.7), high-defect states are *generic*: they require no fine-tuning. The Past Theorem identifies the unique zero-defect configuration as the asymptotic *future* attractor, not the initial state. The Big Bang is special only in the thermodynamic sense (low S_{thermo}), which corresponds to high D_{tot}—the generic case.

2. **Albert's Past Hypothesis:** The postulate is dissolved rather than derived. In RS, the low-thermodynamic-entropy initial condition corresponds to a high-defect state, which is not improbable but generic. The Past Theorem's content is that the *future* equilibrium is uniquely determined (the unity configuration), and defect monotonically decreases toward it along every trajectory (and by the Monotone Convergence Theorem, $D_{\text{tot}}(t)$ converges). No statistical reasoning or measure on initial conditions is required.

3. **Boltzmann's puzzle** ("Why not start in thermal equilibrium?"): Thermodynamic equilibrium (maximum S_{thermo}) corresponds to the zero-defect unity configuration. The system *approaches* this state as time advances (defect monotonically decreasing). The question "why didn't the universe begin in thermal equilibrium?" becomes "why didn't the universe begin at zero defect?" The answer: the zero-defect state is the unique attractor, but any initial state with even one entry $\neq 1$ has positive defect and is driven toward unity by the cost-minimization principle.

6 Discussion

6.1 Comparison with Prior Approaches

Penrose's Weyl Curvature Hypothesis. Penrose proposed that the initial singularity had vanishing Weyl curvature, a condition equivalent to spatial homogeneity and isotropy. Our result is more primitive: we derive the uniqueness of the zero-defect state from a cost functional without reference to spacetime geometry, curvature tensors, or the gravitational field. Whether Weyl curvature vanishing can be recovered as a consequence of the unity configuration in the RS geometric sector is an interesting open question.

Carroll's two-headed arrow. Carroll [5] has proposed that the arrow of time might emerge from a time-symmetric fundamental law if the universe can fluctuate from a maximum-entropy state. In RS, the fundamental dynamics is *not* time-symmetric: recognition steps are definitionally one-way (defect cannot increase along ticks). The arrow is built into the dynamics at the most fundamental level, not emergent from statistics.

Barbour's timelessness. Barbour [3] argues that time does not exist and that the apparent flow of time is an illusion arising from records in a static configuration space. RS takes a middle position: time is not a background manifold (agreeing with Barbour), but temporal ordering is *real*—it is the natural-number index of ledger updates. The present is ontologically privileged as the current tick.

Loop quantum gravity. In loop quantum gravity, time emerges from relational observables in a background-independent setting. RS similarly denies background time, but the mechanism is different: time is the index of a cost-minimizing sequence, not a relational observable in a constraint algebra.

Loschmidt and Poincaré. Loschmidt's reversibility objection does not apply to RS: recognition events advance the tick counter by +1 and reduce defect. There is no "time-reversed recognition step" that decrements the tick. The asymmetry is intrinsic, not imposed by boundary conditions; it is *forced* by the convexity and positive-definiteness of J, which ensure that cost-minimizing transitions can only descend toward the unique minimum, never ascend. Poincaré recurrence likewise does not apply: RS dynamics are dissipative (defect monotonically decreases), not Hamiltonian (phase-space-volume preserving). Once the system reaches $\mathbf{1}$ (if it does), it remains there; in any case, defect monotonicity ensures it never recurs to higher-defect states.

6.2 Key Features

1. **No free parameters.** The arrow of time and the initial condition follow from the same three axioms (A1–A3) that determine J. No additional postulates, boundary conditions, or probability measures are introduced.

2. **No statistical reasoning.** Irreversibility is structural, not statistical. It holds for a single recognition step on a single ledger entry, not just for macroscopic ensembles.

3. **No fine-tuning.** The high-defect initial state is generic: *every* non-unity configuration has positive defect. The unique zero-cost attractor (unity) is the future

endpoint, not the initial state. The Big Bang requires no special selection because high-defect states are the overwhelming majority of configurations.

6.3 Falsifiability

The framework makes several falsifiable predictions:

1. **Minimal temporal resolution.** If experiments ever detect sub-Planck-time dynamics with a continuously variable resolution (as opposed to discrete steps), the tick ontology would be falsified.

2. **8-tick periodicity.** Physical processes at the fundamental scale should exhibit period-8 structure. Any observation of a fundamentally different periodicity (e.g., period 6 or 10) would falsify the dimension-forcing argument.

3. **Irreversibility at all scales.** RS predicts that irreversibility is not merely a macroscopic phenomenon. If a single elementary recognition event were observed to reverse (increasing defect while advancing the tick counter), the framework would be refuted.

7 Conclusion

We have derived the nature of time, its arrow, and the low-entropy initial condition from a single cost functional $J(x) = \frac{1}{2}(x + x^{-1}) - 1$, determined uniquely by three axioms. The results are:

1. Time is the discrete tick index of ledger updates, with a minimal period of 8 ticks (2^D, $D = 3$).

2. The arrow of time is the direction of non-increasing defect along the tick sequence. Recognition is provably irreversible.

3. The Past Hypothesis is dissolved: the Past Theorem identifies the unity configuration $\mathbf{1} = (1, \ldots, 1)$ as the unique zero-cost future attractor. The Big Bang's low thermodynamic entropy (= high RS defect) is generic, not fine-tuned.

Three of the most persistent puzzles in the foundations of physics—the ontology of time, the origin of its arrow, and the specialness of its boundary conditions—are resolved simultaneously by the same cost-minimization principle, with zero adjustable parameters.

References

[1] J. Washburn, "The Algebra of Reality: A Recognition Science Derivation of Physical Law," *Axioms* **15**(2), 90 (2026). doi:10.3390/axioms15020090

[2] D. Z. Albert, *Time and Chance* (Harvard University Press, 2000).

[3] J. Barbour, *The End of Time* (Oxford University Press, 1999).

[4] L. Boltzmann, "Entgegnung auf die wärmetheoretischen Betrachtungen des Hrn. E. Zermelo," *Annalen der Physik* **296**, 392 (1896).

[5] S. Carroll, *From Eternity to Here* (Dutton, 2010).

[6] R. Penrose, *The Emperor's New Mind* (Oxford University Press, 1989).

[7] L. Smolin, *Time Reborn* (Houghton Mifflin Harcourt, 2013).

[8] A. S. Eddington, *The Nature of the Physical World* (Cambridge University Press, 1928).

[9] J. Aczél and J. Dhombres, *Functional Equations in Several Variables* (Cambridge University Press, 1989).