

Comparative Analysis of Palbociclib and Ribociclib: A real world data and Propensity Score-Adjusted Evaluation with endocrine therapy

Author One¹, Author Two¹, Author Three^{2,3}, and Author Four¹

¹ Author one affiliation

² Author two affiliation

³ Author three affiliation

Corresponding author:

Author Four¹

Email address: email@address

ABSTRACT

Keywords: Keyword1; Keyword2; Keyword3

INTRODUCTION

Currently, metastatic breast cancer is difficult to treat. Patients with Hormone Receptor-positive (HR+) and Human Epidermal Growth Factor Receptor 2-negative (HER2-) breast cancer, the most common subtype, typically undergo endocrine therapy. Therefore, new treatments can be very useful in improving quality of life, reducing toxicity, and decreasing scenarios of hormonal resistance. Medications from the group of cyclin-dependent kinase inhibitors appear as a potential improvement in the therapeutic approach to advanced breast cancer. Within this group, there are palbociclib, ribociclib and abemaciclib. Cyclin-dependent kinases 4 and 6 (CDK4/6) are responsible for regulating the cell cycle at the transition between the G1 and S phases. In many neoplasms, this cycle is deregulated, and it promotes uncontrolled cell proliferation. It is then possible for these medications to have better effectiveness. These medications were approved by INFARMED, I.P. after an analysis of the therapeutic value they offer. This decision was made based on data provided by clinical trials done with these medications. The MONALEESA [8, 10, 12] studies were used for ribociclib, PALOMA [13, 9, 4] for palbociclib, and MONARCH [5, 11] for abemaciclib. These studies focused on testing the hypothesis of treating CDK4/6 inhibitors in combination with an aromatase inhibitor or fulvestrant as an alternative to the gold standard. In these research findings, it was determined that there was a notable enhancement in effectiveness, supporting their application in clinical practice. However, this evaluation was based on clinical trials with very specific inclusion and exclusion criteria and in a highly controlled environment. It is then vital to study how these new molecules compare to current practice in terms of treatment effectiveness in a real-world setting. In the meticulously controlled setting of clinical trials, patient selection often skews towards relatively healthier individuals with fewer comorbidities. However, in real-world clinical practice, patients present a diverse range of health profiles, co-existing illnesses, and medication histories that may influence drug efficacy and safety. Real-world data, drawn from electronic health records, insurance claims databases, and patient registries, offers the advantage of reflecting a more heterogeneous patient population, thus potentially uncovering insights not readily apparent in clinical trial settings. Understanding the effectiveness and safety of CDK4/6 inhibitors in real-world conditions is crucial for tailoring more individualized treatment regimens, optimizing outcomes, and enhancing the quality of life for patients with HR+, HER2- breast cancer [7]. Nevertheless, observational studies have inherent limitations, such as confounding by indication, which can lead to biased estimates of treatment effects. To tackle this, there are causality-based assessments that can be employed in order to better estimate the causal effects of treatments. Incorporating statistical techniques like Inverse Probability of Treatment Weighting (IPTW) can play an essential role in enhancing the quality of real-world evidence by accounting for treatment selection bias and balancing observed

covariates between treatment groups. IPTW, grounded in the framework of causal inference, allows for the mimicking of a randomized control trial-like setting within observational studies. By assigning weights to individual patients based on their propensity scores—the likelihood of receiving a particular treatment given a set of observed characteristics—analyses can achieve a balance between different treatment arms, thereby reducing bias and confounding factors. Establishing causality, rather than mere association, is vital for the robust interpretation of real-world data. As we strive to understand the long-term impact, efficacy, and safety of CDK4/6 inhibitors in HR+, HER2- breast cancer, the rigorous application of IPTW and causal inference methods can substantially augment the validity of real-world findings, making them a more reliable basis for clinical decision-making [1, 2] So in this paper, we propose:

- To compare the effectiveness of the CDK4/6 inhibitors drug class in terms of progression-free survival (PFS) and overall survival (OS);
- Assess the Hazard Ratio of using the CDK4/6 inhibitors drug class in terms of PFS and OS.
- To compare the effectiveness of CDK4/6 inhibitors in combination with an aromatase inhibitor or fulvestrant with the current standard of care in terms of PFS and OS in patients with HR+, HER2-advanced breast cancer.
- assess the differences in effectiveness between the three CDK4/6 inhibitors in combination with an aromatase inhibitor or fulvestrant in terms of PFS and OS with causality principles in mind, especially the counterfactual theory and IPTW ;

MATERIALS AND METHODS

0.1 Study Design

This retrospective study was designed in 2022. The aim of the study was to evaluate the clinical benefit and long-term survival of patients with HR+/HER2- that started treatment with CDK4/6 inhibitors plus endocrine therapy in different lines of treatment between the 14th of March 2017 and the 31st of December 2021. The follow-up period was set until June 2022. Inclusion criteria: women and men, Hormone receptor-positive and HER2 negative in the primary tumor or metastatic site after biopsy. Exclusion criteria: Patients that had only ambulatory medication, and patients involved in clinical trials, diagnosed with other neoplasms or with active treatment during the study period. The comparison group was defined by a population of patients, that were treated with hormone therapy as first-line (due to bone metastases) between 2015 and 13 of March 2017.

The evaluation of effectiveness will involve overall survival and progression-free analysis. We will compare the three different cyclin-dependent kinase inhibitors in terms of efficacy in real-world patients and will also compare the effectiveness of this class of drug against traditional endocrine therapy.

0.2 Data collection

All data were collected from original medical records from baseline to last visit or death. The data was collected from Instituto Português de Oncologia – Porto (IPO-P). Table 1 shows a comparison between the groups. Data included for population treated with CDK4/6 inhibitors plus endocrine therapy: demographic information, age at first diagnosis and age at the beginning of treatment, clinical characteristics and performance status by Eastern Cooperative Oncology Group scale (ECOG), treatment line and treatment schema - CDK4/6 inhibitor and endocrine therapy, stage of cancer, site of metastases (bone, soft tissue, visceral, central nervous system-CNS with or without another site). Data included for the population treated with endocrine therapy as first-line: demographic information, age at first diagnosis and age at the beginning of treatment, clinical characteristics and performance status by Eastern Cooperative Oncology Group scale (ECOG), stage of the cancer.

For comparison purposes, we used palbociclib and ribociclib since we had a small number of patients treated with abemaciclib (12).

0.3 Statistical Analysis

R was used for statistical analysis. Demographic, clinical characteristics and side effects were analyzed using descriptive statistics (count, percentages, and median/range). Kaplan–Meier test was used to determine the median PFS and OS in the entire population and subgroups. Log-rank test was used for comparisons of PFS and OS among different subgroups. Cox Regression was used to assess feature importance and impact. All statistical tests were two-sided, and the significance level was 0.05. The evaluation of the proportional hazards assumptions was done by Schoenfeld residues analysis. We applied propensity score weights to achieve a more robust comparison between the two groups of CDK4/6i. We

Table 1. Descriptive statistics of cyclin-dependent kinase inhibitors group and endocrine therapy group. The Drug/combination refers to the actual drug or the combination for CDK4/6

	ET	Palbociclib	Ribociclib
	(N=43)	(N=246)	(N=106)
Age at treatment start			
Mean (SD)	60.1 (12.4)	59.2 (11.7)	58.2 (10.7)
Median [Min, Max]	62.0 [34.0, 85.0]	60.0 [28.0, 84.0]	58.0 [32.0, 79.0]
Bone Only metastases			
No	NA	161 (65 %)	74 (70 %)
Yes	NA	85 (35 %)	32 (30 %)
Missing	43 (100%)	0 (0%)	0 (0%)
Visceral metastasis			
No	NA	121 (49 %)	49 (46 %)
Yes	NA	125 (51 %)	57 (54 %)
Missing	43 (100%)	0 (0%)	0 (0%)
Stage			
I	3 (7 %)	22 (9 %)	7 (7 %)
II	20 (47 %)	75 (30 %)	22 (21 %)
III	11 (26 %)	74 (30 %)	18 (17 %)
IV	2 (5 %)	65 (26 %)	46 (43 %)
Missing	7 (16.3%)	10 (4.1%)	13 (12.3%)
Drug/Combination			
Anastrozol	3 (7 %)	NA	NA
Exemestane	4 (9 %)	NA	NA
Fulvestrant	5 (12 %)	180 (73 %)	10 (9 %)
Letrozol	31 (72 %)	66 (27 %)	96 (91 %)

used the existence of visceral metastases, menopausal status, combination, treatment line, age at treatment start, ECOG, and stage. Likewise, we used the WeightIt package for R [6]. Furthermore, we applied the weights to get the ATE which is $E[Y_i(1) - Y_i(0)]$, the average effect of moving an entire population from untreated to treated, or from one drug to the other. Weights were used instead of matching since it is more suited for calculating ATE and the need to preserve the sample size since it is already small from the start. The formula for calculating the weights was through propensity score weighting with GLM. Multiple comparisons were done with the Benjamini-Hochberg (BH) method.

RESULTS


The median OS in the entire population treated with CDK4/6 inhibitors was 46 months (95%CI 39.4–55.6). Median PFS was 20.1 months (95%CI 18.3–24.2). Following this, we compared Palbociclib and riboclib as first-line treatments. We found that regarding OS, there is no significant difference between the two, but ribociclib is significantly better in terms of PFS (p-value ≤ 0.001) (figure 1). Additionally, we compared the same CDK4/6 inhibitors with letrozol as a combination only. Regarding this scenario, we found out that both were similar in terms of OS and PFS.

We then compared both with a cox regression, where OS shows no significant difference between palbociclib and ribociclib but a significantly better PFS for ribociclib (figure 2) HR 0.60 [95%CI 0.36-0.97] when adjusted to the stage, visceral metastases, age, treatment line, combination and ECOG. This data implies that ribociclib reduces the risk of the disease progression by 40% compared to palbociclib when adjusted for the variables mentioned. The proportional hazards assumption was confirmed with p values all over 0.10.

When comparing endocrine therapy with CDK4/6 inhibitors as first-line treatment (figure 2), we see that only Ribociclib is significantly better in terms of PFS and OS (p-value ≤ 0.001). When comparing palbociclib as the first line, we see that there is no significant difference in terms of PFS and OS (p=0.6 and 0.47). We also applied the same analysis as above, comparing only the letrozol combination with letrozol alone. We found that both ribociclib and palbociclib are significantly better in terms of PFS (HR 0.51 for palbociclib and 0.28 for ribociclib).

When comparing palbociclib and ribociclib adjusted for ATE weights, we found a different scenario

Figure 1. Survival curves for Palbociclib and Ribociclib (1st line) - Progression Free Survival and Overall Survival



from previous assessments. There is a significant difference between the two in terms of OS and PFS (figure 3). We calculated the weights taking into account stage, age at treatment start, treatment line, and ECOG.

The Cox regression adjusted for weights shows that ribociclib is not significantly different from palbociclib for OS. The HR for PFS is 0.54 [0.31-0.94;p=0.029], implying that ribociclib reduces the risk of the disease progression by ~50% compared to palbociclib when adjusted to the stage, combination drug, treatment line, visceral metastasis, age, and ECOG. Proportional hazard assumptions are confirmed as well.

DISCUSSION

This study aimed to evaluate the real-world use of palbociclib and ribociclib in combination with ET for HR+/HER2- and compare this drug class with traditional endocrine therapy. Few real-world evidence studies of palbociclib and ribociclib used in daily clinical practice have been published identifying clinical benefits, patient profiles, and sequencing of treatment, with even less evidence for the Portuguese population.

When comparing with clinical trials, regarding patient profile, in our study, 51% had visceral metastasis and 35% had bone-only metastases compared with 49% and 38% in PALOMA-2, and 60% and 25% in PALOMA-3, respectively [9, 3]. As for ribociclib and bone-only metastases, MONALEESA-7 [12] has 24% and MONALEESA-2 has 40% [8] and our study has 30%. Regarding menopausal status, our study has 20% premenopausal and 80% postmenopausal. The trials were conducted with postmenopausal women only.

Of note, the range of median PFS for first-line palbociclib was 15.5–25.5 months, which is shorter than 27.6 months observed in a post hoc analysis of the PALOMA-2 clinical trial with extended follow-up [9], but in line with RWE studies (13.3–20.2 months) [7]. When assessed with only letrozol as a combination, the median PFS increased to 28.6 months [95% CI 25.5-not reached]. Additionally, analyzing the postmenopausal women subgroup, palbociclib showed a median PFS of 16.3 months [95% CI 12.9 -20]. Furthering analysis of the postmenopausal and with letrozol, the median was 47.6 months [95% 25.6-2-not reached].

As for ribociclib, median survival time was not reached whether in OS and PFS. So we can at least say that the median PFS is longer than 50 months. This is longer than the median progression-free survival of 23.8 months (95% CI 19.2–not reached) reported in the MONALEESA-7 trial [12] and longer than 25.3 months (95% CI 23.0–30.3) in the MONALEESA-2 trial [8]. Regarding the subgroup analysis of postmenopausal women and postmenopausal women treated ribociclib in combination with letrozol, the median was not reached.

When directly comparing ribociclib and palbociclib without any adjustments, one might deduce that ribociclib is superior to palbociclib. However, after adjusting for confounding variables, there is no

Table 2. Cox Regression with palbociclib and Ribociclib - Progression Free Survival and Overall Survival

Characteristic	OS			PFS		
	HR [†]	95% CI [†]	p-value	HR [†]	95% CI [†]	p-value
Drug						
Palbociclib	—	—		—	—	
Ribociclib	1.04	0.54, 2.02	>0.9	0.60	0.36, 0.97	0.039
Combination						
Fulvestrant	—	—		—	—	
Letrozol	0.35	0.18, 0.66	0.001	0.45	0.28, 0.71	<0.001
Treatment Line						
1st Line	—	—		—	—	
2nd+ Lines	1.00	0.62, 1.64	>0.9	1.20	0.82, 1.77	0.3
Stage						
I	—	—		—	—	
II	5.67	1.36, 23.6	0.017	1.89	0.99, 3.62	0.055
III	8.17	1.95, 34.2	0.004	3.02	1.57, 5.80	<0.001
IV	8.06	1.91, 34.0	0.004	2.24	1.15, 4.35	0.017
Visceral Metastasis						
No	—	—		—	—	
Yes	1.75	1.19, 2.56	0.004	1.35	1.00, 1.82	0.049
Age at treatment start	1.00	0.98, 1.02	>0.9	0.99	0.97, 1.00	0.053
ECOG at treatment start						
0	—	—		—	—	
1	1.62	1.05, 2.49	0.030	1.22	0.88, 1.69	0.2
2	3.90	2.05, 7.41	<0.001	1.54	0.86, 2.76	0.15
3	12.4	3.55, 43.1	<0.001	0.39	0.05, 2.87	0.4

[†] HR = Hazard Ratio, CI = Confidence Interval

significant difference between the two inhibitors in terms of Progression-Free Survival (PFS) or Overall Survival (OS) as indicated in 2. This observation is further corroborated by the lower plots in 1, where even a subgroup analysis of CDK4/6i combined solely with letrozole reveals a trend towards nonsignificance.

In the first-line comparison between Endocrine Therapy (ET) and CDK4/6 inhibitors (CDK4/6i), there is no significant impact on Overall Survival (OS) by either treatment, whether the CDK4/6i is combined with both agents or solely with letrozole. With respect to Progression-Free Survival (PFS), ribociclib demonstrates superior efficacy in both combination therapies (HR=0.29) as well as when paired only with letrozole (HR=0.28). Additionally, palbociclib exhibits significant improvement in PFS when combined with letrozole (HR=0.50).

When comparing with propensity scores weighting, we found out that ribociclib is significantly better than palbociclib for PFS and OS, providing a median OS of over 40 months and median PFS of around 42 months. Adjusted for the weighted variables, Ribociclib is no longer significantly better for OS and has a p-value of exactly 0.05 for PFS. This is in line with the results of the Cox regression analysis.

CONCLUSIONS

In conclusion, our findings underscore the efficacy of CDK4/6 inhibitors in real-world settings. While we cannot definitively assert that palbociclib surpasses endocrine therapy in terms of Overall Survival—a facet not extensively explored in prominent clinical trials—we can confidently affirm the impact of CDK4/6i on Progression-Free Survival. This assertion aligns with clinical trial outcomes and real-world data further substantiates these findings.

Delving deeper into the characteristics of the patient population, including safety profiles, economic implications, and quality of life metrics, would be insightful. Additionally, a thorough examination of biomarkers within the population could offer invaluable insights. We intend to explore these facets in subsequent publications.

It's imperative to note that our data is sourced from a singular institution, limiting the generalizability

Figure 2. Survival curves (OS and PFS) comparing endocrine therapy (ET) to CDK4/6 inhibitors as 1st line. p values shown as pairwise vs. ET.





Figure 3. Comparison of palbociclib and ribociclib survival curves adjusted for propensity scores



184 of our results to a broader population. Nonetheless, we posit that this study lays a foundational groundwork
 185 for future research in this domain. While our evidence is rooted in observational data and we've made
 186 adjustments for known confounders, the potential for residual confounding remains. Although the use
 187 of propensity score matching enhances the comparative robustness between the groups, the presence of
 188 unmeasured confounders cannot be entirely ruled out. Furthermore, the small sample size of our study
 189 limits the statistical power of our findings. We hope that our study will serve as a springboard for future
 190 research in this domain, and we look forward to furthering our research in this area.

191 AUTHOR CONTRIBUTIONS

192 A.O., A.T. and A.F. conceived the presented idea; A.O. wrote the main manuscript; All authors have read
 193 and agreed to the published version of the manuscript.

194 INSTITUTIONAL REVIEW

195 This work was approved by the ... Ethics Committee...

196 DATA AVAILABILITY

197 ...

198 FUNDING

199 This work was supported ... For the purpose of open access, the author has applied a CC BY public
200 copyright licence to any Author Accepted Manuscript version arising from this submission.

201 ACKNOWLEDGMENTS

202 ...

203 CONFLICTS OF INTEREST

204 The authors declare no conflict of interest.

205 REFERENCES

- 206 [1] Peter C. Austin. An Introduction to Propensity Score Methods for Reducing the Effects of Confounding
207 in Observational Studies. 46(3):399–424.
- 208 [2] Peter C Austin. The use of propensity score methods with survival or time-to-event outcomes:
209 Reporting measures of effect similar to those used in randomized experiments. *Statistics in Medicine*,
210 33(7):1242–1258, 2014.
- 211 [3] Massimo Cristofanilli, Nicholas C. Turner, Igor Bondarenko, Jungsil Ro, Seock-Ah Im, Norikazu
212 Masuda, Marco Colleoni, Angela DeMichele, Sherene Loi, Sunil Verma, Hiroji Iwata, Nadia Harbeck,
213 Ke Zhang, Kathy Puyana Theall, Yuqiu Jiang, Cynthia Huang Bartlett, Maria Koehler, and Dennis
214 Slamon. Fulvestrant plus palbociclib versus fulvestrant plus placebo for treatment of hormone-
215 receptor-positive, HER2-negative metastatic breast cancer that progressed on previous endocrine
216 therapy (PALOMA-3): Final analysis of the multicentre, double-blind, phase 3 randomised controlled
217 trial. 17(4):425–439.
- 218 [4] Richard S Finn, John P Crown, Istvan Lang, Katalin Boer, Igor M Bondarenko, Sergey O Kulyk, Jo-
219 hannes Ettl, Ravindranath Patel, Tamas Pinter, Marcus Schmidt, Yaroslav Shparyk, Anu R Thummala,
220 Nataliya L Voytko, Camilla Fowst, Xin Huang, Sindy T Kim, Sophia Randolph, and Dennis J Slamon.
221 The cyclin-dependent kinase 4/6 inhibitor palbociclib in combination with letrozole versus letrozole
222 alone as first-line treatment of oestrogen receptor-positive, HER2-negative, advanced breast cancer
223 (PALOMA-1/TRIO-18): A randomised phase 2 study. *The Lancet Oncology*, 16(1):25–35, January
224 2015. <https://linkinghub.elsevier.com/retrieve/pii/S1470204514711593>.
- 225 [5] Matthew P. Goetz, Masakazu Toi, Mario Campone, Joohyuk Sohn, Shani Paluch-Shimon, Jens
226 Huober, In Hae Park, Olivier Trédan, Shin-Cheh Chen, Luis Manso, Orit C. Freedman, Georgina
227 Garnica Jalliffe, Tammy Forrester, Martin Frenzel, Susana Barriga, Ian C. Smith, Nawel Bourayou, and
228 Angelo Di Leo. MONARCH 3: Abemaciclib As Initial Therapy for Advanced Breast Cancer. *Journal*
229 *of Clinical Oncology: Official Journal of the American Society of Clinical Oncology*, 35(32):3638–
230 3646, November 2017.
- 231 [6] Noah Greifer. *WeightIt: Weighting for Covariate Balance in Observational Studies*, 2023.
232 <https://ngreifer.github.io/WeightIt/>, <https://github.com/ngreifer/WeightIt>.
- 233 [7] Nadia Harbeck, Meaghan Bartlett, Dean Spurdén, Becky Hooper, Lin Zhan, Emily Rosta, Chris
234 Cameron, Debanjali Mitra, and Anna Zhou. CDK4/6 inhibitors in HR+/HER2- advanced/metastatic
235 breast cancer: A systematic literature review of real-world evidence studies. 17(16):2107–2122.
- 236 [8] G. Hortobagyi, S. Stemmer, H. Burris, Y. Yap, G. Sonke, S. Paluch-Shimon, M. Campone, K. Petraková,
237 K. Blackwell, E. Winer, W. Janni, S. Verma, P. Conte, C. Arteaga, D. Cameron, S. Mondal, F. Su,
238 M. Miller, M. Elmehiegy, C. Germa, and J. O’Shaughnessy. Updated results from MONALEESA-2, a
239 phase III trial of first-line ribociclib plus letrozole versus placebo plus letrozole in hormone receptor-
240 positive, HER2-negative advanced breast cancer. *Annals of oncology : official journal of the European*
241 *Society for Medical Oncology*, 2018.
- 242 [9] H. S. Rugo, V. Diéras, K. A. Gelmon, R. S. Finn, D. J. Slamon, M. Martin, P. Neven, Y. Shparyk,
243 A. Mori, D. R. Lu, H. Bhattacharyya, C. H. U. a. N. G. Bartlett, S. Iyer, S. Johnston, J. Ettl, and
244 N. Harbeck. Impact of palbociclib plus letrozole on patient-reported health-related quality of life:
245 Results from the PALOMA-2 trial. *Annals of Oncology: Official Journal of the European Society for*
246 *Medical Oncology*, 29(4):888–894, April 2018.

- [10] Dennis J. Slamon, Patrick Neven, Stephen Chia, Peter A. Fasching, Michelino De Laurentiis, Seock-Ah Im, Katarina Petrakova, Giulia Val Bianchi, Francisco J. Esteva, Miguel Martín, Arnd Nusch, Gabe S. Sonke, Luis De la Cruz-Merino, J. Thaddeus Beck, Xavier Pivot, Gena Vidam, Yingbo Wang, Karen Rodriguez Lorenc, Michelle Miller, Tetiana Taran, and Guy Jerusalem. Phase III Randomized Study of Ribociclib and Fulvestrant in Hormone Receptor-Positive, Human Epidermal Growth Factor Receptor 2-Negative Advanced Breast Cancer: MONALEESA-3. *Journal of Clinical Oncology: Official Journal of the American Society of Clinical Oncology*, 36(24):2465–2472, August 2018.
- [11] George W. Sledge, Masakazu Toi, Patrick Neven, Joohyuk Sohn, Kenichi Inoue, Xavier Pivot, Olga Burdaeva, Meena Okera, Norikazu Masuda, Peter A. Kaufman, Han Koh, Eva-Maria Grischke, Martin Frenzel, Yong Lin, Susana Barriga, Ian C. Smith, Nawel Bourayou, and Antonio Llombart-Cussac. MONARCH 2: Abemaciclib in Combination With Fulvestrant in Women With HR+/HER2-Advanced Breast Cancer Who Had Progressed While Receiving Endocrine Therapy. *Journal of Clinical Oncology: Official Journal of the American Society of Clinical Oncology*, 35(25):2875–2884, September 2017.
- [12] Debu Tripathy, Seock-Ah Im, Marco Colleoni, Fabio Franke, Aditya Bardia, Nadia Harbeck, Sara A. Hurvitz, Louis Chow, Joohyuk Sohn, Keun Seok Lee, Saul Campos-Gomez, Rafael Villanueva Vazquez, Kyung Hae Jung, K. Govind Babu, Paul Wheatley-Price, Michelino De Laurentiis, Young-Hyuck Im, Sherko Kuemmel, Nagi El-Saghir, Mei-Ching Liu, Gary Carlson, Gareth Hughes, Ivan Diaz-Padilla, Caroline Germa, Samit Hirawat, and Yen-Shen Lu. Ribociclib plus endocrine therapy for premenopausal women with hormone-receptor-positive, advanced breast cancer (MONALEESA-7): A randomised phase 3 trial. *The Lancet. Oncology*, 19(7):904–915, July 2018.
- [13] Sunil Verma, Cynthia Huang Bartlett, Patrick Schnell, Angela M. DeMichele, Sherene Loi, Jungsil Ro, Marco Colleoni, Hiroji Iwata, Nadia Harbeck, Massimo Cristofanilli, Ke Zhang, Alexandra Thiele, Nicholas C. Turner, and Hope S. Rugo. Palbociclib in Combination With Fulvestrant in Women With Hormone Receptor-Positive/HER2-Negative Advanced Metastatic Breast Cancer: Detailed Safety Analysis From a Multicenter, Randomized, Placebo-Controlled, Phase III Study (PALOMA-3). *The Oncologist*, 21(10):1165–1175, October 2016.

FIGURE TABLE

