Counts, laws, denominators and API field names

Fix report g, information y and normalized nonnegative weights. Choose ν = s (supplied) or P (evaluation).

Raw report frame

$$M = \binom{n}{2} > 0, \quad x_{ij} = \mathbf{1}\{g_i < g_j\}, \quad N(g) = \sum_{i \neq j} x_{ij}$$

$N(g)/M$ is raw disclosure; it is fixed by g.

Realized/action object	Conditional expectation	Rate and condition
Weak direction error (≤)		
$D_k(g, \Theta) = \sum_{i \neq j} x_{ij} \mathbf{1}\{\theta_{ik} \leq \theta_{jk}\}$	$\mathbb{E}_\nu[\cdot \mid y]$ → $E_k^\nu(g) = \mathbb{E}_\nu[D_k \mid y]$	divide → $E_k^\nu(g)/N(g)$, $N(g) > 0$
Strict reversal (<)		
$\sum_{i \neq j} x_{ij} \mathbf{1}\{\theta_{ik} < \theta_{jk}\}$	$\mathbb{E}_\nu[\cdot \mid y]$ → $E_k^{b,\nu}(g)$ Expected strict-error count	divide → $E_k^{b,\nu}(g)/N(g)$, $N(g) > 0$; $E_k^{b,\nu}(g)/Q_k^\nu(g)$, $Q_k^\nu(g) > 0$
Distinct opportunity (≠)		
$\sum_{i \neq j} x_{ij} \mathbf{1}\{\theta_{ik} \neq \theta_{jk}\}$	$\mathbb{E}_\nu[\cdot \mid y]$ → $Q_k^\nu(g)$ Expected distinct mass	weight; divide → $Q_w^\nu(g)/T_w^\nu$, $T_w^\nu > 0$

Weighting and total opportunity

$$Q_w^\nu(g) = \sum_k w_k Q_k^\nu(g), \quad T_w^\nu = \sum_{i<j} \sum_k w_k d_{ij,k}^\nu$$

$$d_{ij,k}^\nu = \Pr_\nu(\theta_{ik} \neq \theta_{jk} \mid y)$$

T uses every original unordered pair; Q uses disclosed directions.

Inspected tie_aware_v1 field-name correspondence

API D ↔ $Q_w^s(g)$; API T ↔ T_w^s

Supplied-law aliases; book D_k remains a realized wrong-count.

$N = 0$

All disclosed counts are zero. N-rates: NA; raw $N/M = 0$. $Q_w^\nu/T_w^\nu = 0$ only if $T_w^\nu > 0$.

$Q_k^\nu = 0$

$E_k^{b,\nu}/Q_k^\nu$ is NA. No conditional-distinct accuracy is assigned.

$T_w^\nu = 0$

Q_w^ν/T_w^ν is NA; neutral risk and R_0^ν are zero. Raw N can be positive outside a guarded family.

When reports or datasets vary, let $A = \sum_k w_k D_k$. E[A]/E[N] and E[A/N | N > 0] generally differ.

Name that averaging law and a positive denominator. Different types can have equal numerical values.