## 1 Energy

**4.2** From the origin O to point P = (1,1) a two dimensional force  $\mathbf{F} = (x^2, 2xy)$  moves a point along three paths where the work done by the force is

$$W = \int_{Q}^{P} \mathbf{F} \cdot d\mathbf{r} = \int_{Q}^{P} F_x dx + F_y dy$$

(a) Splitting the path into two parts  $O \to Q = (1,0)$  and  $Q \to P$ , we have two integrals

$$W = \int_{Q}^{Q} F_x \, \mathrm{d}x + \int_{Q}^{P} F_y \, \mathrm{d}y$$

where the first integral accounts for just the x component of force  $F_x = x^2$  and the second integral accounts for just the y component of force when x = 1;  $F_y = 2(1)y$ . Thus

$$W = \int_0^1 x^2 \, \mathrm{d}x + \int_0^1 2y \, \mathrm{d}y = \frac{4}{3}$$

(b) The path follows the parabola  $y = x^2$  from  $O \to P$ . From dy = 2x dx the integral can be rewritten in terms of just x

$$W = \int_0^1 x^2 dx + \int_0^1 2x(x^2) dy = \frac{1}{3} + \int_0^1 4x^4 dx = \frac{17}{15}$$

(c) Path follows the parametric curve  $x=t^3$  and  $y=t^2$  where the differentials are:  $dx=3t^2\,\mathrm{d}t$  and  $dy=2t\,\mathrm{d}t$ . Thus the work done on the path is

$$W = \int_0^1 (t^6)(3t^2 dt) + \int_0^1 (2t^3)(2t dt) = \frac{1}{3} + \frac{4}{5} = \frac{19}{15}$$

**4.3** Same as Problem 4.2 but with a force  $\mathbf{F} = (-y, x)$  and three different paths from  $P = (1, 0) \rightarrow Q = (0, 1)$ .

(a) This path follows a straight line y=0 from  $P\to O$  and then x=0 from  $O\to Q$ . Thus the work done is

$$W = \int_{P}^{O} F_x \, \mathrm{d}x + \int_{O}^{Q} F_y \, \mathrm{d}y = 0$$

(b) A straight line from  $P \to Q$  is given by y = -x + 1 and the differential dy = -dx. Thus the work done is

$$W = \int_{P}^{Q} F_x \, dx + F_y \, dy = \int_{1}^{0} (-(-x+1)) \, dx + (x)(-dx) = \int_{1}^{0} -1 \, dx = 1$$

(c) The path of a quarter circle centered on the origin in polar coordinates is given by

$$x = r\cos\phi$$
  $y = r\sin\phi$ 

where  $r=1,\,\phi=0\to\pi/2$  and the differentials are

$$dx = \cos\phi dr - r\sin\phi d\phi = -\sin\phi d\phi$$
  $dy = \sin\phi dr + r\cos\phi d\phi = \cos\phi d\phi$ 

Thus the work done is

$$W = \int_{P}^{Q} F_x \, dx + F_y \, dy = \int_{0}^{\pi/2} (-\sin\phi)(-\sin\phi \, d\phi) + (\cos\phi)(\cos\phi \, d\phi) = \int_{0}^{\pi/2} d\phi = \frac{\pi}{2}$$

1

**4.5** (a) Given the force of gravity  $\mathbf{F} = -mg\mathbf{\hat{y}}$  and vertical height from 1 to 2  $h = y_2 - y_1$ , the work done by gravity is

$$W_g(1 \to 2) = \int_1^2 \mathbf{F} \cdot d\mathbf{r} = \int_0^h -mg \, dy = -mgh$$

Since the force  $\mathbf{F}$  depends only on position and the work done by is independent of the path taken, the force is conservative.

(b) The gravitational potential energy of the particle is

$$U_g(\mathbf{r}) = -W_g(0 \to \mathbf{r}) = -\int_0^{\mathbf{r}} \mathbf{F} \cdot d\mathbf{r} = -\int_0^{\mathbf{r}} -mg \, dy = mgy$$

where  $\mathbf{r} = y\hat{\mathbf{y}}$  is the position vector of the particle. The potential energy is a function