

Problems for Griffiths' Electrodynamics

Contents

1	Vector Analysis	2
---	-----------------	---

1 Vector Analysis




Figure 1.1: Three Coplanar Vectors

1.1 (a) When three vectors are coplanar as shown in Figure 1.1, the dot product is

$$\begin{aligned}\mathbf{A} \cdot (\mathbf{B} + \mathbf{C}) &= \mathbf{A} \cdot \mathbf{B} + \mathbf{A} \cdot \mathbf{C} \\ A(B + C) \cos \theta &= AB \cos \theta_B + AC \cos \theta_C\end{aligned}$$

Since $B \cos \theta_B + C \cos \theta_C = (B + C) \cos \theta$ from Figure 1.1, the distributive property holds true. The cross product also holds true since $B \sin \theta_B + C \sin \theta_C = (B + C) \sin \theta$, and multiplying by A on both sides gives the same result as the distributive property:

$$\begin{aligned}\mathbf{A} \times (\mathbf{B} + \mathbf{C}) &= \mathbf{A} \times \mathbf{B} + \mathbf{A} \times \mathbf{C} \\ A(B + C) \sin \theta &= AB \sin \theta_B + AC \sin \theta_C\end{aligned}$$

(b) In the general case in three-dimensional space, each vector has three components: $\mathbf{A} = (A_x, A_y, A_z)$. Therefore,

$$\begin{aligned}\mathbf{A} \cdot (\mathbf{B} + \mathbf{C}) &= \mathbf{A} \cdot (B_x + C_x, B_y + C_y, B_z + C_z) \\ &= A_x(B_x + C_x) + A_y(B_y + C_y) + A_z(B_z + C_z) \\ &= A_x B_x + A_y B_y + A_z B_z + A_x C_x + A_y C_y + A_z C_z \\ &= \mathbf{A} \cdot \mathbf{B} + \mathbf{A} \cdot \mathbf{C}\end{aligned}$$

1.2 Setting $\mathbf{A} = \mathbf{B} = (1, 1, 1)$ and $\mathbf{C} = (1, 1, -1)$:

$$\begin{aligned}(\mathbf{A} \times \mathbf{B}) \times \mathbf{C} &\stackrel{?}{=} \mathbf{A} \times (\mathbf{B} \times \mathbf{C}) \\ 0 &\stackrel{?}{=} (1, 1, 1) \times [(1, 1, 1) \times (1, 1, -1)] \\ 0 &\stackrel{?}{=} (1, 1, 1) \times (-2, 2, 0) \\ 0 &\neq (-2, -2, 4)\end{aligned}$$

where the cross product of parallel vectors $\mathbf{A} \times \mathbf{B} = 0$. Therefore, the cross product is not associative.

1.3 Taking the dot product of a unit cube's body diagonals $\mathbf{A} = (1, 1, 1)$, $\mathbf{B} = (1, 1, -1)$:

$$\begin{aligned}\mathbf{A} \cdot \mathbf{B} &= AB \cos \theta \\ 1 &= 3 \cos \theta \\ \theta &= \arccos 1/3 \approx 70.53^\circ\end{aligned}$$

1.4 The cross product of two vectors coplanar to the shaded plane— $\mathbf{A} = (-1, 2, 0)$, $\mathbf{B} = (-1, 0, 3)$ —is parallel to the normal unit vector $\hat{\mathbf{n}}$ of the plane:

$$\begin{aligned}\mathbf{A} \times \mathbf{B} &= \mathbf{C} \\ \begin{vmatrix} \hat{\mathbf{x}} & \hat{\mathbf{y}} & \hat{\mathbf{z}} \\ -1 & 2 & 0 \\ -1 & 0 & 3 \end{vmatrix} &= (6, 3, 2)\end{aligned}$$

where $\hat{\mathbf{n}} = \mathbf{C}/C$, and $C = \sqrt{6^2 + 3^2 + 2^2} = \sqrt{49} = 7$. Therefore,

$$\hat{\mathbf{n}} = \frac{1}{7}(6, 3, 2)$$

1.5 Proving the “BAC–CAB” rule for three-dimensional vectors:

$$\begin{aligned}\mathbf{A} \times (\mathbf{B} \times \mathbf{C}) &= \mathbf{A} \times \begin{vmatrix} \hat{\mathbf{x}} & \hat{\mathbf{y}} & \hat{\mathbf{z}} \\ B_x & B_y & B_z \\ C_x & C_y & C_z \end{vmatrix} \\ &= \begin{vmatrix} \hat{\mathbf{x}} & \hat{\mathbf{y}} & \hat{\mathbf{z}} \\ A_x & A_y & A_z \\ B_y C_z - B_z C_y & B_z C_x - B_x C_z & B_x C_y - B_y C_x \end{vmatrix}\end{aligned}$$

where the x component is $A_y(B_x C_y - B_y C_x) - A_z(B_z C_x - B_x C_z)$. Similarly,

$$\mathbf{B}(\mathbf{A} \cdot \mathbf{C}) - \mathbf{C}(\mathbf{A} \cdot \mathbf{B}) = \mathbf{B}(A_x C_x + A_y C_y + A_z C_z) - \mathbf{C}(A_x B_x + A_y B_y + A_z B_z)$$

where x component simplifies to

$$B_x(\cancel{A_x C_x} + A_y C_y + A_z C_z) - C_x(\cancel{A_x B_x} + A_y B_y + A_z B_z) = A_y(B_x C_y - B_y C_x) - A_z(B_z C_x - B_x C_z)$$

the same is done for the y and z components. Therefore, the “BAC–CAB” rule holds true.

1.6

$$\begin{aligned}[\mathbf{A} \times (\mathbf{B} \times \mathbf{C})] + [\mathbf{B} \times (\mathbf{C} \times \mathbf{A})] + [\mathbf{C} \times (\mathbf{A} \times \mathbf{B})] &= \cancel{\mathbf{B}(\mathbf{A} \cdot \mathbf{C})} - \mathbf{C}(\mathbf{A} \cdot \mathbf{B}) + \mathbf{C}(\mathbf{B} \cdot \mathbf{A}) \\ &\quad - \mathbf{A}(\mathbf{B} \cdot \mathbf{C}) + \mathbf{A}(\mathbf{C} \cdot \mathbf{B}) - \cancel{\mathbf{B}(\mathbf{C} \cdot \mathbf{A})}\end{aligned}$$

since dot product is associative, the first and last terms cancel out, and the middle terms also cancel out with each other.

$$\begin{aligned}\mathbf{A} \times (\mathbf{B} \times \mathbf{C}) &= (\mathbf{A} \times \mathbf{B}) \times \mathbf{C} \\ \mathbf{B}(\mathbf{A} \cdot \mathbf{C}) - \mathbf{C}(\mathbf{A} \cdot \mathbf{B}) &= \mathbf{B}(\mathbf{A} \cdot \mathbf{C}) - \mathbf{A}(\mathbf{B} \cdot \mathbf{C}) \\ 0 &= -\mathbf{C}(\mathbf{A} \cdot \mathbf{B}) + \mathbf{A}(\mathbf{B} \cdot \mathbf{C}) \\ 0 &= -\mathbf{B}(\mathbf{C} \times \mathbf{A}) = (\mathbf{C} \times \mathbf{A}) \times \mathbf{B}\end{aligned}$$

For the relation to hold true, either the vectors \mathbf{A} and \mathbf{C} are parallel ($\mathbf{A} \times \mathbf{C} = 0$) or \mathbf{B} is perpendicular to both \mathbf{A} and \mathbf{C} ($\mathbf{A} \cdot \mathbf{B} = \mathbf{B} \cdot \mathbf{C} = 0$).