

1 Vector Analysis




Figure 1.1: Three Coplanar Vectors

1.1 (a) When three vectors are coplanar as shown in Figure 1.1, the dot product is

$$\begin{aligned}\mathbf{A} \cdot (\mathbf{B} + \mathbf{C}) &= \mathbf{A} \cdot \mathbf{B} + \mathbf{A} \cdot \mathbf{C} \\ A(B + C) \cos \theta &= AB \cos \theta_B + AC \cos \theta_C\end{aligned}$$

Since $B \cos \theta_B + C \cos \theta_C = (B + C) \cos \theta$ from Figure 1.1, the distributive property holds true. The cross product also holds true since $B \sin \theta_B + C \sin \theta_C = (B + C) \sin \theta$, and multiplying by A on both sides gives the same result as the distributive property:

$$\begin{aligned}\mathbf{A} \times (\mathbf{B} + \mathbf{C}) &= \mathbf{A} \times \mathbf{B} + \mathbf{A} \times \mathbf{C} \\ A(B + C) \sin \theta &= AB \sin \theta_B + AC \sin \theta_C\end{aligned}$$

(b) In the general case in three-dimensional space, each vector has three components: $\mathbf{A} = (A_x, A_y, A_z)$. Therefore,

$$\begin{aligned}\mathbf{A} \cdot (\mathbf{B} + \mathbf{C}) &= \mathbf{A} \cdot (B_x + C_x, B_y + C_y, B_z + C_z) \\ &= A_x(B_x + C_x) + A_y(B_y + C_y) + A_z(B_z + C_z) \\ &= A_x B_x + A_y B_y + A_z B_z + A_x C_x + A_y C_y + A_z C_z \\ &= \mathbf{A} \cdot \mathbf{B} + \mathbf{A} \cdot \mathbf{C}\end{aligned}$$

1.2 Setting $\mathbf{A} = \mathbf{B} = (1, 1, 1)$ and $\mathbf{C} = (1, 1, -1)$:

$$\begin{aligned}(\mathbf{A} \times \mathbf{B}) \times \mathbf{C} &\stackrel{?}{=} \mathbf{A} \times (\mathbf{B} \times \mathbf{C}) \\ 0 &\stackrel{?}{=} (1, 1, 1) \times [(1, 1, 1) \times (1, 1, -1)] \\ 0 &\stackrel{?}{=} (1, 1, 1) \times (-2, 2, 0) \\ 0 &\neq (-2, -2, 4)\end{aligned}$$

where the cross product of parallel vectors $\mathbf{A} \times \mathbf{B} = 0$. Therefore, the cross product is not associative.

1.3 Taking the dot product of a unit cube's body diagonals $\mathbf{A} = (1, 1, 1)$, $\mathbf{B} = (1, 1, -1)$:

$$\begin{aligned}\mathbf{A} \cdot \mathbf{B} &= AB \cos \theta \\ 1 &= 3 \cos \theta \\ \theta &= \arccos 1/3 \approx 70.53^\circ\end{aligned}$$

1.4 The cross product of two vectors coplanar to the shaded plane— $\mathbf{A} = (-1, 2, 0)$, $\mathbf{B} = (-1, 0, 3)$ —is parallel to the normal unit vector $\hat{\mathbf{n}}$ of the plane:

$$\begin{aligned}\mathbf{A} \times \mathbf{B} &= \mathbf{C} \\ \begin{vmatrix} \hat{\mathbf{x}} & \hat{\mathbf{y}} & \hat{\mathbf{z}} \\ -1 & 2 & 0 \\ -1 & 0 & 3 \end{vmatrix} &= (6, 3, 2)\end{aligned}$$

where $\hat{\mathbf{n}} = \mathbf{C}/C$, and $C = \sqrt{6^2 + 3^2 + 2^2} = \sqrt{49} = 7$. Therefore,

$$\hat{\mathbf{n}} = \frac{1}{7}(6, 3, 2)$$

1.5 Proving the “BAC–CAB” rule for three-dimensional vectors:

$$\begin{aligned}\mathbf{A} \times (\mathbf{B} \times \mathbf{C}) &= \mathbf{A} \times \begin{vmatrix} \hat{\mathbf{x}} & \hat{\mathbf{y}} & \hat{\mathbf{z}} \\ B_x & B_y & B_z \\ C_x & C_y & C_z \end{vmatrix} \\ &= \begin{vmatrix} \hat{\mathbf{x}} & \hat{\mathbf{y}} & \hat{\mathbf{z}} \\ A_x & A_y & A_z \\ B_y C_z - B_z C_y & B_z C_x - B_x C_z & B_x C_y - B_y C_x \end{vmatrix}\end{aligned}$$

where the x component is

$$A_y(B_x C_y - B_y C_x) - A_z(B_z C_x - B_x C_z)$$

From the “BAC–CAB” rule,

$$\mathbf{B}(\mathbf{A} \cdot \mathbf{C}) - \mathbf{C}(\mathbf{A} \cdot \mathbf{B}) = \mathbf{B}(A_x C_x + A_y C_y + A_z C_z) - \mathbf{C}(A_x B_x + A_y B_y + A_z B_z)$$

where x component simplifies to

$$B_x(A_x C_x + A_y C_y + A_z C_z) - C_x(A_x B_x + A_y B_y + A_z B_z) = A_y(B_x C_y - B_y C_x) - A_z(B_z C_x - B_x C_z)$$

So,

$$[\mathbf{A} \times (\mathbf{B} \times \mathbf{C})]_x = [\mathbf{B}(\mathbf{A} \cdot \mathbf{C}) - \mathbf{C}(\mathbf{A} \cdot \mathbf{B})]_x$$

and similiary for the y and z components. Therefore, the “BAC–CAB” rule holds true.

1.6

$$\begin{aligned}[\mathbf{A} \times (\mathbf{B} \times \mathbf{C})] + [\mathbf{B} \times (\mathbf{C} \times \mathbf{A})] + [\mathbf{C} \times (\mathbf{A} \times \mathbf{B})] &= \cancel{\mathbf{B}(\mathbf{A} \cdot \mathbf{C})} - \mathbf{C}(\mathbf{A} \cdot \mathbf{B}) + \mathbf{C}(\mathbf{B} \cdot \mathbf{A}) = 0 \\ &= -\mathbf{A}(\mathbf{B} \cdot \mathbf{C}) + \mathbf{A}(\mathbf{C} \cdot \mathbf{B}) - \cancel{\mathbf{B}(\mathbf{C} \cdot \mathbf{A})}\end{aligned}$$

since dot product is associative, the first and last terms cancel out, and the middle terms also cancel out with each other.

$$\begin{aligned}\mathbf{A} \times (\mathbf{B} \times \mathbf{C}) &= (\mathbf{A} \times \mathbf{B}) \times \mathbf{C} \\ \mathbf{B}(\mathbf{A} \cdot \mathbf{C}) - \mathbf{C}(\mathbf{A} \cdot \mathbf{B}) &= \mathbf{B}(\mathbf{A} \cdot \mathbf{C}) - \mathbf{A}(\mathbf{B} \cdot \mathbf{C}) \\ 0 &= -\mathbf{C}(\mathbf{A} \cdot \mathbf{B}) + \mathbf{A}(\mathbf{B} \cdot \mathbf{C}) \\ 0 &= -\mathbf{B}(\mathbf{C} \times \mathbf{A}) = (\mathbf{C} \times \mathbf{A}) \times \mathbf{B}\end{aligned}$$

For the relation to hold true, either the vectors \mathbf{A} and \mathbf{C} are parallel ($\mathbf{A} \times \mathbf{C} = 0$) or \mathbf{B} is perpendicular to both \mathbf{A} and \mathbf{C} ($\mathbf{A} \cdot \mathbf{B} = \mathbf{B} \cdot \mathbf{C} = 0$).

1.7 Finding the separation vector \mathbf{z} :

$$\begin{aligned}\mathbf{z} &= \mathbf{r} - \mathbf{r}' = (4, 6, 8) - (2, 8, 7) = (2, -2, 1) \\ z &= \sqrt{2^2 + (-2)^2 + 1^2} = 3 \\ \hat{\mathbf{z}} &= \left(\frac{2}{3}, -\frac{2}{3}, \frac{1}{3} \right)\end{aligned}$$

1.8 (a)

$$\begin{aligned}\bar{A}_y \bar{B}_y + \bar{A}_z \bar{B}_z &= (A_y \cos \phi + A_z \sin \phi)(B_y \cos \phi + B_z \sin \phi) \\ &\quad + (-A_y \sin \phi + A_z \cos \phi)(-B_y \sin \phi + B_z \cos \phi) \\ &= A_y B_y \cos^2 \phi + A_z B_z \sin^2 \phi + \cancel{A_y B_z \sin \phi \cos \phi} + \cancel{A_z B_y \sin \phi \cos \phi} \\ &\quad + A_y B_y \sin^2 \phi - \cancel{A_y B_z \sin \phi \cos \phi} - \cancel{A_z B_y \sin \phi \cos \phi} + A_z B_z \cos^2 \phi \\ &= A_y B_y (\sin^2 \phi + \cos^2 \phi) + A_z B_z (\sin^2 \phi + \cos^2 \phi) \\ \bar{A}_y \bar{B}_y + \bar{A}_z \bar{B}_z &= A_y B_y + A_z B_z\end{aligned}$$

(b) To preserve length $|\bar{\mathbf{A}}| = |\mathbf{A}|$. Squaring both sides and expanding gives

$$\bar{A}_x^2 + \bar{A}_y^2 + \bar{A}_z^2 = A_x^2 + A_y^2 + A_z^2$$

in summation form,

$$\sum_{i=1}^3 \bar{A}_i \bar{A}_i = \sum_{i=1}^3 \left(\sum_{j=1}^3 R_{ij} A_j \right) \left(\sum_{k=1}^3 R_{ik} A_k \right) = \sum_{i=1}^3 \sum_{j=1}^3 \sum_{k=1}^3 R_{ij} R_{ik} A_j A_k$$

For the length to be preserved, the indices j and k must be equal. Therefore,

$$R_{ij} R_{ik} = \delta_{jk}$$

where δ_{ij} is the Kronecker delta. Thus the length is preserved if the rotation matrix is orthogonal, i.e.

$$R_{ij} R_{ik} = (R^T)_{ji} R_{ik} = \delta_{jk} \quad \text{or} \quad R^T R = I$$




Figure 1.2: Rotation of 120° about an axis through the origin and point $(1, 1, 1)$

1.9 From Figure 1.2, the rotation is equivalent to changing the position of the basis vectors $\hat{\mathbf{x}} \rightarrow \hat{\mathbf{z}}$, $\hat{\mathbf{y}} \rightarrow \hat{\mathbf{x}}$, and $\hat{\mathbf{z}} \rightarrow \hat{\mathbf{y}}$. Therefore, the rotation matrix is a permutation matrix:

$$R = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}$$

1.10 (a) Under a **translation** of coordinates $\bar{y} = y - a$, the origin O and terminus $A = (x, y, z)$ of some vector are translated to

$$\begin{aligned} O &\rightarrow O' = (0, -a, 0) \\ A &\rightarrow A' = (x, y - a, z) \end{aligned}$$

therefore, the translated vector is

$$\overline{O'A'} = (x, y - a, z) - (0, -a, 0) = (x, y, z) = \overline{OA} = \mathbf{A}$$

which is the same as the original vector, so

$$\begin{pmatrix} \bar{A}_x \\ \bar{A}_y \\ \bar{A}_z \end{pmatrix} = \begin{pmatrix} A_x \\ A_y \\ A_z \end{pmatrix}$$

(b) Under **inversion** of coordinates, only the terminus changes

$$\begin{aligned} O &\rightarrow O' = (0, 0, 0) \\ A &\rightarrow A' = (-x, -y, -z) \end{aligned}$$

therefore, the inverted vector is

$$\overline{O'A'} = (-x, -y, -z) \quad \text{or} \quad \begin{pmatrix} \bar{A}_x \\ \bar{A}_y \\ \bar{A}_z \end{pmatrix} = \begin{pmatrix} -A_x \\ -A_y \\ -A_z \end{pmatrix}$$

(c) The components of cross product under inversion

$$\bar{\mathbf{A}} \times \bar{\mathbf{B}} = \begin{pmatrix} \bar{A}_y \bar{B}_z - \bar{A}_z \bar{B}_y \\ \bar{A}_z \bar{B}_x - \bar{A}_x \bar{B}_z \\ \bar{A}_x \bar{B}_y - \bar{A}_y \bar{B}_x \end{pmatrix} = \begin{pmatrix} A_y B_z - A_z B_y \\ A_z B_x - A_x B_z \\ A_x B_y - A_y B_x \end{pmatrix}$$

which is the same as the original cross product $\mathbf{A} \times \mathbf{B}$. The cross product of two pseudovectors is also a pseudovector. Torque $\tau = \mathbf{r} \times \mathbf{F}$ and magnetic force $\mathbf{F} = q\mathbf{v} \times \mathbf{B}$ are examples of pseudovectors.

(d) Scalar triple product under inversion

$$\begin{aligned} \bar{\mathbf{A}} \cdot (\bar{\mathbf{B}} \times \bar{\mathbf{C}}) &= -\mathbf{A} \cdot (-\mathbf{B} \times -\mathbf{C}) \\ &= -\mathbf{A} \cdot (\mathbf{B} \times \mathbf{C}) \end{aligned}$$

the scalar triple product changes sign under inversion.

1.11 (a) Finding gradient of $f(x, y, z) = x^2 + y^3 + z^4$:

$$\begin{aligned} \nabla f &= \frac{\partial f}{\partial x} \hat{\mathbf{x}} + \frac{\partial f}{\partial y} \hat{\mathbf{y}} + \frac{\partial f}{\partial z} \hat{\mathbf{z}} \\ &= 2x \hat{\mathbf{x}} + 3y^2 \hat{\mathbf{y}} + 4z^3 \hat{\mathbf{z}} \end{aligned}$$

(b) Gradient of $f(x, y, z) = x^2 y^3 z^4$:

$$\nabla f = 2xy^3z^4 \hat{\mathbf{x}} + 3x^2y^2z^4 \hat{\mathbf{y}} + 4x^2y^3z^3 \hat{\mathbf{z}}$$

(c) Gradient of $f(x, y, z) = e^x \sin(y) \ln(z)$:

$$\nabla f = e^x \hat{\mathbf{x}} + e^x \cos(y) \ln(z) \hat{\mathbf{y}} + \frac{e^x \sin(y)}{z} \hat{\mathbf{z}}$$

1.12 The height of the hill (in feet) is given by the function

$$h(x, y) = 10(2xy - 3x^2 - 4y^2 - 18x + 28y + 12)$$

where y is north and x is east in miles. The gradient of h is

$$\nabla h = 10(2y - 6x - 18)\hat{\mathbf{x}} + 10(2x - 8y + 28)\hat{\mathbf{y}}$$

(a) The top of the hill is a stationary point, so the summit is found by setting the gradient to zero which gives the system of equations

$$0 = 2y - 6x - 18 \qquad 0 = 2x - 8y + 28$$

adding the first equation to 3 times the second equation gives

$$\begin{aligned} 0 &= 2y - 6x - 18 + 3(2x - 8y + 28) \\ 0 &= -22y + 66 \\ y &= 3 \end{aligned}$$

substituting $y = 3$ into the first equation

$$0 = 2(3) - 6x - 18 \rightarrow x = -2$$

Therefore, the top of the hill is at $(-2, 3)$ or 2 miles west and 3 miles north of the origin.

(b) The height of the hill is simply $h(-2, 3) = 10(12) = 720$ feet.

(c) The steepness of the hill at $h(1, 1)$ is given by the magnitude of the gradient

$$\begin{aligned} |\nabla h| &= 10\sqrt{(2y - 6x - 18)^2 + (2x - 8y + 28)^2} \\ &= 10\sqrt{(2 - 6 - 18)^2 + (2 - 8 + 28)^2} \\ &= 10\sqrt{(-22)^2 + (22)^2} = 220\sqrt{2} \approx 311 \text{ ft/mi} \end{aligned}$$

The direction of the steepest slope is given by the vector in the direction of the gradient at the point $\nabla h(1, 1) = 220(-\mathbf{x} + \mathbf{y})$, or simply northwest.

1.13 Given the separation vector

$$\mathbf{z} = (x - x')\hat{\mathbf{x}} + (y - y')\hat{\mathbf{y}} + (z - z')\hat{\mathbf{z}} \quad \text{and} \quad z = \sqrt{(x - x')^2 + (y - y')^2 + (z - z')^2}$$

(a) Show that $\nabla(z^2) = 2\mathbf{z}$:

$$z^2 = (x - x')^2 + (y - y')^2 + (z - z')^2$$

and each component of the gradient is

$$\begin{aligned} \frac{\partial}{\partial x}(z^2) &= 2(x - x') \\ \frac{\partial}{\partial y}(z^2) &= 2(y - y') \\ \frac{\partial}{\partial z}(z^2) &= 2(z - z') \end{aligned}$$

so

$$\nabla(z^2) = 2\mathbf{z}$$

(b) Show $\nabla(1/z) = -\hat{\mathbf{z}}/z^2$:

$$\frac{1}{z} = [(x - x')^2 + (y - y')^2 + (z - z')^2]^{-1/2} = [(\quad)]^{-1/2}$$

looking at the x component of the gradient (using chain rule),

$$\begin{aligned}\frac{\partial}{\partial x}\left(\frac{1}{z}\right) &= -\frac{1}{2}[(\)]^{-3/2} \cdot 2(x-x') \\ &= -\frac{(x-x')}{z^3}\end{aligned}$$

and similarly for the y and z components:

$$\begin{aligned}\nabla\left(\frac{1}{z}\right) &= \frac{\partial}{\partial x}\left(\frac{1}{z}\right)\hat{\mathbf{x}} + \frac{\partial}{\partial y}\left(\frac{1}{z}\right)\hat{\mathbf{y}} + \frac{\partial}{\partial z}\left(\frac{1}{z}\right)\hat{\mathbf{z}} \\ &= -\frac{1}{z^3}[(x-x')\hat{\mathbf{x}} + (y-y')\hat{\mathbf{y}} + (z-z')\hat{\mathbf{z}}] = -\frac{\mathbf{z}}{z^3}\end{aligned}$$

Finally, substituting the unit vector $\hat{\mathbf{z}} = \mathbf{z}/z$ gives us

$$\nabla\left(\frac{1}{z}\right) = -\frac{\hat{\mathbf{z}}}{z^2}$$

(c) The general formula for $\nabla(z^n)$:

$$\nabla(z^n) = n z^{n-1} \cdot \nabla(z)$$

where

$$\begin{aligned}\nabla(z) &= \frac{\partial}{\partial x}(z)\hat{\mathbf{x}} + \frac{\partial}{\partial y}(z)\hat{\mathbf{y}} + \frac{\partial}{\partial z}(z)\hat{\mathbf{z}} \\ &= \frac{1}{2}[(\)]^{-1/2} \cdot 2(x-x')\hat{\mathbf{x}} + \dots \quad [\text{similar to part (b)}] \\ &= \frac{\mathbf{z}}{z} = \hat{\mathbf{z}}\end{aligned}$$

So the general formula is

$$\nabla(z^n) = n z^{n-1} \hat{\mathbf{z}}$$

1.14 Given the rotation matrix

$$\begin{pmatrix} \bar{y} \\ \bar{z} \end{pmatrix} = \begin{pmatrix} \cos \phi & \sin \phi \\ -\sin \phi & \cos \phi \end{pmatrix} \begin{pmatrix} y \\ z \end{pmatrix}$$

or the two equations

$$\begin{aligned}\bar{y} &= y \cos \phi + z \sin \phi \\ \bar{z} &= -y \sin \phi + z \cos \phi\end{aligned}$$

differentiating with respect to \bar{y} and \bar{z} respectively gives

$$\begin{aligned}1 &= \frac{\partial y}{\partial \bar{y}} \cos \phi + \frac{\partial z}{\partial \bar{y}} \sin \phi \\ 1 &= -\frac{\partial y}{\partial \bar{z}} \sin \phi + \frac{\partial z}{\partial \bar{z}} \cos \phi\end{aligned}$$

this can only be true if

$$\frac{\partial y}{\partial \bar{y}} = \cos \phi, \quad \frac{\partial z}{\partial \bar{y}} = \sin \phi \quad \text{and} \quad \frac{\partial y}{\partial \bar{z}} = -\sin \phi, \quad \frac{\partial z}{\partial \bar{z}} = \cos \phi$$

which satisfies the trig identity $\sin^2 \phi + \cos^2 \phi = 1$. Differentiating f with respect to the rotated coordinates \bar{y} and \bar{z} is given by

$$\begin{aligned}\frac{\partial f}{\partial \bar{y}} &= \frac{\partial f}{\partial y} \frac{\partial y}{\partial \bar{y}} + \frac{\partial f}{\partial z} \frac{\partial z}{\partial \bar{y}} = \frac{\partial f}{\partial y} \cos \phi + \frac{\partial f}{\partial z} \sin \phi \\ \frac{\partial f}{\partial \bar{z}} &= \frac{\partial f}{\partial y} \frac{\partial y}{\partial \bar{z}} + \frac{\partial f}{\partial z} \frac{\partial z}{\partial \bar{z}} = -\frac{\partial f}{\partial y} \sin \phi + \frac{\partial f}{\partial z} \cos \phi\end{aligned}$$

therefore, the gradient of f transforms as a vector under rotations given by

$$\overline{\nabla}f = \frac{\partial f}{\partial \bar{y}}\hat{\mathbf{y}} + \frac{\partial f}{\partial \bar{z}}\hat{\mathbf{z}} = \left(\frac{\partial f}{\partial y}\cos\phi + \frac{\partial f}{\partial z}\sin\phi\right)\hat{\mathbf{y}} + \left(-\frac{\partial f}{\partial y}\sin\phi + \frac{\partial f}{\partial z}\cos\phi\right)\hat{\mathbf{z}}$$

or in matrix form

$$\overline{\nabla}f = \begin{pmatrix} \cos\phi & -\sin\phi \\ \sin\phi & \cos\phi \end{pmatrix} \nabla f$$

where the gradient is a column vector

$$\nabla f = \begin{pmatrix} \frac{\partial f}{\partial y} \\ \frac{\partial f}{\partial z} \end{pmatrix}$$

1.15 (a) Calculating divergence of $\mathbf{v}_a = x^2\hat{\mathbf{x}} + 3xz^2\hat{\mathbf{y}} - 2xz\hat{\mathbf{z}}$:

$$\begin{aligned} \nabla \cdot \mathbf{v}_a &= \frac{\partial v_{ax}}{\partial x} + \frac{\partial v_{ay}}{\partial y} + \frac{\partial v_{az}}{\partial z} \\ &= 2x + 0 - 2x = 0 \end{aligned}$$

(b) $\mathbf{v}_b = xy\hat{\mathbf{x}} + 2yz\hat{\mathbf{y}} + 3zx\hat{\mathbf{z}}$:

$$\nabla \cdot \mathbf{v}_b = y + 2z + 3x$$

(c) $\mathbf{v}_c = y^2\hat{\mathbf{x}} + (2xy + z^2)\hat{\mathbf{y}} + 2yz\hat{\mathbf{z}}$:

$$\nabla \cdot \mathbf{v}_c = 0 + 2x + 2y = 2(x + y)$$




Figure 1.3: Sketch of the vector field $\mathbf{v} = \hat{\mathbf{r}}/r^2$

1.16 Given

$$r = \sqrt{x^2 + y^2 + z^2} \quad \text{and} \quad \hat{\mathbf{r}} = \frac{\mathbf{r}}{r} = \frac{x\hat{\mathbf{x}} + y\hat{\mathbf{y}} + z\hat{\mathbf{z}}}{\sqrt{x^2 + y^2 + z^2}}$$

where $\mathbf{r} = x\hat{\mathbf{x}} + y\hat{\mathbf{y}} + z\hat{\mathbf{z}}$ is the position vector. The vector functions is

$$v = \frac{\hat{\mathbf{r}}}{r^2} = \frac{\mathbf{r}}{r^3}$$

Finally adding the two equations together gives

$$\begin{aligned}
\nabla \cdot \bar{\mathbf{v}} &= \frac{\partial \bar{v}_y}{\partial y} + \frac{\partial \bar{v}_z}{\partial z} \\
&= \frac{\partial v_y}{\partial y} \cos^2 \phi + \frac{\partial v_y}{\partial z} \sin \phi \cos \phi + \frac{\partial v_z}{\partial y} \sin \phi \cos \phi + \frac{\partial v_z}{\partial z} \sin^2 \phi \\
&\quad + \frac{\partial v_y}{\partial y} \sin^2 \phi - \frac{\partial v_y}{\partial z} \sin \phi \cos \phi - \frac{\partial v_z}{\partial y} \sin \phi \cos \phi + \frac{\partial v_z}{\partial z} \cos^2 \phi \\
&= (\sin^2 \phi + \cos^2 \phi) \left[\frac{\partial v_y}{\partial y} + \frac{\partial v_z}{\partial z} \right] \\
&= \frac{\partial v_y}{\partial y} + \frac{\partial v_z}{\partial z}
\end{aligned}$$

which shows that the divergence transforms as a scalar under rotations.

1.18 Curl of vector functions from Problem 1.15: (a) $\mathbf{v}_a = x^2 \hat{\mathbf{x}} + 3xz^2 \hat{\mathbf{y}} - 2xz \hat{\mathbf{z}}$:

$$\begin{aligned}
\nabla \times \mathbf{v}_a &= \begin{vmatrix} \hat{\mathbf{x}} & \hat{\mathbf{y}} & \hat{\mathbf{z}} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ x^2 & 3xz^2 & -2xz \end{vmatrix} \\
&= \hat{\mathbf{x}}(0 - 6xz) - \hat{\mathbf{y}}(-2z - 0) + \hat{\mathbf{z}}(3z^2 - 0) \\
&= -6xz \hat{\mathbf{x}} + 2z \hat{\mathbf{y}} + 3z^2 \hat{\mathbf{z}}
\end{aligned}$$

(b) $\mathbf{v}_b = xy \hat{\mathbf{x}} + 2yz \hat{\mathbf{y}} + 3zx \hat{\mathbf{z}}$:

$$\begin{aligned}
\nabla \times \mathbf{v}_b &= \begin{vmatrix} \hat{\mathbf{x}} & \hat{\mathbf{y}} & \hat{\mathbf{z}} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ xy & 2yz & 3zx \end{vmatrix} \\
&= \hat{\mathbf{x}}(0 - 2y) - \hat{\mathbf{y}}(3z - 0) + \hat{\mathbf{z}}(0 - x) \\
&= -2y \hat{\mathbf{x}} - 3z \hat{\mathbf{y}} - x \hat{\mathbf{z}}
\end{aligned}$$

(c) $\mathbf{v}_c = y^2 \hat{\mathbf{x}} + (2xy + z^2) \hat{\mathbf{y}} + 2yz \hat{\mathbf{z}}$:

$$\begin{aligned}
\nabla \times \mathbf{v}_c &= \begin{vmatrix} \hat{\mathbf{x}} & \hat{\mathbf{y}} & \hat{\mathbf{z}} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ y^2 & 2xy + z^2 & 2yz \end{vmatrix} \\
&= \hat{\mathbf{x}}(2z - 2z) - \hat{\mathbf{y}}(0 - 0) + \hat{\mathbf{z}}(2y - 2y) \\
&= 0
\end{aligned}$$

1.19 From Figure 1.4, the sign of $\partial v_x / \partial y$ is positive, and the sign of $\partial v_y / \partial x$ is negative. Therefore, the curl

$$\nabla \times \mathbf{v} = \hat{\mathbf{z}} \left(\frac{\partial v_x}{\partial y} - \frac{\partial v_y}{\partial x} \right)$$

is in the negative z direction, or into the page. This is consistent with the right-hand rule as the thumb points into the page.

1.20 Proof that the vector function

$$\mathbf{g} = \frac{\hat{\mathbf{r}}}{r^2} = \frac{\mathbf{r}}{r^3} = \frac{x \hat{\mathbf{x}} + y \hat{\mathbf{y}} + z \hat{\mathbf{z}}}{r^3}$$

has zero divergence and curl given

$$\frac{\partial r}{\partial x_i} = \frac{x_i}{r} \quad \text{and} \quad \frac{\partial x}{\partial y} = \frac{y}{x} = 0$$

(iv)

$$\begin{aligned}
\nabla \cdot (\mathbf{A} \times \mathbf{B}) &= \nabla \cdot \begin{vmatrix} \hat{\mathbf{x}} & \hat{\mathbf{y}} & \hat{\mathbf{z}} \\ A_x & A_y & A_z \\ B_x & B_y & B_z \end{vmatrix} \\
&= \nabla \cdot [(A_y B_z - A_z B_y)\hat{\mathbf{x}} + (A_z B_x - A_x B_z)\hat{\mathbf{y}} + (A_x B_y - A_y B_x)\hat{\mathbf{z}}] \\
&= \frac{\partial}{\partial x}(A_y B_z - A_z B_y) + \frac{\partial}{\partial y}(A_z B_x - A_x B_z) + \frac{\partial}{\partial z}(A_x B_y - A_y B_x) \\
&= \left(\frac{\partial A_y}{\partial x} B_z + A_y \frac{\partial B_z}{\partial x} - \frac{\partial A_z}{\partial x} B_y - A_z \frac{\partial B_y}{\partial x} \right) + \left(\frac{\partial A_z}{\partial y} B_x + A_z \frac{\partial B_x}{\partial y} - \frac{\partial A_x}{\partial y} B_z - A_x \frac{\partial B_z}{\partial y} \right) \\
&\quad + \left(\frac{\partial A_x}{\partial z} B_y + A_x \frac{\partial B_y}{\partial z} - \frac{\partial A_y}{\partial z} B_x - A_y \frac{\partial B_x}{\partial z} \right) \\
&= B_x \left(\frac{\partial A_z}{\partial y} - \frac{\partial A_y}{\partial z} \right) + B_y \left(\frac{\partial A_x}{\partial z} - \frac{\partial A_z}{\partial x} \right) + B_z \left(\frac{\partial A_y}{\partial x} - \frac{\partial A_x}{\partial y} \right) \\
&\quad + A_x \left(\frac{\partial B_y}{\partial z} - \frac{\partial B_z}{\partial y} \right) + A_y \left(\frac{\partial B_z}{\partial x} - \frac{\partial B_x}{\partial z} \right) + A_z \left(\frac{\partial B_x}{\partial y} - \frac{\partial B_y}{\partial x} \right) \\
&= \mathbf{B} \cdot (\nabla \times \mathbf{A}) + \mathbf{A} \cdot (\nabla \times \mathbf{B})
\end{aligned}$$

(v)

$$\begin{aligned}
\nabla \times (f\mathbf{A}) &= \begin{vmatrix} \hat{\mathbf{x}} & \hat{\mathbf{y}} & \hat{\mathbf{z}} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ fA_x & fA_y & fA_z \end{vmatrix} \\
&= \hat{\mathbf{x}} \left(\frac{\partial}{\partial y}(fA_z) - \frac{\partial}{\partial z}(fA_y) \right) - \hat{\mathbf{y}} \left(\frac{\partial}{\partial x}(fA_z) - \frac{\partial}{\partial z}(fA_x) \right) + \hat{\mathbf{z}} \left(\frac{\partial}{\partial x}(fA_y) - \frac{\partial}{\partial y}(fA_x) \right) \\
&= \hat{\mathbf{x}} \left(f \frac{\partial A_z}{\partial y} + A_z \frac{\partial f}{\partial y} - f \frac{\partial A_y}{\partial z} - A_y \frac{\partial f}{\partial z} \right) - \hat{\mathbf{y}} \left(f \frac{\partial A_z}{\partial x} + A_z \frac{\partial f}{\partial x} - f \frac{\partial A_x}{\partial z} - A_x \frac{\partial f}{\partial z} \right) \\
&\quad + \hat{\mathbf{z}} \left(f \frac{\partial A_y}{\partial x} + A_y \frac{\partial f}{\partial x} - f \frac{\partial A_x}{\partial y} - A_x \frac{\partial f}{\partial y} \right) \\
&= f \left[\hat{\mathbf{x}} \left(\frac{\partial A_z}{\partial y} - \frac{\partial A_y}{\partial z} \right) + \hat{\mathbf{y}} \left(\frac{\partial A_x}{\partial z} - \frac{\partial A_z}{\partial x} \right) + \hat{\mathbf{z}} \left(\frac{\partial A_y}{\partial x} - \frac{\partial A_x}{\partial y} \right) \right] \\
&\quad - \hat{\mathbf{x}} \left(A_y \frac{\partial f}{\partial z} - A_z \frac{\partial f}{\partial y} \right) + \hat{\mathbf{y}} \left(A_z \frac{\partial f}{\partial x} - A_x \frac{\partial f}{\partial z} \right) - \hat{\mathbf{z}} \left(A_x \frac{\partial f}{\partial y} - A_y \frac{\partial f}{\partial x} \right) \\
&= f(\nabla \times \mathbf{A}) - \mathbf{A} \times (\nabla f)
\end{aligned}$$

1.22 (a) If \mathbf{A} and \mathbf{B} are two vector functions, then

$$\begin{aligned}
(\mathbf{A} \cdot \nabla)\mathbf{B} &= \left(A_x \frac{\partial}{\partial x} + A_y \frac{\partial}{\partial y} + A_z \frac{\partial}{\partial z} \right) (B_x \hat{\mathbf{x}} + B_y \hat{\mathbf{y}} + B_z \hat{\mathbf{z}}) \\
&= \hat{\mathbf{x}} \left(A_x \frac{\partial B_x}{\partial x} + A_y \frac{\partial B_x}{\partial y} + A_z \frac{\partial B_x}{\partial z} \right) + \hat{\mathbf{y}} \left(A_x \frac{\partial B_y}{\partial x} + A_y \frac{\partial B_y}{\partial y} + A_z \frac{\partial B_y}{\partial z} \right) \\
&\quad + \hat{\mathbf{z}} \left(A_x \frac{\partial B_z}{\partial x} + A_y \frac{\partial B_z}{\partial y} + A_z \frac{\partial B_z}{\partial z} \right)
\end{aligned}$$

This means that the direction of \mathbf{A} points in the direction of where \mathbf{B} moves fastest.

(b)

$$(\hat{\mathbf{r}} \cdot \nabla)\hat{\mathbf{r}} = \frac{1}{r} \left(x \frac{\partial}{\partial x} + y \frac{\partial}{\partial y} + z \frac{\partial}{\partial z} \right) \frac{(x\hat{\mathbf{x}} + y\hat{\mathbf{y}} + z\hat{\mathbf{z}})}{r}$$

looking at the x component,

$$\begin{aligned}\frac{\partial}{\partial x}\left(\frac{\mathbf{r}}{r}\right) &= \frac{1}{r}\frac{\partial}{\partial x}(x\hat{\mathbf{x}} + y\hat{\mathbf{y}} + z\hat{\mathbf{z}}) + \mathbf{r}\frac{\partial}{\partial x}\left(\frac{1}{r}\right) \\ &= \frac{\hat{\mathbf{x}}}{r} - \mathbf{r}\frac{x^2}{r^3}\end{aligned}$$

therefore,

$$\begin{aligned}(\hat{\mathbf{r}} \cdot \nabla)\hat{\mathbf{r}} &= \frac{1}{r}\left[x\left(\frac{\hat{\mathbf{x}}}{r} - \mathbf{r}\frac{x}{r^3}\right) + y\left(\frac{\hat{\mathbf{y}}}{r} - \mathbf{r}\frac{y}{r^3}\right) + z\left(\frac{\hat{\mathbf{z}}}{r} - \mathbf{r}\frac{z}{r^3}\right)\right] \\ &= \frac{1}{r}\left[\frac{x\hat{\mathbf{x}} + y\hat{\mathbf{y}} + z\hat{\mathbf{z}}}{r} - \mathbf{r}\frac{x^2 + y^2 + z^2}{r^3}\right] \\ &= \frac{1}{r}\left[\frac{\mathbf{r}}{r} - \frac{\mathbf{r}}{r}\right] = 0\end{aligned}$$

(c)

$$\begin{aligned}(v_a \cdot \nabla)v_b &= \left(x^2\frac{\partial}{\partial x} + 3xz^2\frac{\partial}{\partial y} - 2xz\frac{\partial}{\partial z}\right)(xy\hat{\mathbf{x}} + 2yz\hat{\mathbf{y}} + 3zx\hat{\mathbf{z}}) \\ &= x^2(y\hat{\mathbf{x}} + 0 + 3z\hat{\mathbf{z}}) + 3xz^2(x\hat{\mathbf{x}} + 2z\hat{\mathbf{y}} + 0) - 2xz(0 + 2y\hat{\mathbf{y}} + 3x\hat{\mathbf{z}}) \\ &= (x^2y + 3x^2z^2)\hat{\mathbf{x}} + (6xz^3 - 4xyz)\hat{\mathbf{y}} + (3x^2z - 6x^2z)\hat{\mathbf{z}} \\ &= x^2(y + 3z^2)\hat{\mathbf{x}} + 2xz(3z^2 - 2y)\hat{\mathbf{y}} - 3x^2z\hat{\mathbf{z}}\end{aligned}$$

1.23 Proving the product rule for (ii) given the x component of the left hand side is

$$\begin{aligned}[\nabla(\mathbf{A} \cdot \mathbf{B})]_x &= \frac{\partial(\mathbf{A} \cdot \mathbf{B})}{\partial x}\hat{\mathbf{x}} \\ &= \frac{\partial}{\partial x}(A_xB_x + A_yB_y + A_zB_z)\hat{\mathbf{x}} \\ &= A_x\frac{\partial B_x}{\partial x} + B_x\frac{\partial A_x}{\partial x} + A_y\frac{\partial B_y}{\partial x} + B_y\frac{\partial A_y}{\partial x} + A_z\frac{\partial B_z}{\partial x} + B_z\frac{\partial A_z}{\partial x}\end{aligned}$$

Finding the x component of the right hand side of (ii)

$$\begin{aligned}[\mathbf{A} \times (\nabla \times \mathbf{B})]_x &= \left[\mathbf{A} \times \begin{bmatrix} \hat{\mathbf{x}} & \hat{\mathbf{y}} & \hat{\mathbf{z}} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ B_x & B_y & B_z \end{bmatrix}\right]_x \\ &= \left[\begin{bmatrix} \hat{\mathbf{x}} & \hat{\mathbf{y}} & \hat{\mathbf{z}} \\ A_x & A_y & A_z \\ () & -(\frac{\partial B_z}{\partial x} - \frac{\partial B_x}{\partial z}) & \frac{\partial B_y}{\partial x} - \frac{\partial B_x}{\partial y} \end{bmatrix}\right]_x \\ &= A_y\left(\frac{\partial B_y}{\partial x} - \frac{\partial B_x}{\partial y}\right) - A_z\left(\frac{\partial B_x}{\partial z} - \frac{\partial B_z}{\partial x}\right)\end{aligned}$$

and

$$[\mathbf{B} \times (\nabla \times \mathbf{A})]_x = B_y\left(\frac{\partial A_y}{\partial x} - \frac{\partial A_x}{\partial y}\right) - B_z\left(\frac{\partial A_x}{\partial z} - \frac{\partial A_z}{\partial x}\right)$$

From Problem 1.22:

$$[(\mathbf{A} \cdot \nabla)\mathbf{B}] = A_x\frac{\partial B_x}{\partial x} + A_y\frac{\partial B_x}{\partial y} + A_z\frac{\partial B_x}{\partial z}$$

and likewise,

$$[(\mathbf{B} \cdot \nabla)\mathbf{A}] = B_x\frac{\partial A_x}{\partial x} + B_y\frac{\partial A_x}{\partial y} + B_z\frac{\partial A_x}{\partial z}$$

Adding all four equations gives

$$\begin{aligned}
[\mathbf{A} \times (\nabla \times \mathbf{B}) + \mathbf{B} \times (\nabla \times \mathbf{A}) + (\mathbf{A} \cdot \nabla)\mathbf{B} + (\mathbf{B} \cdot \nabla)\mathbf{A}]_x = \\
A_y \left(\frac{\partial B_y}{\partial x} - \frac{\partial B_x}{\partial y} \right) - A_z \left(\frac{\partial B_x}{\partial z} - \frac{\partial B_z}{\partial x} \right) + B_y \left(\frac{\partial A_y}{\partial x} - \frac{\partial A_x}{\partial y} \right) - B_z \left(\frac{\partial A_x}{\partial z} - \frac{\partial A_z}{\partial x} \right) \\
+ A_x \frac{\partial B_x}{\partial x} + A_y \frac{\partial B_x}{\partial y} + A_z \frac{\partial B_x}{\partial z} + B_x \frac{\partial A_x}{\partial x} + B_y \frac{\partial A_x}{\partial y} + B_z \frac{\partial A_x}{\partial z} \\
= A_x \frac{\partial B_x}{\partial x} + A_y \left(\frac{\partial B_y}{\partial x} - \cancel{\frac{\partial B_x}{\partial y}} + \cancel{\frac{\partial B_x}{\partial y}} \right) + A_z \left(\frac{\partial B_z}{\partial x} - \cancel{\frac{\partial B_x}{\partial z}} + \cancel{\frac{\partial B_x}{\partial z}} \right) \\
+ B_x \frac{\partial A_x}{\partial x} + B_y \left(\frac{\partial A_y}{\partial x} - \cancel{\frac{\partial A_x}{\partial y}} + \cancel{\frac{\partial A_x}{\partial y}} \right) + B_z \left(\frac{\partial A_z}{\partial x} - \cancel{\frac{\partial A_x}{\partial z}} + \cancel{\frac{\partial A_x}{\partial z}} \right) \\
= A_x \frac{\partial B_x}{\partial x} + B_x \frac{\partial A_x}{\partial x} + A_y \frac{\partial B_x}{\partial y} + B_y \frac{\partial A_x}{\partial y} + A_z \frac{\partial B_x}{\partial z} + B_z \frac{\partial A_x}{\partial z} \\
= [\nabla(\mathbf{A} \cdot \mathbf{B})]_x
\end{aligned}$$

and likewise for the y and z components.

For (vi), the x on the left hand side is

$$\begin{aligned}
[\nabla \times (\mathbf{A} \times \mathbf{B})]_x &= \left[\nabla \times \begin{bmatrix} \hat{\mathbf{x}} & \hat{\mathbf{y}} & \hat{\mathbf{z}} \\ A_x & A_y & A_z \\ B_x & B_y & B_z \end{bmatrix} \right]_x \\
&= \left[\begin{bmatrix} \hat{\mathbf{x}} & \hat{\mathbf{y}} & \hat{\mathbf{z}} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ () & -(A_x B_z - A_z B_x) & A_x B_y - A_y B_x \end{bmatrix} \right]_x \\
&= \frac{\partial}{\partial y} (A_x B_y - A_y B_x) - \frac{\partial}{\partial z} (A_z B_x - A_x B_z) \\
&= A_x \frac{\partial B_y}{\partial y} + B_y \frac{\partial A_x}{\partial y} - A_y \frac{\partial B_x}{\partial y} - B_x \frac{\partial A_y}{\partial y} \\
&\quad - A_z \frac{\partial B_x}{\partial z} - B_x \frac{\partial A_z}{\partial z} + A_x \frac{\partial B_z}{\partial z} + B_z \frac{\partial A_x}{\partial z} + \\
&= A_x \left(\frac{\partial B_y}{\partial y} + \frac{\partial B_z}{\partial z} \right) - A_y \frac{\partial B_x}{\partial y} - A_z \frac{\partial B_x}{\partial z} - B_x \left(\frac{\partial A_y}{\partial y} + \frac{\partial A_z}{\partial z} \right) + B_y \frac{\partial A_x}{\partial y} + B_z \frac{\partial A_x}{\partial z}
\end{aligned}$$

On the right hand side, first we find the x component of the two new operations:

$$\begin{aligned}
[A(\nabla \cdot \mathbf{B})]_x &= \left[A \left(\frac{\partial B_x}{\partial x} + \frac{\partial B_y}{\partial y} + \frac{\partial B_z}{\partial z} \right) \right]_x \\
&= A_x \left(\frac{\partial B_x}{\partial x} + \frac{\partial B_y}{\partial y} + \frac{\partial B_z}{\partial z} \right)
\end{aligned}$$

and likewise,

$$[B(\nabla \cdot \mathbf{A})]_x = B_x \left(\frac{\partial A_x}{\partial x} + \frac{\partial A_y}{\partial y} + \frac{\partial A_z}{\partial z} \right)$$

Therefore, $[(\mathbf{B} \cdot \nabla)\mathbf{A} - (\mathbf{A} \cdot \nabla)\mathbf{B} + A(\nabla \cdot \mathbf{B}) - B(\nabla \cdot \mathbf{A})]_x =$

$$\begin{aligned}
& B_x \frac{\partial A_x}{\partial x} + B_y \frac{\partial A_x}{\partial y} + B_z \frac{\partial A_x}{\partial z} - \left(A_x \frac{\partial B_x}{\partial x} + A_y \frac{\partial B_x}{\partial y} + A_z \frac{\partial B_x}{\partial z} \right) \\
& + A_x \left(\frac{\partial B_x}{\partial x} + \frac{\partial B_y}{\partial y} + \frac{\partial B_z}{\partial z} \right) - \left(B_x \left(\frac{\partial A_x}{\partial x} + \frac{\partial A_y}{\partial y} + \frac{\partial A_z}{\partial z} \right) \right) \\
& = A_x \left(\cancel{\frac{\partial B_x}{\partial x}} + \frac{\partial B_y}{\partial y} + \frac{\partial B_z}{\partial z} - \cancel{\frac{\partial B_x}{\partial x}} \right) - A_y \frac{\partial B_x}{\partial y} - A_z \frac{\partial B_x}{\partial z} \\
& - B_x \left(\cancel{\frac{\partial A_x}{\partial x}} + \frac{\partial A_y}{\partial y} + \frac{\partial A_z}{\partial z} - \cancel{\frac{\partial A_x}{\partial x}} \right) + B_y \frac{\partial A_x}{\partial y} + B_z \frac{\partial A_x}{\partial z} \\
& = A_x \left(\frac{\partial B_y}{\partial y} + \frac{\partial B_z}{\partial z} \right) - A_y \frac{\partial B_x}{\partial y} - A_z \frac{\partial B_x}{\partial z} - B_x \left(\frac{\partial A_y}{\partial y} + \frac{\partial A_z}{\partial z} \right) + B_y \frac{\partial A_x}{\partial y} + B_z \frac{\partial A_x}{\partial z} \\
& = [\nabla \times (\mathbf{A} \times \mathbf{B})]_x
\end{aligned}$$

and likewise for the y and z components.

1.24 Deriving the three quotient rules from the product rule: The gradient is

$$\begin{aligned}
\nabla \left(\frac{f}{g} \right) &= \nabla (fg^{-1}) = f\nabla(g^{-1}) + g^{-1}\nabla(f) \\
&= f(-g^{-2}\nabla(g)) + g^{-1}\nabla(f) \\
&= -\frac{f}{g^2}\nabla(g) + \frac{g}{g} \frac{1}{g}\nabla(f) \\
&= \frac{g\nabla f - f\nabla g}{g^2}
\end{aligned}$$

the divergence

$$\begin{aligned}
\nabla \cdot \left(\frac{\mathbf{A}}{g} \right) &= \nabla \cdot (Ag^{-1}) = A(\nabla \cdot g^{-1}) + g^{-1}(\nabla \cdot \mathbf{A}) \\
&= A(-g^{-2}(\nabla \cdot g)) + \frac{g}{g}g^{-1}(\nabla \cdot \mathbf{A}) \\
&= \frac{g(\nabla \cdot \mathbf{A}) - A\nabla \cdot g}{g^2}
\end{aligned}$$

and the curl

$$\begin{aligned}
\left[\nabla \times \left(\frac{\mathbf{A}}{g} \right) \right]_x &= \frac{\partial}{\partial y} \left(\frac{A_z}{g} \right) - \frac{\partial}{\partial z} \left(\frac{A_y}{g} \right) \\
&= \frac{g \frac{\partial A_z}{\partial y} - A_z \frac{\partial g}{\partial y}}{g^2} - \frac{g \frac{\partial A_y}{\partial z} - A_y \frac{\partial g}{\partial z}}{g^2} \\
&= \frac{1}{g^2} \left[g \left(\frac{\partial A_z}{\partial y} - \frac{\partial A_y}{\partial z} \right) + \left(A_y \frac{\partial g}{\partial z} - A_z \frac{\partial g}{\partial y} \right) \right] \\
&= \frac{g[\nabla \times \mathbf{A}]_x - \mathbf{A} \times [\nabla g]_x}{g^2}
\end{aligned}$$

and likewise for the y and z components. Therefore,

$$\nabla \times \left(\frac{\mathbf{A}}{g} \right) = \frac{g(\nabla \times \mathbf{A}) - \mathbf{A} \times (\nabla g)}{g^2}$$

1.25 (a) Calculating (iv) for the functions

$$\mathbf{A} = x\hat{\mathbf{x}} + 2y\hat{\mathbf{y}} + 3z\hat{\mathbf{z}}; \quad \mathbf{B} = 3y\hat{\mathbf{x}} - 2x\hat{\mathbf{y}}$$

LHS:

$$\begin{aligned}
\nabla \cdot (\mathbf{A} \times \mathbf{B}) &= \nabla \cdot \begin{vmatrix} \hat{\mathbf{x}} & \hat{\mathbf{y}} & \hat{\mathbf{z}} \\ x & 2y & 3z \\ 3y & -2x & 0 \end{vmatrix} \\
&= \nabla \cdot [(0 + 6xz)\hat{\mathbf{x}} - (0 - 9yz)\hat{\mathbf{y}} + (-2x^2 - 6y^2)\hat{\mathbf{z}}] \\
&= \frac{\partial}{\partial x}(6xz) + \frac{\partial}{\partial y}(9yz) + \frac{\partial}{\partial z}(-2x^2 - 6y^2) \\
&= 6z + 9z + 0 = 15z
\end{aligned}$$

RHS:

$$\begin{aligned}
\mathbf{B} \cdot (\nabla \times \mathbf{A}) &= \mathbf{B} \cdot \begin{vmatrix} \hat{\mathbf{x}} & \hat{\mathbf{y}} & \hat{\mathbf{z}} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ x & 2y & 3z \end{vmatrix} \\
&= \mathbf{B} \cdot [(0)\hat{\mathbf{x}} - (0)\hat{\mathbf{y}} + (0)\hat{\mathbf{z}}] = 0
\end{aligned}$$

and

$$\begin{aligned}
\mathbf{A} \cdot (\nabla \times \mathbf{B}) &= \mathbf{A} \cdot \begin{vmatrix} \hat{\mathbf{x}} & \hat{\mathbf{y}} & \hat{\mathbf{z}} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ 3y & -2x & 0 \end{vmatrix} \\
&= \mathbf{A} \cdot [(0)\hat{\mathbf{x}} - (0)\hat{\mathbf{y}} + (-2 - 3)\hat{\mathbf{z}}] \\
&= 3z(-5) = -15z
\end{aligned}$$

therefore,

$$\mathbf{B} \cdot (\nabla \times \mathbf{A}) - \mathbf{A} \cdot (\nabla \times \mathbf{B}) = 0 - (-15z) = 15z$$

(b) Checking (ii): LHS

$$\begin{aligned}
\nabla(\mathbf{A} \cdot \mathbf{B}) &= \nabla(x(3y) + 2y(-2x) + 3z(0)) \\
&= \nabla(3xy - 4xy) = \nabla(-xy) \\
&= -y\hat{\mathbf{x}} - x\hat{\mathbf{y}}
\end{aligned}$$

RHS:

$$\begin{aligned}
\mathbf{A} \times (\nabla \times \mathbf{B}) &= \mathbf{A} \times \begin{vmatrix} \hat{\mathbf{x}} & \hat{\mathbf{y}} & \hat{\mathbf{z}} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ 3y & -2x & 0 \end{vmatrix} \\
&= \mathbf{A} \times [(0)\hat{\mathbf{x}} - (0)\hat{\mathbf{y}} + (-5)\hat{\mathbf{z}}] \\
&= \begin{vmatrix} \hat{\mathbf{x}} & \hat{\mathbf{y}} & \hat{\mathbf{z}} \\ x & 2y & 3z \\ 0 & 0 & -5 \end{vmatrix} = -10y\hat{\mathbf{x}} + 5x\hat{\mathbf{y}}
\end{aligned}$$

and

$$\mathbf{B} \times (\nabla \times \mathbf{A}) = \mathbf{B} \times \begin{vmatrix} \hat{\mathbf{x}} & \hat{\mathbf{y}} & \hat{\mathbf{z}} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ x & 2y & 3z \end{vmatrix} = \mathbf{B} \times [(0)\hat{\mathbf{x}} - (0)\hat{\mathbf{y}} + (0)\hat{\mathbf{z}}] = 0$$

and

$$\begin{aligned}
(\mathbf{A} \cdot \nabla)\mathbf{B} &= \left(x\frac{\partial}{\partial x} + 2y\frac{\partial}{\partial y} + 3z\frac{\partial}{\partial z}\right)(3y\hat{\mathbf{x}} - 2x\hat{\mathbf{y}}) \\
&= 6y\hat{\mathbf{x}} - 2x\hat{\mathbf{y}}
\end{aligned}$$

and

$$\begin{aligned}(\mathbf{B} \cdot \nabla) \mathbf{A} &= \left(3y \frac{\partial}{\partial x} - 2x \frac{\partial}{\partial y} \right) (x\hat{\mathbf{x}} + 2y\hat{\mathbf{y}} + 3z\hat{\mathbf{z}}) \\ &= 3y\hat{\mathbf{x}} - 4x\hat{\mathbf{y}}\end{aligned}$$

therefore,

$$\begin{aligned}\mathbf{A} \times (\nabla \times \mathbf{B}) + \mathbf{B} \times (\nabla \times \mathbf{A}) + (\mathbf{A} \cdot \nabla) \mathbf{B} + (\mathbf{B} \cdot \nabla) \mathbf{A} &= (-10y\hat{\mathbf{x}} + 5x\hat{\mathbf{y}}) + (6y\hat{\mathbf{x}} - 2x\hat{\mathbf{y}}) + (3y\hat{\mathbf{x}} - 4x\hat{\mathbf{y}}) \\ &= -y\hat{\mathbf{x}} - x\hat{\mathbf{y}}\end{aligned}$$

(c) For rule (vi), the left hand side is

$$\begin{aligned}\nabla \times (\mathbf{A} \times \mathbf{B}) &= \nabla \times \begin{vmatrix} \hat{\mathbf{x}} & \hat{\mathbf{y}} & \hat{\mathbf{z}} \\ x & 2y & 3z \\ 3y & -2x & 0 \end{vmatrix} \\ &= \nabla \times [6xz\hat{\mathbf{x}} + 9yz\hat{\mathbf{y}} + (-2x^2 - 6y^2)\hat{\mathbf{z}}] \\ &= \begin{vmatrix} \hat{\mathbf{x}} & \hat{\mathbf{y}} & \hat{\mathbf{z}} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ 6xz & 9yz & -2x^2 - 6y^2 \end{vmatrix} \\ &= \hat{\mathbf{x}}(-12y - 9y) - \hat{\mathbf{y}}(-4x - 6x) + \hat{\mathbf{z}}(0) \\ &= -21y\hat{\mathbf{x}} + 10x\hat{\mathbf{y}}\end{aligned}$$

and on the right hand side, the new terms are

$$\mathbf{A}(\nabla \cdot \mathbf{B}) = \mathbf{A}[0 + 0] = 0$$

and

$$\begin{aligned}\mathbf{B}(\nabla \cdot \mathbf{A}) &= \mathbf{B}[1 + 2 + 3] = 6\mathbf{B} \\ &= 6(3y\hat{\mathbf{x}} - 2x\hat{\mathbf{y}}) = 18y\hat{\mathbf{x}} - 12x\hat{\mathbf{y}}\end{aligned}$$

combining these with the terms from (iv) gives

$$\begin{aligned}(\mathbf{B} \cdot \nabla) \mathbf{A} - (\mathbf{A} \cdot \nabla) \mathbf{B} + (\mathbf{A} \cdot \nabla) \mathbf{B} - (\mathbf{B} \cdot \nabla) \mathbf{A} &= (3y\hat{\mathbf{x}} - 4x\hat{\mathbf{y}}) - (6y\hat{\mathbf{x}} - 2x\hat{\mathbf{y}}) + 0 - (18y\hat{\mathbf{x}} - 12x\hat{\mathbf{y}}) \\ &= -21y\hat{\mathbf{x}} + 10x\hat{\mathbf{y}}\end{aligned}$$

1.26 Given the Laplacian of a scalar function T is

$$\nabla^2 T = \nabla \cdot (\nabla T) = \frac{\partial^2 T}{\partial x^2} + \frac{\partial^2 T}{\partial y^2} + \frac{\partial^2 T}{\partial z^2}$$

(a) $T_a = x^2 + 2xy + 3z + 4$:

$$\nabla^2 T_a = 2 + 0 + 0 = 2$$

(b) $T_b = \sin x \sin y \sin z$:

$$\frac{\partial^2 T_b}{\partial x^2} = \frac{\partial^2 T_b}{\partial y^2} = \frac{\partial^2 T_b}{\partial z^2} = -\sin x \sin y \sin z = -T_b$$

$$\nabla^2 T_b = -3T_b$$

(c) $T_c = e^{-5x} \sin 4y \cos 3z$: The components are

$$\frac{\partial^2 T_c}{\partial x^2} = 25e^{-5x} \sin 4y \cos 3z = 25T_c$$

$$\frac{\partial^2 T_c}{\partial y^2} = -16e^{-5x} \sin 4y \cos 3z = -16T_c$$

$$\frac{\partial^2 T_c}{\partial z^2} = -9e^{-5x} \sin 4y \cos 3z = -9T_c$$

therefore

$$\nabla^2 T_c = T_c(25 - 16 - 9) = 0$$

(d) $\mathbf{v} = x^2\hat{\mathbf{x}} + 3xz^2\hat{\mathbf{y}} - 2xz\hat{\mathbf{z}}$: The laplacian of a vector function is

$$\nabla^2 \mathbf{v} = (\nabla^2 v_x)\hat{\mathbf{x}} + (\nabla^2 v_y)\hat{\mathbf{y}} + (\nabla^2 v_z)\hat{\mathbf{z}}$$

and the components are

$$\nabla^2 v_x = \frac{\partial^2 v_x}{\partial x^2} + \frac{\partial^2 v_x}{\partial y^2} + \frac{\partial^2 v_x}{\partial z^2} = 2 + 0 + 0 = 2$$

$$\nabla^2 v_y = 0 + 0 + 6x = 6x$$

$$\nabla^2 v_z = 0$$

therefore

$$\nabla^2 \mathbf{v} = 2\hat{\mathbf{x}} + 6x\hat{\mathbf{y}}$$

1.27 The divergence of curl is always zero:

$$\begin{aligned} \nabla \cdot (\nabla \times \mathbf{v}) &= \nabla \cdot \begin{vmatrix} \hat{\mathbf{x}} & \hat{\mathbf{y}} & \hat{\mathbf{z}} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ v_x & v_y & v_z \end{vmatrix} \\ &= \nabla \cdot \left(\hat{\mathbf{x}} \left(\frac{\partial v_z}{\partial y} - \frac{\partial v_y}{\partial z} \right) - \hat{\mathbf{y}} \left(\frac{\partial v_z}{\partial x} - \frac{\partial v_x}{\partial z} \right) + \hat{\mathbf{z}} \left(\frac{\partial v_y}{\partial x} - \frac{\partial v_x}{\partial y} \right) \right) \\ &= \frac{\partial}{\partial x} \left(\frac{\partial v_z}{\partial y} - \frac{\partial v_y}{\partial z} \right) + \frac{\partial}{\partial y} \left(\frac{\partial v_x}{\partial z} - \frac{\partial v_z}{\partial x} \right) + \frac{\partial}{\partial z} \left(\frac{\partial v_y}{\partial x} - \frac{\partial v_x}{\partial y} \right) \\ &= \left[\frac{\partial}{\partial x} \left(\frac{\partial v_z}{\partial y} \right) - \frac{\partial}{\partial y} \left(\frac{\partial v_z}{\partial x} \right) \right] + \left[\frac{\partial}{\partial y} \left(\frac{\partial v_x}{\partial z} \right) - \frac{\partial}{\partial z} \left(\frac{\partial v_x}{\partial y} \right) \right] + \left[\frac{\partial}{\partial z} \left(\frac{\partial v_y}{\partial x} \right) - \frac{\partial}{\partial x} \left(\frac{\partial v_y}{\partial z} \right) \right] \\ \nabla \cdot (\nabla \times \mathbf{v}) &= 0 \end{aligned}$$

where the last step hinges on the equality of cross derivatives:

$$\frac{\partial}{\partial x_i} \left(\frac{\partial v}{\partial x_j} \right) = \frac{\partial}{\partial x_j} \left(\frac{\partial v}{\partial x_i} \right)$$

Checking for $v_a = x^2\hat{\mathbf{x}} + 2xz^2\hat{\mathbf{y}} - 2xz\hat{\mathbf{z}}$:

$$\begin{aligned} \nabla \cdot (\nabla \times v_a) &= \nabla \cdot \begin{vmatrix} \hat{\mathbf{x}} & \hat{\mathbf{y}} & \hat{\mathbf{z}} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ x^2 & 2xz^2 & -2xz \end{vmatrix} \\ &= \nabla \cdot [\hat{\mathbf{x}}(0 - 4xz) - \hat{\mathbf{y}}(-2z - 0) + \hat{\mathbf{z}}(2z^2 - 0)] \\ &= \nabla \cdot \left[\frac{\partial}{\partial x}(-4xz) + \frac{\partial}{\partial y}(2z) + \frac{\partial}{\partial z}(2z^2) \right] \\ &= -4z + 0 + 4z = 0 \end{aligned}$$

1.28 The curl of gradient is always zero:

$$\begin{aligned} \nabla \times (\nabla T) &= \nabla \times \left(\frac{\partial T}{\partial x}\hat{\mathbf{x}} + \frac{\partial T}{\partial y}\hat{\mathbf{y}} + \frac{\partial T}{\partial z}\hat{\mathbf{z}} \right) \\ &= \begin{vmatrix} \hat{\mathbf{x}} & \hat{\mathbf{y}} & \hat{\mathbf{z}} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ \frac{\partial T}{\partial x} & \frac{\partial T}{\partial y} & \frac{\partial T}{\partial z} \end{vmatrix} \\ &= \hat{\mathbf{x}} \left[\frac{\partial}{\partial y} \left(\frac{\partial T}{\partial z} \right) - \frac{\partial}{\partial z} \left(\frac{\partial T}{\partial y} \right) \right] - \hat{\mathbf{y}} \left[\frac{\partial}{\partial x} \left(\frac{\partial T}{\partial z} \right) - \frac{\partial}{\partial z} \left(\frac{\partial T}{\partial x} \right) \right] + \hat{\mathbf{z}} \left[\frac{\partial}{\partial x} \left(\frac{\partial T}{\partial y} \right) - \frac{\partial}{\partial y} \left(\frac{\partial T}{\partial x} \right) \right] \\ \nabla \times (\nabla T) &= 0 \end{aligned}$$

where the last step uses the equality of cross derivatives again. Checking for $T = x^2y^3z^4$:

$$\frac{\partial T}{\partial x} = 2xy^3z^4, \quad \frac{\partial T}{\partial y} = 3x^2y^2z^4, \quad \text{and} \quad \frac{\partial T}{\partial z} = 4x^2y^3z^3$$

and

$$\begin{aligned} \nabla \times (\nabla T) &= \nabla \times (2xy^3z^4\hat{\mathbf{x}} + 3x^2y^2z^4\hat{\mathbf{y}} + 4x^2y^3z^3\hat{\mathbf{z}}) \\ &= \begin{vmatrix} \hat{\mathbf{x}} & \hat{\mathbf{y}} & \hat{\mathbf{z}} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ 2xy^3z^4 & 3x^2y^2z^4 & 4x^2y^3z^3 \end{vmatrix} \\ &= \hat{\mathbf{x}}(12x^2y^2z^4 - 12x^2y^2z^4) - \hat{\mathbf{y}}(8x^2y^3z^3 - 8x^2y^3z^3) + \hat{\mathbf{z}}(6x^2y^3z^3 - 6x^2y^3z^3) \\ &= 0 \end{aligned}$$

1.29 Calculating the line integral of the function $\mathbf{v} = x^2\hat{\mathbf{x}} + 2yz\hat{\mathbf{y}} + y^2\hat{\mathbf{z}}$: from the origin to point $(1, 1, 1)$ along three different paths:

(a) $a = (0, 0, 0) \rightarrow b = (1, 0, 0) \rightarrow c = (1, 1, 0) \rightarrow d = (1, 1, 1)$ split to three paths:

(i) From $a \rightarrow b$: $dl = dx\hat{\mathbf{x}}$ and $\mathbf{v} = x^2\hat{\mathbf{x}}$.

(ii) From $b \rightarrow c$: $dl = dy\hat{\mathbf{y}}$ and $\mathbf{v} = 2yz\hat{\mathbf{y}} = 0$ since $z = 0$.

(iii) From $c \rightarrow d$: $dl = dz\hat{\mathbf{z}}$ and $\mathbf{v} = y^2\hat{\mathbf{z}} = 1\hat{\mathbf{z}}$ since $y = 1$.

$$\begin{aligned} \int_a^b \mathbf{v} \cdot d\mathbf{l} &= \int_0^1 x^2 dx = \frac{1}{3} \\ \int_b^c \mathbf{v} \cdot d\mathbf{l} &= \int_0^1 0 dy = 0 \\ \int_c^d \mathbf{v} \cdot d\mathbf{l} &= \int_0^1 1 dz = 1 \\ \int_a^d \mathbf{v} \cdot d\mathbf{l} &= \frac{1}{3} + 0 + 1 = \frac{4}{3} \end{aligned}$$

(b) $a = (0, 0, 0) \rightarrow b = (0, 0, 1) \rightarrow c = (0, 1, 1) \rightarrow d = (1, 1, 1)$ split to three paths:

(i) From $a \rightarrow b$: $dl = dz\hat{\mathbf{z}}$ and $\mathbf{v} = y^2\hat{\mathbf{z}} = 0$ since $y = 0$.

(ii) From $b \rightarrow c$: $dl = dy\hat{\mathbf{y}}$ and $\mathbf{v} = 2yz\hat{\mathbf{y}} = 2y\hat{\mathbf{y}}$ since $y = 1$.

(iii) From $c \rightarrow d$: $dl = dx\hat{\mathbf{x}}$ and $\mathbf{v} = x^2\hat{\mathbf{x}}$.

$$\begin{aligned} \int_a^b \mathbf{v} \cdot d\mathbf{l} &= \int_0^1 0 dz = 0 \\ \int_b^c \mathbf{v} \cdot d\mathbf{l} &= \int_0^1 2y dy = 1 \\ \int_c^d \mathbf{v} \cdot d\mathbf{l} &= \int_0^1 x^2 dx = \frac{1}{3} \\ \int_a^d \mathbf{v} \cdot d\mathbf{l} &= 0 + 1 + \frac{1}{3} = \frac{4}{3} \end{aligned}$$

(c) A straight line: Since $x = y = z$ and $dx = dy = dz$,

$dl = dx(\hat{\mathbf{x}} + \hat{\mathbf{y}} + \hat{\mathbf{z}})$ and $\mathbf{v} = x^2\hat{\mathbf{x}} + 2yz\hat{\mathbf{y}} + y^2\hat{\mathbf{z}} = 4x^2\hat{\mathbf{x}}$.

$$\int_a^d \mathbf{v} \cdot d\mathbf{l} = \int_0^1 4x^2 dx = \frac{4}{3}$$

(d) The line integral around the closed loop:

$$\oint \mathbf{v} \cdot d\mathbf{l} = \int_{(a)} \mathbf{v} \cdot d\mathbf{l} - \int_{(b)} \mathbf{v} \cdot d\mathbf{l} = \frac{4}{3} - \frac{4}{3} = 0$$

1.30 Surface integral of $\mathbf{v} = 2xz\hat{\mathbf{x}} + (x+2)\hat{\mathbf{y}} + y(z^2-3)\hat{\mathbf{z}}$ over the bottom of the box:
 $z = 0$, $d\mathbf{A} = dx dy \hat{\mathbf{z}}$ $\mathbf{v} \cdot d\mathbf{A} = y(z^2-3) dx dy = -3y dx dy$, so

$$\int \mathbf{v} \cdot d\mathbf{A} = \int_0^2 dx \int_0^2 -3y dy = -12$$

From Example 1.7 (v) has the same boundary line but a different surface integral, so the surface integral does not depend on just the boundary line. The surface over the closed surface of the box is

$$\oint \mathbf{v} \cdot d\mathbf{A} = 20 + 12 = 32$$

since the direction of $d\mathbf{A}$ on the bottom side is in the negative z direction for it to point ‘outward’.

1.31 Calculating the volume integral of $T = z^2$ over the tetrahedron with vertices $(0,0,0)$, $(1,0,0)$, $(0,1,0)$, and $(0,0,1)$:

The equation of the plane containing the three vertices $A = (1,0,0)$, $B = (0,1,0)$, and $C(0,0,1)$:
The vector normal to this plane $\mathbf{n} = (a,b,c)$ is the cross product of two vectors in the plane given by $\mathbf{AB} = (-1,1,0)$ and $\mathbf{AC} = (-1,0,1)$:

$$\mathbf{n} = \begin{vmatrix} x & y & z \\ -1 & 1 & 0 \\ -1 & 0 & 1 \end{vmatrix} = (1,1,1)$$

the equation of the plane is therefore

$$\begin{aligned} \mathbf{n} \cdot (\mathbf{r} - \mathbf{r}_o) &= 0 \\ (1,1,1) \cdot [(x,y,z) - A] &= 0 \\ x + y + z &= 1 \end{aligned}$$

therefore, the boundary for x is $x = 0$ and $x = 1 - y - z$; for y is $y = 0$ and $y = 1 - z$; and for z is $z = 0$ and $z = 1$. The volume integral is therefore

$$\begin{aligned} \int T dV &= \int_0^1 z^2 dz \int_0^{1-z} dy \int_0^{1-y-z} dx \\ &= \int_0^1 z^2 dz \int_0^{1-z} (1-y-z) dy \\ &= \int_0^1 z^2 dz \left(y - y^2/2 - yz \Big|_0^{1-z} \right) \\ &= \int_0^1 z^2 [(1-z) - (1-z)^2/2 - z(1-z)] dz \\ &= \int_0^1 z^2 (1-z-1/2-z^2/2+z-z+z^2) dz \\ &= \int_0^1 z^2 (1/2-z+z^2/2) dz \\ &= \int_0^1 (z^2/2 - z^3 + z^4/2) dz \\ &= z^3/6 - z^4/4 + z^5/10 \Big|_0^1 \\ &= 1/6 - 1/4 + 1/10 = 1/60 \end{aligned}$$

1.32 Given $T = x^2 + 4xy + 2yz^3$,

$$\frac{\partial T}{\partial x} = 2x + 4y, \quad \frac{\partial T}{\partial y} = 4x + 2z^3, \quad \text{and} \quad \frac{\partial T}{\partial z} = 6yz^2$$

(iii) $(0, 1, 1) \rightarrow b$:

$$z : 0 \rightarrow 1; \quad y = z = 1, \quad dy = dz = 0; \quad dl = dx \hat{\mathbf{x}}; \quad \nabla T \cdot d\mathbf{l} = (2x + 4) dx$$

and

$$\int_d^b \nabla T \cdot d\mathbf{l} = \int_0^1 (2x + 4) dx = 5$$

therefore

$$\int_a^b \nabla T \cdot d\mathbf{l} = 0 + 2 + 5 = 7$$

(c) the parabolic path $z = x^2$; $y = x$:

$$dx = dy, \quad \text{and} \quad dz = 2x dx; \quad d\mathbf{l} = dx \hat{\mathbf{x}} + dx \hat{\mathbf{y}} + 2x dx \hat{\mathbf{z}}$$

and

$$\begin{aligned} \nabla T \cdot d\mathbf{l} &= (2x + 4y) dx + (4x + 2z^3) dx + (6yz^2) 2x dx \\ &= 6x dx + (4x + 2x^6) dx + (12x^6) dx \\ &= 10x dx + 14x^6 dx \end{aligned}$$

therefore

$$\begin{aligned} \int_a^b \nabla T \cdot d\mathbf{l} &= \int_0^1 (10x + 14x^6) dx \\ &= 5x^2 + 2x^7 \Big|_0^1 = 7 \end{aligned}$$

1.33 Checking the divergence theorem: For the function

$$\mathbf{v} = (xy)\hat{\mathbf{x}} + (2yz)\hat{\mathbf{y}} + (3zx)\hat{\mathbf{z}}$$

the divergence is

$$\nabla \cdot \mathbf{v} = y + 2z + 3x$$

so the volume integral is

$$\begin{aligned} \int_V \nabla \cdot \mathbf{v} d\tau &= \int_0^2 \int_0^2 \int_0^2 (y + 2z + 3x) dx dy dz \\ &= \int_0^2 \int_0^2 (2y + 4z + 6) dy dz \\ &= \int_0^2 (4 + 8z + 12) dz \\ &= 8 + 16 + 24 \\ \int_V \nabla \cdot \mathbf{v} d\tau &= 48 \end{aligned}$$

The surface integral is evaluated over the six faces of the cube noted by Figure 1.29:

(i) $x = 2$, $d\mathbf{A} = dy dz \hat{\mathbf{x}}$, $\mathbf{v} \cdot d\mathbf{A} = 2y dy dz$;

$$\int \mathbf{v} \cdot d\mathbf{A} = \int_0^2 \int_0^2 2y dy dz = 8$$

1.34 Testing Stokes' theorem for the function

$$\mathbf{v} = (xy)\hat{\mathbf{x}} + (2yz)\hat{\mathbf{y}} + (3zx)\hat{\mathbf{z}}$$

using the triangular shaded area bounded by the vertices $O = (0, 0, 0)$, $A = (0, 2, 0)$, and $B = (0, 0, 2)$:

$$\begin{aligned}\nabla \times \mathbf{v} &= (0 - 2y)\hat{\mathbf{x}} - (3z - 0)\hat{\mathbf{y}} + (0 - x)\hat{\mathbf{z}} \quad \text{and} \quad d\mathbf{A} = dz dy \hat{\mathbf{x}} \\ &= -2y\hat{\mathbf{x}} - 3z\hat{\mathbf{y}} - x\hat{\mathbf{z}}\end{aligned}$$

$x = 0$ on this surface, and the limits of integration are $y : 0 \rightarrow 2$ and $z = 0 \rightarrow z = 2 - y$:

$$(\nabla \times \mathbf{v}) \cdot d\mathbf{A} = -2y dz dy$$

Thus, the flux of the curl through the surface is

$$\begin{aligned}\int_S (\nabla \times \mathbf{v}) \cdot d\mathbf{A} &= \int_0^2 \int_0^{2-y} -2y dz dy \\ &= \int_0^2 -2y(2 - y) dy \\ &= -2y^2 + \frac{2}{3}y^3 \Big|_0^2 = -8/3\end{aligned}$$

The line integral is evaluated over the three sides of the triangle:

(i) On the path OA :

$$x = z = 0; d\mathbf{l} = dy \hat{\mathbf{y}}; \mathbf{v} \cdot d\mathbf{l} = 2yz dy = 0;$$

$$\int_{OA} \mathbf{v} \cdot d\mathbf{l} = 0$$

(ii) On the path AB :

$$x = 0, y = 2 - z; dy = -dz; d\mathbf{l} = -dz (\hat{\mathbf{y}} - \hat{\mathbf{z}}); \mathbf{v} \cdot d\mathbf{l} = -2yz dz = -2(2 - z)z dz = (2z^2 - 4z) dz;$$

$$\int_{AB} \mathbf{v} \cdot d\mathbf{l} = \int_0^2 (2z^2 - 4z) dz = -8/3$$

(iii) On the path BO :

$$x = y = 0; d\mathbf{l} = dz \hat{\mathbf{z}}; \mathbf{v} \cdot d\mathbf{l} = 0;$$

$$\int_{BO} \mathbf{v} \cdot d\mathbf{l} = 0$$

So, the circulation of \mathbf{v} around the triangle is

$$\oint \mathbf{v} \cdot d\mathbf{l} = 0 + -8/3 + 0 = -8/3$$

thus,

$$\int_S (\nabla \times \mathbf{v}) \cdot d\mathbf{A} = \oint \mathbf{v} \cdot d\mathbf{l}$$

1.35 By Corollary 1, the function

$$\mathbf{v} = (2xz + 3y^2)\hat{\mathbf{y}} + (4yz^2)\hat{\mathbf{z}}$$

with curl

$$\nabla \times \mathbf{v} = (4z^2 - 2x)\hat{\mathbf{x}} + 2z\hat{\mathbf{z}}$$

or rewritten as

$$\mathbf{B} \cdot (\nabla \times \mathbf{A}) = \nabla \cdot (\mathbf{A} \times \mathbf{B}) + \mathbf{A} \cdot (\nabla \times \mathbf{B})$$

integrating both sides over a volume:

$$\int_V \mathbf{B} \cdot (\nabla \times \mathbf{A}) dV = \int_V \nabla \cdot (\mathbf{A} \times \mathbf{B}) dV + \int_V \mathbf{A} \cdot (\nabla \times \mathbf{B}) dV$$

Using the divergence theorem $\int_V (\nabla \cdot \mathbf{v}) dV = \oint_S \mathbf{v} \cdot d\mathbf{a}$:

$$\int_V \mathbf{B} \cdot (\nabla \times \mathbf{A}) dV = \oint_S (\mathbf{A} \times \mathbf{B}) \cdot d\mathbf{a} + \int_V \mathbf{A} \cdot (\nabla \times \mathbf{B}) dV$$

1.37 Given the relation of Cartesian to spherical coordinates:

$$x = r \sin \theta \cos \phi, \quad y = r \sin \theta \sin \phi, \quad \text{and} \quad z = r \cos \theta$$

To find the formula for r , take the sum of the squares of the three equations; Solve for θ using the third equation; and solve for ϕ by dividing the second equation by the first:

$$r = \sqrt{x^2 + y^2 + z^2}, \quad \theta = \arccos \frac{z}{r}, \quad \text{and} \quad \phi = \arctan \frac{y}{x}$$

1.38 From the position vector

$$\begin{aligned} \mathbf{r} &= x\hat{\mathbf{x}} + y\hat{\mathbf{y}} + z\hat{\mathbf{z}} \\ \mathbf{r} &= r \sin \theta \cos \phi \hat{\mathbf{x}} + r \sin \theta \sin \phi \hat{\mathbf{y}} + r \cos \theta \hat{\mathbf{z}} \\ \mathbf{r} &= r \hat{\mathbf{r}}(\theta, \phi) \end{aligned}$$

where the unit vector $\hat{\mathbf{r}}(\theta, \phi)$ is dependent on θ and ϕ . The new basis vectors are in the same direction as the partial derivatives with respect to r , θ , and ϕ , so

$$\hat{\mathbf{r}} = \frac{e_r}{|e_r|}, \quad \hat{\boldsymbol{\theta}} = \frac{e_\theta}{|e_\theta|}, \quad \text{and} \quad \hat{\boldsymbol{\phi}} = \frac{e_\phi}{|e_\phi|}$$

The partial derivatives are

$$\begin{aligned} e_r &= \frac{\partial \mathbf{r}}{\partial r} = \sin \theta \cos \phi \hat{\mathbf{x}} + \sin \theta \sin \phi \hat{\mathbf{y}} + \cos \theta \hat{\mathbf{z}} \\ e_\theta &= \frac{\partial \mathbf{r}}{\partial \theta} = r \cos \theta \cos \phi \hat{\mathbf{x}} + r \cos \theta \sin \phi \hat{\mathbf{y}} - r \sin \theta \hat{\mathbf{z}} \\ e_\phi &= \frac{\partial \mathbf{r}}{\partial \phi} = -r \sin \theta \sin \phi \hat{\mathbf{x}} + r \sin \theta \cos \phi \hat{\mathbf{y}} \end{aligned}$$

and the magnitude

$$\begin{aligned} |e_r| &= \sqrt{\sin^2 \theta \cos^2 \phi + \sin^2 \theta \sin^2 \phi + \cos^2 \theta} \\ &= \sqrt{\sin^2 \theta (\cos^2 \phi + \sin^2 \phi) + \cos^2 \theta} \\ &= \sqrt{\sin^2 \theta + \cos^2 \theta} = 1 \\ |e_\theta| &= \sqrt{r^2 (\cos^2 \theta \cos^2 \phi + \cos^2 \theta \sin^2 \phi + \sin^2 \theta)} \\ &= \sqrt{r^2 (\cos^2 \theta (\cos^2 \phi + \sin^2 \phi) + \sin^2 \theta)} \\ &= \sqrt{r^2 (\cos^2 \theta + \sin^2 \theta)} = r \\ |e_\phi| &= \sqrt{r^2 (\sin^2 \theta \sin^2 \phi + \sin^2 \theta \cos^2 \phi)} \\ &= \sqrt{r^2 \sin^2 \theta (\sin^2 \phi + \cos^2 \phi)} \\ &= \sqrt{r^2 \sin^2 \theta} = r \sin \theta \end{aligned}$$

thus, the unit vectors are:

$$\begin{aligned}\hat{\mathbf{r}} &= \sin \theta \cos \phi \hat{\mathbf{x}} + \sin \theta \sin \phi \hat{\mathbf{y}} + \cos \theta \hat{\mathbf{z}} \\ \hat{\boldsymbol{\theta}} &= \cos \theta \cos \phi \hat{\mathbf{x}} + \cos \theta \sin \phi \hat{\mathbf{y}} - \sin \theta \hat{\mathbf{z}} \\ \hat{\boldsymbol{\phi}} &= -\sin \phi \hat{\mathbf{x}} + \cos \phi \hat{\mathbf{y}}\end{aligned}$$

or in matrix form:

$$\begin{pmatrix} \hat{\mathbf{r}} \\ \hat{\boldsymbol{\theta}} \\ \hat{\boldsymbol{\phi}} \end{pmatrix} = \begin{pmatrix} \sin \theta \cos \phi & \sin \theta \sin \phi & \cos \theta \\ \cos \theta \cos \phi & \cos \theta \sin \phi & -\sin \theta \\ -\sin \phi & \cos \phi & 0 \end{pmatrix} \begin{pmatrix} \hat{\mathbf{x}} \\ \hat{\mathbf{y}} \\ \hat{\mathbf{z}} \end{pmatrix}$$

where $a = Qx$ is an orthogonal matrix, so $Q^T = Q^{-1}$ and $Q^T Q = I$. Multiplying both sides by Q^T :

$$Q^T a = Q^T Q x \rightarrow x = Q^T a$$

thus, the inverse formula is

$$\begin{pmatrix} \hat{\mathbf{x}} \\ \hat{\mathbf{y}} \\ \hat{\mathbf{z}} \end{pmatrix} = \begin{pmatrix} \sin \theta \cos \phi & \cos \theta \cos \phi & -\sin \phi \\ \sin \theta \sin \phi & \cos \theta \sin \phi & \cos \phi \\ \cos \theta & -\sin \theta & 0 \end{pmatrix} \begin{pmatrix} \hat{\mathbf{r}} \\ \hat{\boldsymbol{\theta}} \\ \hat{\boldsymbol{\phi}} \end{pmatrix}$$

or in vector form:

$$\begin{aligned}\hat{\mathbf{x}} &= \sin \theta \cos \phi \hat{\mathbf{r}} + \cos \theta \cos \phi \hat{\boldsymbol{\theta}} - \sin \phi \hat{\boldsymbol{\phi}} \\ \hat{\mathbf{y}} &= \sin \theta \sin \phi \hat{\mathbf{r}} + \cos \theta \sin \phi \hat{\boldsymbol{\theta}} + \cos \phi \hat{\boldsymbol{\phi}} \\ \hat{\mathbf{z}} &= \cos \theta \hat{\mathbf{r}} - \sin \theta \hat{\boldsymbol{\theta}}\end{aligned}$$

1.39 (a) Divergence theorem for $\mathbf{v}_1 = r^2 \hat{\mathbf{r}}$ using a volume of a sphere of radius R centered at the origin: The divergence is

$$\nabla \cdot \mathbf{v}_1 = \frac{1}{r^2} \frac{\partial}{\partial r} (r^2(r^2)) = 4r$$

and the volume and surface elements are

$$dV = r^2 \sin \theta \, dr \, d\theta \, d\phi, \quad \text{and} \quad d\mathbf{a}_1 = (r^2 \sin \theta) \, d\theta \, d\phi \, \hat{\mathbf{r}}$$

So

$$\begin{aligned}\int_V (\nabla \cdot \mathbf{v}_1) \cdot dV &= \int_{r=0}^R \int_{\theta=0}^{\pi} \int_{\phi=0}^{2\pi} 4r(r^2 \sin \theta) \, d\theta \, d\phi \, dr \\ &= \int_0^R 4r^3 \, dr \int_0^{\pi} \sin \theta \, d\theta \int_0^{2\pi} d\phi \\ &= (R^4)(2)(2\pi) = 4\pi R^4\end{aligned}$$

and

$$\begin{aligned}\oint_S \mathbf{v}_1 \cdot d\mathbf{a}_1 &= \int_{\theta=0}^{\pi} \int_{\phi=0}^{2\pi} r^2(r^2 \sin \theta) \, d\theta \, d\phi \\ &= r^4 \int_0^{\pi} \sin \theta \, d\theta \int_0^{2\pi} d\phi \\ &= (r^4)(2)(2\pi) = 4\pi r^4\end{aligned}$$

where $r = R$ on the surface of the sphere. Therefore,

$$\int_V (\nabla \cdot \mathbf{v}_1) \cdot dV = \oint_S \mathbf{v}_1 \cdot d\mathbf{a}_1$$

(b) For $\mathbf{v}_2 = (1/r^2)\hat{\mathbf{r}}$:

$$\nabla \cdot \mathbf{v}_2 = \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 (1/r^2)) = 0$$

So

$$\int_V (\nabla \cdot \mathbf{v}_2) \cdot dV = 0$$

and

$$\begin{aligned} \oint_S \mathbf{v}_2 \cdot d\mathbf{a}_1 &= \int_{\theta=0}^{\pi} \int_{\phi=0}^{2\pi} \frac{1}{r^2} (r^2 \sin \theta) d\theta d\phi \\ &= \int_0^{\pi} \sin \theta d\theta \int_0^{2\pi} d\phi = 4\pi \end{aligned}$$

1.40 Given the function

$$\mathbf{v} = (r \cos \theta)\hat{\mathbf{r}} + (r \sin \theta)\hat{\boldsymbol{\theta}} + (r \sin \theta \cos \phi)\hat{\boldsymbol{\phi}}$$

the divergence in spherical coordinates is

$$\begin{aligned} \nabla \cdot \mathbf{v} &= \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 v_r) + \frac{1}{r \sin \theta} \frac{\partial}{\partial \theta} (\sin \theta v_\theta) + \frac{1}{r \sin \theta} \frac{\partial}{\partial \phi} (v_\phi) \\ &= \frac{1}{r^2} \frac{\partial}{\partial r} (r^3 \cos \theta) + \frac{1}{r \sin \theta} \frac{\partial}{\partial \theta} (r \sin^2 \theta) + \frac{1}{r \sin \theta} \frac{\partial}{\partial \phi} (r \sin \theta \cos \phi) \\ &= 3 \cos \theta + 2 \cos \theta - \sin \phi \\ &= 5 \cos \theta - \sin \phi \end{aligned}$$

Checking the divergence theorem using a volume of a inverted hemispherical bowl of radius R , resting on the xy plane and centered at the origin:

The volume and surface elements are

$$dV = r^2 \sin \theta dr d\theta d\phi, \quad \text{and} \quad d\mathbf{a}_1 = r^2 \sin \theta d\theta d\phi \hat{\mathbf{r}}$$

The volume integral for the hemisphere is

$$\begin{aligned} \int_V (\nabla \cdot \mathbf{v}) \cdot dV &= \int_{r=0}^R \int_{\theta=0}^{\pi/2} \int_{\phi=0}^{2\pi} (5 \cos \theta - \sin \phi) (r^2 \sin \theta) d\theta d\phi dr \\ &= \int_0^R r^2 dr \int_0^{\pi/2} \sin \theta \int_0^{2\pi} (5 \cos \theta - \sin \phi) d\phi d\theta \\ &= \frac{R^3}{3} \int_0^{\pi/2} 5\phi \cos \theta + \cos \phi \Big|_0^{2\pi} d\theta \\ &= \frac{R^3}{3} (10\pi) \int_0^{\pi/2} \sin \theta \cos \theta d\theta \end{aligned}$$

$$\begin{aligned} \text{using } u = \sin \theta, \quad du = \cos \theta d\theta; \quad \int u du &= \frac{u^2}{2} \\ &= \frac{5\pi R^3}{3} \sin^2 \theta \Big|_0^{\pi/2} = \frac{5\pi R^3}{3} \end{aligned}$$

The surface integral is split into two parts: the top surface of the hemisphere and the circular base.

(i) The top surface of the hemisphere where $r = R$:

$$\begin{aligned} \oint_S \mathbf{v} \cdot d\mathbf{a}_1 &= \int_{\theta=0}^{\pi/2} \int_{\phi=0}^{2\pi} (r \cos \theta) (r^2 \sin \theta) d\theta d\phi \\ &= r^3 \int_0^{\pi/2} \sin \theta \cos \theta d\theta \int_0^{2\pi} d\phi \\ &= \pi r^3 = \pi R^3 \end{aligned}$$

(ii) The circular base of the hemisphere where $\theta = \pi/2$ and $\mathbf{a}_2 = r \, dr \, d\phi \, \hat{\boldsymbol{\theta}}$:

$$\begin{aligned}\oint_S \mathbf{v} \cdot d\mathbf{a}_2 &= \int_{r=0}^R \int_{\phi=0}^{2\pi} (r \sin \theta) r \, dr \, d\phi \\ &= \sin(\pi/2) \int_0^R r^2 \, dr \int_0^{2\pi} d\phi \\ &= (1) \frac{R^3}{3} (2\pi) = \frac{2\pi R^3}{3}\end{aligned}$$

So, the total surface integral is

$$\oint_S \mathbf{v} \cdot d\mathbf{a}_1 + \oint_S \mathbf{v} \cdot d\mathbf{a}_2 = \pi R^3 + \frac{2\pi R^3}{3} = \frac{5\pi R^3}{3}$$

1.41

$$T = r(\cos \theta + \sin \theta \cos \phi)$$

The partial derivatives are:

$$\begin{aligned}\frac{\partial T}{\partial r} &= \cos \theta + \sin \theta \cos \phi \\ \frac{\partial T}{\partial \theta} &= r(-\sin \theta + \cos \theta \cos \phi) \\ \frac{\partial T}{\partial \phi} &= -r \sin \theta \sin \phi\end{aligned}$$

thus, the gradient of T in spherical is

$$\begin{aligned}\nabla T &= \frac{\partial T}{\partial r} \hat{\mathbf{r}} + \frac{1}{r} \frac{\partial T}{\partial \theta} \hat{\boldsymbol{\theta}} + \frac{1}{r \sin \theta} \frac{\partial T}{\partial \phi} \hat{\boldsymbol{\phi}} \\ &= (\cos \theta + \sin \theta \cos \phi) \hat{\mathbf{r}} + (-\sin \theta + \cos \theta \cos \phi) \hat{\boldsymbol{\theta}} - (\sin \phi) \hat{\boldsymbol{\phi}}\end{aligned}$$

and partial derivative in the laplacian are:

$$\begin{aligned}\frac{\partial}{\partial r} \left(r^2 \frac{\partial T}{\partial r} \right) &= \frac{\partial}{\partial r} (r^2 (\cos \theta + \sin \theta \cos \phi)) = 2r(\cos \theta + \sin \theta \cos \phi) \\ \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial T}{\partial \theta} \right) &= r \frac{\partial}{\partial \theta} (\sin \theta (-\sin \theta + \cos \theta \cos \phi)) = r \frac{\partial}{\partial \theta} (-\sin^2 \theta + \sin \theta \cos \theta \cos \phi) \\ &= -2r \sin \theta \cos \theta + r \cos^2 \theta \cos \phi - r \sin^2 \theta \cos \phi \\ \frac{\partial^2 T}{\partial \phi^2} &= -r \sin \theta \cos \phi\end{aligned}$$

The laplacian of T in spherical is

$$\nabla^2 T = \frac{1}{r^2} \frac{\partial}{\partial r} \left(r^2 \frac{\partial T}{\partial r} \right) + \frac{1}{r^2 \sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial T}{\partial \theta} \right) + \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2 T}{\partial \phi^2}$$

Simplifying each term: (i) The first term:

$$\frac{2}{r} (\cos \theta + \sin \theta \cos \phi)$$

(ii) The second term:

$$-\frac{2}{r} \cos \theta + \frac{\cos^2 \theta \cos \phi}{r \sin \theta} - \frac{1}{r} \sin \theta \cos \phi$$

(iii) The third term:

$$-\frac{\cos \phi}{r \sin \theta}$$

adding all three terms:

$$\begin{aligned}\nabla^2 T &= \frac{2}{r}(\cos \theta + \sin \theta \cos \phi) - \frac{2}{r} \cos \theta + \frac{\cos^2 \theta \cos \phi}{r \sin \theta} - \frac{1}{r} \sin \theta \cos \phi - \frac{\cos \phi}{r \sin \theta} \\ &= \frac{2}{r}(\sin \theta \cos \phi) + \frac{\cos^2 \theta \cos \phi - \cos \phi}{r \sin \theta} - \frac{1}{r} \sin \theta \cos \phi \\ &= \frac{2}{r \sin \theta}(\sin^2 \theta \cos \phi) + \frac{\cos^2 \theta \cos \phi - \cos \phi}{r \sin \theta} - \frac{1}{r \sin \theta} \sin^2 \theta \cos \phi \\ &= \frac{1}{r \sin \theta}(\sin^2 \theta \cos \phi + \cos^2 \theta \cos \phi - \cos \phi) \\ &= \frac{1}{r \sin \theta}(\cos \phi)(\sin^2 \theta + \cos^2 \theta - 1) = 0\end{aligned}$$

Converting T to Cartesian coordinates:

$$x = r \sin \theta \cos \phi, \quad y = r \sin \theta \sin \phi, \quad \text{and} \quad z = r \cos \theta$$

So

$$T = z + x$$

The laplacian of T in Cartesian coordinates is

$$\nabla^2 T = \frac{\partial^2 T}{\partial x^2} + \frac{\partial^2 T}{\partial y^2} + \frac{\partial^2 T}{\partial z^2} = 0$$

Testing the gradient theorem using the path $O \rightarrow A = (2, 0, 0) \rightarrow B = (0, 2, 0) \rightarrow C = (0, 0, 2)$: Given the general infinitesimal displacement

$$d\mathbf{l} = dr \hat{\mathbf{r}} + r d\theta \hat{\boldsymbol{\theta}} + r \sin \theta d\phi \hat{\boldsymbol{\phi}}$$

and the gradient of T in spherical coordinates

$$\nabla T = (\cos \theta + \sin \theta \cos \phi) \hat{\mathbf{r}} + (-\sin \theta + \cos \theta \cos \phi) \hat{\boldsymbol{\theta}} + (-\sin \phi) \hat{\boldsymbol{\phi}}$$

(i) On the path OA :

$$\theta = \pi/2, \quad \phi = 0, \quad d\mathbf{l} = dr \hat{\mathbf{r}}; \quad (\nabla T) \cdot d\mathbf{l} = 1 dr$$

So

$$\int_{OA} (\nabla T) \cdot d\mathbf{l} = \int_0^2 1 dr = 2$$

(ii) On the path AB :

$$r = 2, \quad \theta = \pi/2, \quad d\mathbf{l} = 2 d\phi \hat{\boldsymbol{\phi}}; \quad (\nabla T) \cdot d\mathbf{l} = -2 \sin \phi d\phi$$

So

$$\int_{AB} (\nabla T) \cdot d\mathbf{l} = \int_0^{\pi/2} -2 \sin \phi d\phi = -2$$

(iii) On the path BC :

$$r = 2, \quad \phi = \pi/2, \quad d\mathbf{l} = 2 d\theta \hat{\boldsymbol{\theta}}; \quad (\nabla T) \cdot d\mathbf{l} = -2 \sin \theta d\theta$$

thus, the unit vectors are:

$$\begin{aligned}\hat{\mathbf{s}} &= \cos \phi \hat{\mathbf{x}} + \sin \phi \hat{\mathbf{y}} \\ \hat{\phi} &= -\sin \phi \hat{\mathbf{x}} + \cos \phi \hat{\mathbf{y}} \\ \hat{\mathbf{z}} &= \hat{\mathbf{z}}\end{aligned}$$

The cylindrical unit vectors in terms of $\hat{\mathbf{x}}$, $\hat{\mathbf{y}}$, and $\hat{\mathbf{z}}$ in matrix form:

$$\begin{pmatrix} \hat{\mathbf{s}} \\ \hat{\phi} \\ \hat{\mathbf{z}} \end{pmatrix} = \begin{pmatrix} \cos \phi & \sin \phi & 0 \\ -\sin \phi & \cos \phi & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} \hat{\mathbf{x}} \\ \hat{\mathbf{y}} \\ \hat{\mathbf{z}} \end{pmatrix}$$

Which is an orthogonal matrix $a = Qx$, so the Cartesian unit vectors is found by multiplying a by the transpose of Q :

$$\begin{pmatrix} \hat{\mathbf{x}} \\ \hat{\mathbf{y}} \\ \hat{\mathbf{z}} \end{pmatrix} = \begin{pmatrix} \cos \phi & -\sin \phi & 0 \\ \sin \phi & \cos \phi & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} \hat{\mathbf{s}} \\ \hat{\phi} \\ \hat{\mathbf{z}} \end{pmatrix}$$

or in vector form:

$$\begin{aligned}\hat{\mathbf{x}} &= \cos \phi \hat{\mathbf{s}} - \sin \phi \hat{\phi} \\ \hat{\mathbf{y}} &= \sin \phi \hat{\mathbf{s}} + \cos \phi \hat{\phi} \\ \hat{\mathbf{z}} &= \hat{\mathbf{z}}\end{aligned}$$

1.43 (a) Finding the divergence of the function

$$\mathbf{v} = s(2 + \sin^2 \phi) \hat{\mathbf{s}} + s \sin \phi \cos \phi \hat{\phi} + 3z \hat{\mathbf{z}}$$

$$\begin{aligned}\nabla \cdot \mathbf{v} &= \frac{1}{s} \frac{\partial}{\partial s}(s v_s) + \frac{1}{s} \frac{\partial v_\phi}{\partial \phi} + \frac{\partial v_z}{\partial z} \\ &= \frac{1}{s} \frac{\partial}{\partial s}(s(s(2 + \sin^2 \phi))) + \frac{1}{s} \frac{\partial}{\partial \phi}(s \sin \phi \cos \phi) + \frac{\partial}{\partial z}(3z) \\ &= 2(2 + \sin^2 \phi) + \cos^2 \phi - \sin^2 \phi + 3 \\ &= 7 + \sin^2 \phi + \cos^2 \phi = 8\end{aligned}$$

(b) Testing divergence theorem using a quarter cylinder of radius 2 and height 5 in quadrant I:
LHS: The volume elements is

$$dV = s \, ds \, d\phi \, dz,$$

so the volume integral is

$$\begin{aligned}\int_V (\nabla \cdot \mathbf{v}) \cdot dV &= \int_{s=0}^2 \int_{\phi=0}^{\pi/2} \int_{z=0}^5 8(s \, ds \, d\phi \, dz) \\ &= 8 \int_0^2 s \, ds \int_0^{\pi/2} d\phi \int_0^5 dz \\ &= 8(2)(\pi/2)(5) = 40\pi\end{aligned}$$

RHS: There are 5 surfaces: the top, bottom, and 3 sides.

(i) The top surface:

$$z = 5, \quad da = s \, ds \, d\phi \, \hat{\mathbf{z}}; \quad \mathbf{v} \cdot d\mathbf{a} = 15s \, ds \, d\phi$$

So

$$\oint_S \mathbf{v} \cdot d\mathbf{a} = \int_{\phi=0}^{\pi/2} \int_{s=0}^2 15s \, ds \, d\phi = 15 \int_0^{\pi/2} d\phi \int_0^2 s \, ds = 15\pi$$

(ii) The bottom surface:

$$z = 0, \quad d\mathbf{a} = s \, ds \, d\phi \, \hat{\mathbf{z}}; \quad \mathbf{v} \cdot d\mathbf{a} = 0 \quad \text{thus} \quad \oint_S \mathbf{v} \cdot d\mathbf{a} = 0$$

(iii) The surface on the xy plane:

$$\phi = \pi/2 \quad d\mathbf{a} = ds \, dz \, \hat{\phi}; \quad \mathbf{v} \cdot d\mathbf{a} = 0 \quad \text{thus} \quad \oint_S \mathbf{v} \cdot d\mathbf{a} = 0$$

(iv) The surface on the xz plane:

$$\phi = 0, \quad d\mathbf{a} = -ds \, dz \, \hat{\phi}; \quad \mathbf{v} \cdot d\mathbf{a} = 0 \quad \text{thus} \quad \oint_S \mathbf{v} \cdot d\mathbf{a} = 0$$

(v) The curved surface:

$$s = 2, \quad d\mathbf{a} = 2 \, d\phi \, dz \, \hat{\mathbf{s}}; \quad \mathbf{v} \cdot d\mathbf{a} = 4(2 + \sin^2 \phi) \, d\phi \, dz$$

So

$$\begin{aligned} \oint_S \mathbf{v} \cdot d\mathbf{a} &= \int_{z=0}^5 \int_{\phi=0}^{\pi/2} 4(2 + \sin^2 \phi) \, d\phi \, dz \\ &= 4 \int_0^5 dz \int_0^{\pi/2} (4(1/2) + \sin^2 \phi) \, d\phi \\ \text{using } \sin^2 \phi &= \frac{1 - \cos(2\phi)}{2} \\ &= 4 \int_0^5 dz \int_0^{\pi/2} (2 + (1 - \cos(2\phi))/2) \, d\phi \\ &= 10 \int_0^{\pi/2} (5 - \cos(2\phi)) \, d\phi \\ &= 10 \left(5\phi - \frac{\sin(2\phi)}{2} \Big|_0^{\pi/2} \right) = 25\pi \end{aligned}$$

So the total surface integral is

$$\oint_S \mathbf{v} \cdot d\mathbf{a} = 15\pi + 0 + 0 + 0 + 25\pi = 40\pi$$

which is the same as the volume integral, so the divergence theorem holds.

(c) The curl of \mathbf{v} in cylindrical coordinates is

$$\begin{aligned} \nabla \times \mathbf{v} &= \frac{1}{s} \begin{vmatrix} \hat{\mathbf{s}} & s\hat{\phi} & \hat{\mathbf{z}} \\ \frac{\partial}{\partial s} & \frac{\partial}{\partial \phi} & \frac{\partial}{\partial z} \\ v_s & (s)v_\phi & v_z \end{vmatrix} \\ &= \frac{1}{s} \begin{vmatrix} \hat{\mathbf{s}} & s\hat{\phi} & \hat{\mathbf{z}} \\ \frac{\partial}{\partial s} & \frac{\partial}{\partial \phi} & \frac{\partial}{\partial z} \\ s(2 + \sin^2 \phi) & (s)s \sin \phi \cos \phi & 3z \end{vmatrix} \\ &= 0\hat{\mathbf{s}} - \frac{1}{s}(0 - 0) + \frac{1}{s}(2s \sin \phi \cos \phi - 2s \sin \phi \cos \phi) \\ &= 0 \end{aligned}$$

1.48 (a)

$$\int (r^2 + \mathbf{r} \cdot \mathbf{a} + a^2) \delta^3(\mathbf{r} - \mathbf{a}) dV = a^2 + \mathbf{a} \cdot \mathbf{a} + a^2 = 3a^2$$

(b) Given V is a cube of side 2 centered at the origin and $\mathbf{b} = 4\hat{\mathbf{y}} + 3\hat{\mathbf{z}}$:

$$\begin{aligned} \int_V |\mathbf{r} - \mathbf{b}|^2 \delta^3(5\mathbf{r}) dV &= \int_0^1 \int_0^1 \int_0^1 |\mathbf{r} - \mathbf{b}|^2 \delta(5x) \delta(5y) \delta(5z) dx dy dz \\ &= \frac{1}{5^3} |\mathbf{b}|^2 = \frac{1}{5^3} (4^2 + 3^2) = \frac{1}{5} \end{aligned}$$

(c) Given V is a sphere of radius 6 centered at the origin and $\mathbf{c} = 5\hat{\mathbf{x}} + 3\hat{\mathbf{y}} + 2\hat{\mathbf{z}}$:

The magnitude $c = \sqrt{5^2 + 3^2 + 2^2} = \sqrt{38}$, is outside the sphere (magnitude $r = \sqrt{36}$). Therefore,

$$\int_V [r^4 + r^2(\mathbf{r} \cdot \mathbf{c}) + c^4] \delta^3(\mathbf{r} - \mathbf{c}) dV = 0$$

(d) Given V is a sphere of radius 1.5 centered at $(2, 2, 2)$ and

$$\mathbf{d} = (1, 2, 3), \quad \mathbf{e} = (3, 2, 1)$$

Checking if the delta function is inside the sphere:

$$|\mathbf{e} - (2, 2, 2)| = \sqrt{1 + 0 + 1} = \sqrt{2} < 1.5$$

so the delta function is inside the sphere and the integral is

$$\begin{aligned} \int_V \mathbf{r} \cdot (\mathbf{d} - \mathbf{r}) \delta^3(\mathbf{e} - \mathbf{r}) dV &= \int_V \mathbf{r} \cdot (\mathbf{d} - \mathbf{r}) \delta^3(\mathbf{r} - \mathbf{e}) dV \\ &= \mathbf{e} \cdot (\mathbf{d} - \mathbf{e}) \\ &= (3, 2, 1) \cdot (-2, 0, 2) = -4 \end{aligned}$$

1.49 Evaluating the integral

$$J = \int_V e^{-r} \left(\nabla \cdot \frac{\hat{\mathbf{r}}}{r^2} \right) dV$$

where V is a sphere of radius R centered at the origin.

(i)

$$J = \int_V e^{-r} 4\pi \delta^3(\mathbf{r}) dV = 4\pi e^0 = 4\pi$$

(ii) Using the product rule of divergence:

$$\nabla \cdot (f\mathbf{A}) = f(\nabla \cdot \mathbf{A}) + \mathbf{A} \cdot (\nabla f)$$

integrating over a volume and using the divergence theorem:

$$\int \nabla \cdot (f\mathbf{A}) dV = \int f(\nabla \cdot \mathbf{A}) dV + \int \mathbf{A} \cdot (\nabla f) dV = \oint f\mathbf{A} \cdot d\mathbf{a}$$

or

$$\int_V f(\nabla \cdot \mathbf{A}) dV = - \int_V \mathbf{A} \cdot (\nabla f) dV + \oint f\mathbf{A} \cdot d\mathbf{a}$$

where

$$f = e^{-r} \quad \text{and} \quad \mathbf{A} = \frac{\hat{\mathbf{r}}}{r^2}$$

The gradient of a scalar:

$$\mathbf{F}_3 = -\nabla V_3$$

so

$$\frac{\partial V_3}{\partial x} = -yz, \quad \frac{\partial V_3}{\partial y} = -zx, \quad \frac{\partial V_3}{\partial z} = -xy$$

A particular solution is

$$V_3 = -xyz$$

The curl of a vector (a little harder):

$$\mathbf{F}_3 = \nabla \times \mathbf{V}_3$$

so

$$\frac{\partial A_z}{\partial y} - \frac{\partial A_y}{\partial z} = yz, \quad \frac{\partial A_x}{\partial z} - \frac{\partial A_z}{\partial x} = zx, \quad \frac{\partial A_y}{\partial x} - \frac{\partial A_x}{\partial y} = xy$$

integrating the first equation with respect to y :

$$\begin{aligned} \frac{\partial A_z}{\partial y} &= yz + \frac{\partial A_y}{\partial z} \\ \int \frac{\partial A_z}{\partial y} dy &= \int yz dy + \int \frac{\partial A_y}{\partial z} dy \\ A_z &= \frac{1}{2}y^2z + f \end{aligned}$$

where the function f is an arbitrary function. Integrating the first equation again but with respect to z gives:

$$A_y = -\frac{1}{2}yz^2 + g$$

Repeating for all three equations:

$$\begin{aligned} A_z &= \frac{1}{2}y^2z + f \quad \text{and} \quad A_y = -\frac{1}{2}yz^2 + g \\ A_x &= \frac{1}{2}z^2x + h \quad \text{and} \quad A_z = -\frac{1}{2}zx^2 + i \\ A_y &= \frac{1}{2}x^2y + j \quad \text{and} \quad A_x = -\frac{1}{2}xy^2 + k \end{aligned}$$

a particular solution can be found by setting $f = g = h = i = j = k = 0$, and each component is a linear combination of the two possible solutions:

$$\begin{aligned} 2A_x &= \frac{1}{2}z^2x + f - \frac{1}{2}xy^2 + k \\ 2A_x &= \frac{1}{2}z^2x - \frac{1}{2}xy^2 \\ A_x &= \frac{1}{4}(z^2x - xy^2) \end{aligned}$$

and similarly for A_y and A_z gives the vector potential:

$$\mathbf{A} = \frac{1}{4}[x(z^2 - y^2)\hat{\mathbf{x}} + y(x^2 - z^2)\hat{\mathbf{y}} + z(y^2 - x^2)\hat{\mathbf{z}}]$$

1.51 (d) \rightarrow (a): Given $\mathbf{F} = -\nabla V$, the curl of the gradient is always zero;

$$\nabla \times \mathbf{F} = \nabla \times (-\nabla V) = 0$$

(a) \rightarrow (c): From Stokes' theorem;

$$\int_S (\nabla \times \mathbf{F}) \cdot d\mathbf{a} = \oint \mathbf{F} \cdot d\mathbf{l} = 0$$

(c) \rightarrow (b):

$$\oint \mathbf{F} \cdot d\mathbf{l} = \int_a^b \mathbf{F} \cdot d\mathbf{l} - \int_a^b \mathbf{F} \cdot d\mathbf{l} = 0$$

where the integrals are two different paths but equal, and thus independent of path. (b) \rightarrow (c) and (c) \rightarrow (a) are just the same steps in reverse.

1.52 (d) \rightarrow (a): Given $\mathbf{F} = \nabla \times A$, the divergence of curl is always zero;

$$\nabla \cdot (\nabla \times A) = 0$$

(a) \rightarrow (c): From the divergence theorem;

$$\int_V (\nabla \cdot \mathbf{F}) dV = \oint \mathbf{F} \cdot d\mathbf{a} = 0$$

(c) \rightarrow (b):

$$\oint \mathbf{F} \cdot d\mathbf{a} = \int \mathbf{F} \cdot d\mathbf{a}_1 + \int \mathbf{F} \cdot d\mathbf{a}_2 = 0$$

where $\mathbf{a}_2 = -\mathbf{a}_1$ and the integrals are two different surfaces but equal, and thus depend only on the boundary. (b) \rightarrow (c) and (c) \rightarrow (a) are just the same steps in reverse.

1.53 (a) From Problem 1.15,

$$\begin{aligned}\mathbf{v}_a &= x^2 \hat{\mathbf{x}} + 3xz^2 \hat{\mathbf{y}} - 2xz \hat{\mathbf{z}} \\ \mathbf{v}_b &= xy \hat{\mathbf{x}} + 2yz \hat{\mathbf{y}} + 3zx \hat{\mathbf{z}} \\ \mathbf{v}_c &= y^2 \hat{\mathbf{x}} + (2xy + z^2) \hat{\mathbf{y}} + 2yz \hat{\mathbf{z}}\end{aligned}$$

and the curl of each vector field:

$$\begin{aligned}\nabla \times \mathbf{v}_a &\neq 0 \\ \nabla \times \mathbf{v}_b &\neq 0 \\ \nabla \times \mathbf{v}_c &= 0\end{aligned}$$

Thus \mathbf{v}_c can be written as a gradient of some scalar potential $\mathbf{v}_c = -\nabla V$:

$$\begin{aligned}\frac{\partial V}{\partial x} &= -y^2; & V &= -y^2 x + f \\ \frac{\partial V}{\partial y} &= -(2xy + z^2); & V &= -y^2 x - yz^2 + g \\ \frac{\partial V}{\partial z} &= -2yz; & V &= -yz^2 + h\end{aligned}$$

where f , g , and h are arbitrary functions, thus one solution can be found setting $g = 0$ and solving for f or h ;

$$V = -y^2 x - yz^2$$

1.61 (a) Show that

$$\int_V (\nabla T) dV = \oint_S T d\mathbf{a}$$

Letting $\mathbf{v} = \mathbf{c}T$ where \mathbf{c} is a constant: using the product rule of divergence:

$$\begin{aligned}\nabla \cdot (f\mathbf{A}) &= f(\nabla \cdot \mathbf{A}) + \mathbf{A} \cdot (\nabla f) \\ \nabla \cdot (T\mathbf{c}) &= T(\nabla \cdot \mathbf{c}) + \mathbf{c} \cdot (\nabla T) \\ \nabla \cdot (T\mathbf{c}) &= \mathbf{c} \cdot (\nabla T)\end{aligned}$$

where $\nabla \cdot \mathbf{c} = 0$ since \mathbf{c} is a constant. Integrating over a volume and using the divergence theorem:

$$\int_V \nabla \cdot (T\mathbf{c}) dV = \int_V \mathbf{c} \cdot (\nabla T) dV = \oint_S T\mathbf{c} \cdot d\mathbf{a}$$

or moving the constant \mathbf{c} outside the integral:

$$\begin{aligned}\mathbf{c} \int_V \nabla T dV &= \mathbf{c} \oint_S T \cdot d\mathbf{a} \\ \int_V \nabla T dV &= \oint_S T \cdot d\mathbf{a}\end{aligned}$$

(b) Show that

$$\int_V (\nabla \times \mathbf{v}) dV = - \oint_S \mathbf{v} \times d\mathbf{a}$$

replacing \mathbf{v} with $\mathbf{v} \times \mathbf{c}$ in the divergence theorem:

$$\int_V \nabla \cdot (\mathbf{v} \times \mathbf{c}) dV = \oint_S (\mathbf{v} \times \mathbf{c}) \cdot d\mathbf{a} \quad (1.1)$$

and using the product rule of divergence:

$$\begin{aligned}\nabla \cdot (\mathbf{v} \times \mathbf{c}) &= \mathbf{c} \cdot (\nabla \times \mathbf{v}) - \mathbf{v} \cdot (\nabla \times \mathbf{c}) \\ \nabla \cdot (\mathbf{v} \times \mathbf{c}) &= \mathbf{c} \cdot (\nabla \times \mathbf{v})\end{aligned}$$

where $\nabla \times \mathbf{c} = 0$ since \mathbf{c} is a constant. So subbing into the divergence theorem and moving the constant \mathbf{c} outside the integral:

$$\mathbf{c} \int_V (\nabla \times \mathbf{v}) dV = \oint_S (\mathbf{v} \times \mathbf{c}) \cdot d\mathbf{a}$$

and using the scalar triple product identity:

$$(\mathbf{v} \times \mathbf{c}) \cdot d\mathbf{a} = \mathbf{v} \cdot (\mathbf{c} \times d\mathbf{a}) = \mathbf{c} \cdot (d\mathbf{a} \times \mathbf{v}) = -\mathbf{c} \cdot (\mathbf{v} \times d\mathbf{a})$$

so

$$\begin{aligned}\mathbf{c} \int_V (\nabla \times \mathbf{v}) dV &= -\mathbf{c} \oint_S (\mathbf{v} \times d\mathbf{a}) \\ \int_V (\nabla \times \mathbf{v}) dV &= - \oint_S (\mathbf{v} \times d\mathbf{a})\end{aligned}$$

(c) [Green's identity] Show that

$$\int_V [T\nabla^2 U dV + (\nabla T) \cdot (\nabla U)] dV = \oint_S (T\nabla U) \cdot d\mathbf{a}$$

letting $\mathbf{v} = T\nabla U$ in the divergence theorem:

$$\int_V \nabla \cdot (T\nabla U) dV = \oint_S (T\nabla U) \cdot d\mathbf{a}$$

using the product rule of divergence:

$$\nabla \cdot (T\nabla U) = T(\nabla \cdot \nabla U) + (\nabla U) \cdot (\nabla T)$$

$$\nabla \cdot (T\nabla U) = T(\nabla^2 U) + (\nabla T) \cdot (\nabla U)$$

where $\nabla \cdot \nabla U = \nabla^2 U$ and the dot product is commutative. Subbing into the divergence theorem:

$$\int_V [T\nabla^2 U dV + (\nabla T) \cdot (\nabla U)] dV = \oint_S (T\nabla U) \cdot d\mathbf{a}$$

(d) [Green's second identity] Show that

$$\int_V (T\nabla^2 U - U\nabla^2 T) dV = \oint_S (T\nabla U - U\nabla T) \cdot d\mathbf{a}$$

using the product rule of divergence:

$$\nabla \cdot (U\nabla T) = U\nabla^2 T + (\nabla T) \cdot (\nabla U)$$

and letting $\mathbf{v} = U\nabla T$ in the divergence theorem:

$$\int_V \nabla \cdot (U\nabla T) dV = \oint_S (U\nabla T) \cdot d\mathbf{a}$$

$$\int_V [U\nabla^2 T + (\nabla T) \cdot (\nabla U)] dV = \oint_S (U\nabla T) \cdot d\mathbf{a}$$

and taking the difference with the result from (c):

$$\begin{aligned} \oint_S (T\nabla U) \cdot d\mathbf{a} - \oint_S (U\nabla T) \cdot d\mathbf{a} &= \oint_S (T\nabla U - U\nabla T) \cdot d\mathbf{a} \\ \int_V [T\nabla^2 U dV - U\nabla^2 T] dV &= \oint_S (T\nabla U - U\nabla T) \cdot d\mathbf{a} \end{aligned}$$

where $(\nabla T) \cdot (\nabla U)$ cancels out on the left hand side.

(e) Show that

$$\int_S \nabla T \times d\mathbf{a} = - \oint_P T d\mathbf{l}$$

letting $\mathbf{v} = \mathbf{c}T$ in Stokes' theorem:

$$\int_S \nabla \times (\mathbf{c}T) \cdot d\mathbf{a} = \oint_P (\mathbf{c}T) \cdot d\mathbf{l}$$

using the product rule of curl:

$$\nabla \times (\mathbf{c}T) = T(\nabla \times \mathbf{c}) - \mathbf{c} \times (\nabla T)$$

$$\nabla \times (\mathbf{c}T) = -\mathbf{c} \times (\nabla T)$$

where $\nabla \times \mathbf{c} = 0$ since \mathbf{c} is a constant. And subbing into Stokes' theorem:

$$\int_S (-\mathbf{c} \times (\nabla T)) \cdot d\mathbf{a} = \oint_P (\mathbf{c}T) \cdot d\mathbf{l}$$

and using the scalar triple product identity

$$(-\mathbf{c} \times (\nabla T)) \cdot d\mathbf{a} = -\mathbf{c} \cdot (\nabla T \times d\mathbf{a})$$

thus the constant \mathbf{c} can move out of the integral and cancel out giving the result

$$\int_S \nabla T \times d\mathbf{a} = - \oint_P T d\mathbf{l}$$

Thus the vector area of the two surfaces are equal as long as the boundary is the same.

(d) Show that the integral around the boundary line

$$\mathbf{a} = \frac{1}{2} \oint_S \mathbf{r} \times d\mathbf{l}$$

A cone subtended by the loop at the origin (vertex at the origin) can be split into infinitesimal triangular wedges with base $d\mathbf{l}$ and height \mathbf{r} , and the area of each wedge is

$$dA = \frac{1}{2} \mathbf{r} \times d\mathbf{l}$$

since the cross product is the area of the parallelogram formed by the two vectors or in this case the area of the triangle. Summing over all the wedges along the closed loop gives the total area

$$\frac{1}{2} \oint_S \mathbf{r} \times d\mathbf{l} = \mathbf{a} = \oint_S dA$$

(e) Show that

$$\oint (\mathbf{c} \cdot \mathbf{r}) d\mathbf{l} = \mathbf{a} \cdot \mathbf{c}$$

for any constant vector \mathbf{c} . Letting $T = \mathbf{c} \cdot \mathbf{r}$ from the result in 1.61(e):

$$\begin{aligned} \int \nabla T \times d\mathbf{a} &= - \oint T d\mathbf{l} \\ \int \nabla(\mathbf{c} \cdot \mathbf{r}) \times d\mathbf{a} &= - \oint (\mathbf{c} \cdot \mathbf{r}) d\mathbf{l} \end{aligned}$$

using the product rule of gradients:

$$\nabla(\mathbf{c} \cdot \mathbf{r}) = \mathbf{c} \times (\nabla \times \mathbf{r}) + \mathbf{r} \times (\nabla \times \mathbf{c}) + (\mathbf{c} \cdot \nabla) \mathbf{r} + (\mathbf{r} \cdot \nabla) \mathbf{c}$$

where $\mathbf{r} = (x, y, z)$; $\nabla \times \mathbf{r} = 0$, $\nabla \times \mathbf{c} = 0$, $\nabla \mathbf{c} = 0$, and

$$(\mathbf{c} \cdot \nabla) \mathbf{r} = \mathbf{c}$$

So

$$\nabla(\mathbf{c} \cdot \mathbf{r}) = \mathbf{c}$$

therefore

$$\begin{aligned} - \oint (\mathbf{c} \cdot \mathbf{r}) d\mathbf{l} &= \int \mathbf{c} \times d\mathbf{a} \\ &= \mathbf{c} \times \int d\mathbf{a} \\ &= \mathbf{c} \times \mathbf{a} \\ &= -\mathbf{a} \times \mathbf{c} \\ \oint (\mathbf{c} \cdot \mathbf{r}) d\mathbf{l} &= \mathbf{a} \times \mathbf{c} \end{aligned}$$

1.63 (a) Finding the divergence of

$$\mathbf{v} = \frac{\hat{\mathbf{r}}}{r}$$

$$\nabla \cdot \mathbf{v} = \frac{1}{r^2} \frac{\partial}{\partial r} \left(r^2 \frac{1}{r} \right) = \frac{1}{r^2}$$

