Ludwig-Maximilians-Universität München

Bachelors Thesis

The role of excited atomic states in multiphoton ionization

Johannes Porsch



Abstract

Contents

Al	Abstract												
List of Figures													
1	Introduction	3											
2	Theory 2.1 Basic Formalism	4 5 6 7 8											
3	3 Conclusion												
\mathbf{A}	Appendix A	10											
В	Appendix B	11											
Bi	ibliography	12											

List of Figures

2.1	Sine function					 	_					_	_				_	_						_		_			_	_		4
	DIIIC IUIICUIOII	 	•	•	•	 	•	•	•	•	•	•	•	•	•	•	•	•		•	•	•	•	•	•	•	•	•	•	•	•	1

1. Introduction

2. Theory

$$\partial_t u = \mathcal{H}(t)\lambda \tag{2.1}$$




Figure 2.1: Sine function

2.1 Basic Formalism

Basic Definitions of schröfinger qe, light dyson series, and sotrong field s matrix

2.2 Strong Field Approximation

clear defintition of the strong field approximation, and the assumptions that are made.

2.3 Strong Field Ionization

Derivation of

$$\lim_{t \to \infty} |\Psi(t)\rangle = -i \int d^3p |\mathbf{p}\rangle \int_{-\infty}^{\infty} dt' e^{-\frac{i}{2} \int_{t'}^{\infty} [\mathbf{p} + \mathbf{A}(t')]^2 dt'} e^{iI_{\mathbf{p}}t'} \langle \mathbf{p} + \mathbf{A}(t') | \hat{\mathbf{d}} \cdot \mathbf{E}(t') | \Psi_0 \rangle \qquad (2.2)$$

2.4 Multiphoton Ionization

Different types of Ionization, tunneling Ionization, multiphoton

3. Conclusion

Lorem ipsum dolor sit amet, consetetur sadipscing elitr, sed diam nonumy eirmod tempor invidunt ut labore et dolore magna aliquyam erat, sed diam voluptua. At vero eos et accusam et justo duo dolores et ea rebum. Stet clita kasd gubergren, no sea takimata sanctus est Lorem ipsum dolor sit amet. Lorem ipsum dolor sit amet, consetetur sadipscing elitr, sed diam nonumy eirmod tempor invidunt ut labore et dolore magna aliquyam erat, sed diam voluptua. At vero eos et accusam et justo duo dolores et ea rebum. Stet clita kasd gubergren, no sea takimata sanctus est Lorem ipsum dolor sit amet.

$$\partial \mathbf{A} = \mathfrak{B}$$

$$\int_{\mathbb{R}^d} |f(x)|^2 \, \mathrm{d}x = \int_{\mathbb{R}^d} |\mathcal{F}f(\xi)|^2 \, \mathrm{d}\xi \tag{3.1}$$

$$i\partial_t u = \mathcal{H}(t) |a\rangle \lambda$$
 (3.2)

for this calculation [1] was used

A. Appendix A

B. Appendix B

Bibliography

[1] Seung Beom Park, Kyungseung Kim, Wosik Cho, Sung In Hwang, Igor Ivanov, Chang Hee Nam, and Kyung Taec Kim. Direct sampling of a light wave in air. *Optica*, 5(4):402–408, Apr 2018.