NEUR 411 · NEUROETHOLOGY · FALL 2026

Lecture 91

Sea lions: brain/neuro anatomy/organization, rhythmic entrainment, auditory–motor coordination, learned commands, and tests of beat synchronization

Date TBD

Jordan Lopez, Ph.D.
Professor of Neurobiology & Biomedical Engineering





Photo: California sea lion Ronan at UC Santa Cruz's Long Marine Laboratory. Photo by Colleen Reichmuth; NOAA/NMFS 23554.

Ronan learns to align movement with sound

Rhythmic entrainment is the adjustment of a repeating movement to an external rhythm. Ronan, a California sea lion, learned to move her head up and down while hearing regularly spaced sounds. The lowest point of each movement supplied a repeatable timing marker.

Cook and colleagues trained this behavior at 80 and 120 beats per minute, then tested other rates and musical recordings. A movement that stays aligned across unfamiliar rates requires flexible timing rather than the repetition of one memorized motor speed.

Ronan lived at the University of California, Santa Cruz. The experimental task underlying her public reputation as a dancing sea lion combined a controlled sound sequence, concealed trainers, video measurements, and food reinforcement for appropriately timed behavior.




Cook et al. (2013), Fig. 1C. Head-bob timing after training at 80 bpm.

Matching tempo differs from matching the beat

Tempo is the repetition rate of the sound sequence, expressed in beats per minute. Matching tempo means producing the same average number of movements per unit time. Two repeating events can share a tempo while occurring at different points within each cycle.

Phase describes the position of a movement relative to the sound cycle. Synchronization requires a stable timing relationship, ideally with the selected movement landmark near sound onset. The 2013 study measured both the movement rate and its alignment with individual beats.

At different unfamiliar tempos, Ronan maintained organized phase relationships even when her head reached its lowest point before or after the sound. These offsets distinguish successful coordination with the repeating stimulus from exact coincidence at the beat itself.




Cook et al. (2013), Fig. 2. Movement timing at five unfamiliar tempos.

A folded cortex overlies a large cerebellum

Neuroanatomy is the organization of the nervous system and its structures. **Cerebral cortex** is the outer sheet of the cerebrum. Sawyer and colleagues examined a juvenile California sea-lion brain whose cortical surface contained numerous folds. **Gyri** are the raised folds; **sulci** are the grooves separating them. The two hemispheres had similar but nonidentical groove patterns.

The fixed specimen was 10.2 cm long and 10.1 cm wide near its posterior end, compared with 6.0 cm across the anterior portion. These are dimensions of the preserved juvenile specimen, rather than measurements of Ronan or a standard adult brain.

A large **cerebellum**, a brain structure involved in movement coordination, lay beneath the posterior cerebral region. The study localized sensory cortex using tissue stains and anatomical landmarks. Folding alone did not locate functional regions; microscopic organization supplied the boundaries of the somatosensory area.




Sawyer et al. (2016), Fig. 7G–I. Dorsal, isometric, and ventral views of a juvenile sea-lion brain.

Whisker follicles connect touch to the trigeminal nerve

Vibrissae are specialized whiskers whose follicles receive sensory innervation. The examined sea lions had 38 whiskers arranged in six rows on each cheek pad. Their facial array provides many separate points at which contact or water movement can deform a whisker.

A stained follicle contained a blood-filled cavernous sinus and a deep vibrissal nerve entering its base. This structure differs from an ordinary coat-hair follicle. Whisker deformation is therefore delivered to an innervated sensory organ embedded in the skin rather than detected by the hair shaft itself.

The **trigeminal nerve** carries facial somatosensory signals, meaning information about touch and related bodily sensation. Its thick sensory root lay beside a separate, smaller motor root. **Axons** are nerve-cell fibers that conduct signals; **myelin** is their insulating sheath. Microscopy identified thinly and thickly myelinated axons in the pathway carrying facial information into the brainstem.




Sawyer et al. (2016), Fig. 2A–F. Whiskers, an innervated follicle, and sensory and motor roots.

Brainstem modules organize facial sensory input

The **principal sensory trigeminal nucleus** is a brainstem relay for facial somatosensory information. In sea-lion tissue, this nucleus contained clusters separated by relatively cell-sparse partitions. Related modules also occurred in the spinal trigeminal nuclei, which extend the facial representation through the brainstem.

Nissl staining reveals neuronal cell bodies. **Cytochrome oxidase staining** marks metabolic organization, while vesicular glutamate transporter 1 staining identified dense afferent terminal regions. **Glutamate** is a neurotransmitter, a chemical signal released at synapses. This transporter packages it in presynaptic vesicles. The three preparations converged on a modular architecture instead of a uniform mass of tissue.

The arrangement resembled the facial whisker array, especially the six-row organization in the interpolaris subdivision. Individual sections did not yield an exact whisker-to-module assignment. The anatomical finding is a partitioned facial sensory representation, with dense incoming terminals concentrated within its modules.




Sawyer et al. (2016), Fig. 4I–K. Principal sensory nucleus stained for cells, metabolism, and VGLUT1.

The thalamus separates face and body representations

The **thalamus** contains relays between ascending sensory pathways and cerebral cortex. The sea-lion **ventral posterior nucleus** was distinguished from surrounding tissue by large neuronal cell bodies and dense staining for cytochrome oxidase and vesicular glutamate transporter 2.

Its **ventroposterior medial subdivision**, or VPM, occupied a large part of the nucleus and corresponds to facial representation in the mammalian somatosensory pathway. The neighboring **ventroposterior lateral subdivision**, or VPL, corresponds to bodily representation. Their boundaries changed across successive sections rather than forming identical shapes at every depth.

Lightly stained partitions divided the nucleus internally. The sea-lion preparation established these subdivisions anatomically, without a physiological point-by-point body map. Facial sensation thus has an extensive thalamic territory between its brainstem representation and the cortical sensory region.




Sawyer et al. (2016), Fig. 6B–D. Matched sections identify the somatosensory thalamic subdivisions.

Sensory cortical layer 4 receives dense input

Primary somatosensory cortex, or S1, is a cerebral region receiving bodily sensory information. Sawyer and colleagues localized a presumptive S1 region just anterior to the major vertical fissure using its microscopic organization and position within the folded sea-lion cortex.

Cortical layers are bands that differ in neuronal populations and connections. In this region, layer 4 contained densely packed cells and stained strongly for cytochrome oxidase and vesicular glutamate transporter 2. Layer 5 was relatively cell-sparse, whereas layer 6 was thick and cell-dense.

The VGLUT2-rich layer identifies a concentration of glutamatergic input terminals associated with primary sensory cortex. Tissue organization therefore connected a macroscopically folded surface with a specific sensory territory. Reconstructing successive stained sections placed that territory relative to the coronal and suprasylvian sulci.




Sawyer et al. (2016), Fig. 8E–G. Layer-specific metabolic, neuronal, and VGLUT2 staining.

Brain rhythms change with arousal state

Electroencephalography, or EEG, records voltage fluctuations through electrodes near the head. Williams and colleagues adapted the electrode arrangement to California sea lions and identified patterns in neurologically normal animals alongside recordings from animals with neurological disease.

Wakeful recordings included low-amplitude **beta activity**, faster than 13 Hz, interspersed with **alpha activity** at 8–13 Hz. Slow-wave sleep contained high-amplitude **delta activity**, slower than 4 Hz, and sleep spindles around 12–14 Hz. The same nervous system can therefore produce different temporal activity patterns across states.

These EEG rhythms are electrical activity patterns, whereas Ronan’s entrainment is a measured relationship between movement and an external sound sequence. The distinction separates physiological frequency from behavioral beat keeping. Sedation and arousal were tracked because they altered the EEG state being interpreted.




Williams et al. (2023), Fig. 2A–B. Sea-lion EEG during wakefulness and slow-wave sleep.

Touch exploration changes with the sensory task

Active sensing means moving a sensory organ to acquire information relevant to a task. Lo, a California sea lion, used different head and whisker movements when discriminating texture, size, or visual brightness. Blindfolds restricted the texture and size tasks to tactile exploration.

For texture, Lo swept laterally across the surfaces. For size, she moved toward the object edges and made larger whisker movements. Average whisker excursions were about 34° for texture and 47° for size, compared with 24° during visual brightness discrimination.

The target remained the intermediate stimulus among two distractors. A visual control reduced the need to explore by touch, and head and whisker movements declined. Task-specific movement therefore changed how the sensory surface was sampled, linking perceptual demands to the motor strategy used to solve them.



Milne et al. (2021), Fig. 1A–B. Lo explores texture and size with different whisker trajectories.



Milne et al. (2021), Movie 1. Lo explores texture, size, and visual brightness.

A learned command initiates sensory-guided choice

A **learned command** is a cue associated with a trained action or task. In Lo's discrimination experiment, the trainer used "find it" to initiate exploration after the stimuli had been placed underwater. The command started a task whose solution still depended on information from the objects.

Lo indicated her choice by resting her nose on the selected stimulus for more than 3 s and retracting her whiskers. Correct choices produced a whistle followed by fish. The whistle served as a **conditioned reinforcer**, a previously learned signal marking success before the primary food reward arrived.

Stimulus positions changed between trials, so the same command could not be solved by approaching one permanent location. After learning, decisions during exploration took less than 1 s and accuracy approached complete target identification. The command controlled task initiation, while sensory discrimination controlled which object was selected.




Milne et al. (2021), Fig. 1C. Lo selects a brightness target with reduced tactile exploration.

Whisker adjustments precede head corrections

Ball balancing requires continuous adjustments as the supported object changes orientation. Milne and Grant recorded California sea lions named Gina, Anya, and Elmo performing this familiar trained action. The animals received ball-balancing commands and maintained the action until a trainer marked completion.

Whisker asymmetry describes the difference between average whisker angles on the two sides of the face. Changes in this measure followed ball orientation more closely and sooner than changes in head orientation. The reported average delays were approximately 27 ms for whiskers and 129 ms for the head.

Balancing therefore recruits control of the sensory array as well as movement of the larger support surface. Whisker movements need not merely follow the head passively. The supplied recording preserves the brief analyzed interval at its original speed, in which ball position and whisker configuration change together.



Milne and Grant (2014), Fig. 1b–c. Ball, head, and whisker positions during balancing.




Milne and Grant (2014), ESM 1. The original brief ball-balancing analysis interval.

Banking turns recruit the hindflippers


Sea lions propel ordinary forward swimming with their foreflippers, while the hindflippers can trail behind. A **banking turn** combines rolling the body with changing the swimming direction. Leahy and colleagues trained sea lions to complete 180° turns beneath an underwater viewing window.

At turn onset, the animal abducted its foreflippers, moving them away from the body, and rolled approximately 90°. The hindflippers spread and rotated, expanding their surface area. Their position behind the body's center of gravity allows a hydrodynamic force to influence body orientation as well as the path of travel.

The hindflippers' average **angle of attack**, their orientation relative to the approaching flow, was about 28° during turning. They remained spread during the maneuver and were adducted, brought together, as forward swimming resumed. Coordinated control therefore changes the role of an appendage with the behavioral requirement.



Leahy et al. (2021), Fig. 3A, E, I. Successive stages of a sea-lion banking turn.




Leahy et al. (2021), Movie 3. A trained sea lion executes a 180° left banking turn.

Leaping alternates propulsion and body stabilization

Porpoising is a swimming pattern that repeatedly carries the body above and below the water surface. A trained California sea lion performed consecutive leaps while cameras recorded aerial and submerged phases. The full cycle lasted about 1.5 s in the analyzed captive performance.

The foreflippers generated propulsive strokes during submerged swimming. The hindflippers spread beneath the water and helped stabilize the posterior body region. At the fully aerial crest, the flippers were held against the body or folded rather than continuously generating force against water.

Water entry began with the nose along a curved path, followed by repositioning the foreflippers and hindflippers for the submerged phase. Hindflipper angle of attack during the analyzed submerged phase averaged about 15°. The coordinated cycle thus changes the timing and orientation of several body parts as the surrounding medium changes.



Leahy et al. (2021), Fig. 1. Sea-lion hindflipper geometry and spread configuration.




Leahy et al. (2021), Movie 1. A captive sea lion completes a porpoising bout.

Wild porpoising retains the coordinated movement pattern

Wild California sea lions in Monterey Bay also executed repeated porpoising leaps. The field recordings and the trained-animal recordings supplied different contexts for the same broad locomotor pattern. Both involved alternating aerial movement with submerged propulsion and a changing hindflipper posture.

In the field sequence, the legs generally remained more extended during the aerial phase, and the hindflippers cleared the water. The reported average leap length was 0.81 body lengths and the aerial height was 0.06 body lengths. Expressing distances in body lengths permits comparison across differently sized animals.

The captive sea lion made longer and higher leaps and used a different entry angle. Training conditions can therefore change the scale of a movement without eliminating the coordinated pattern. The original field movie supplies the natural locomotor comparison; these measurements concern body control rather than synchronization to an acoustic pulse.



Leahy et al. (2021), Fig. 3D, H, L. Changing flipper configuration during a maneuver.




Leahy et al. (2021), Movie 2. Wild sea lions porpoise in Monterey Bay.

Musical pulse guides movement beyond the training sound

A musical recording contains overlapping instruments and events between the regular beats. Ronan coordinated her head movements with Everybody at about 108 bpm and Boogie Wonderland at 130 bpm. Keeping the repeated response organized required selecting a temporal regularity within this more complex acoustic material.

The songs had different exposure histories. Everybody received explicit training after earlier unsuccessful exposure. Boogie Wonderland had been heard briefly before the later transfer test, without that exposure producing entrained bobbing. Prior exposure and reinforced successful performance are distinct learning events.

After training at the native tempo of Boogie Wonderland, Ronan also followed altered versions spanning 104–143 bpm. The same musical material guided different movement rates. Flexibility across these versions links the trained action to the current pulse instead of one fixed speed learned from a familiar recording.




Cook et al. (2013), Fig. 4A–B. Head-bob timing with Everybody and Boogie Wonderland.

A dancing reputation rests on controlled experiments

The public description of Ronan as a dancing sea lion refers to a trained head-bob response measured against sound timing. Her scientific contribution was not simply that she moved while music played, but that her movement adapted across rates and persisted through controlled tests.

Novel tempos, musical recordings, silent omissions, and extended trials each challenged a different simple explanation of the behavior. Together they connected the repeating motor response to temporal regularities that varied independently of the original two training examples.

Ronan's study site was Long Marine Laboratory at UC Santa Cruz. The original researchers' recordings are the relevant evidence for her timing performance; a publicity recording supplies behavioral context, while quantified trial measurements establish the accuracy, consistency, and transfer of that coordination.



Cook et al. (2013), Fig. 6E–F. Extended coordination with music at 137 and 124 bpm.




Cook et al. (2013), official UCSC study video. Ronan transfers trained synchronization to music.

Perturbations separate phase from period correction

A **phase perturbation** shifts where the sound sequence falls in time while preserving its eventual repetition rate. A **tempo perturbation** changes the interval between successive sounds. These manipulations challenge different components of an already coordinated movement.

Rouse and colleagues introduced one unexpected change during a trial, after a stable initial run of beats. Ronan then had to modify the movement she was already producing. The response immediately after the change exposed the dynamics of correction.

Period is the duration of one complete cycle. A repeating motor process can correct timing position quickly while changing its preferred period more slowly. Comparing phase and tempo perturbations therefore tested a coupled-oscillator hypothesis: the sound sequence influences a repeating movement process, or **oscillator**. Transient behavior measures how ongoing coordination changes after an unexpected timing disturbance.




Rouse et al. (2016), Fig. 3. Responses to mid-trial timing changes.

The perturbation stimulus uses brief paired tones

The stimulus clicks combined tones at 659 and 1319 Hz for 10 ms. Repeating these brief events created a clear temporal sequence. The main baseline rate was 85 bpm, a rate Ronan had not previously encountered in training or testing.

Perturbations changed a timing interval by 3, 8, 15, or 25 percent in either direction. The change occurred between the sixteenth and twenty-fifth beat, with its position varied across conditions so that a fixed point in the sequence would not predict it.

At the baseline rate, the normal interval was about 706 ms. Controlled changes to that interval introduced experimentally specified timing errors into a sequence that Ronan was already following, allowing the subsequent head bobs to reveal how she recovered coordination.




Rouse et al. (2016), Fig. 1. Phase distributions before and after timing changes.

A phase shift moves the beat without a new tempo

In a phase-shift trial, a sound occurs earlier or later than expected, but later sounds return to the original spacing. The stimulus's temporal position changes while its continuing tempo remains fixed, separating an onset-position error from a lasting change in speed.

Ronan adjusted her head-bob timing after both advances and delays. Her movement regained an organized relationship with the shifted sequence within a few beats, rather than staying locked to the timing established before the perturbation.

This recovery is **phase correction**, the adjustment of an ongoing cycle's position relative to an external event. Because the subsequent sound interval returns to baseline, successful recovery can be achieved by a transient movement adjustment without adopting a permanently different repetition rate.



Rouse et al. (2016), Fig. 1, phase conditions. Timing distributions after phase advances and delays.




Rouse et al. (2016), Supplementary Video 1. Ronan adjusts to phase and tempo changes.

A tempo shift changes every later sound interval

A tempo perturbation alters the spacing of every subsequent beat. If the head-bob rhythm kept its original period, the relative position of movement and sound would drift increasingly across the trial, even after the first timing error had been corrected.

Ronan adapted after all the tested tempo changes. The response had to address the continuing mismatch in cycle duration, not merely the placement of a single unexpectedly early or late sound within an otherwise unchanged sequence.

The authors' sign convention referred to interval length: a positive interval change made the beat slower, while a negative change made it faster. Keeping interval and tempo distinct is necessary for interpreting which perturbations required Ronan to slow her repeated movement and which required her to accelerate it.




Rouse et al. (2016), Fig. 1, tempo conditions. Timing distributions after persistent rate changes.

The first transfer response already tracks the change

The first presentation of each perturbation provided a test before repeated experience with that exact change. Ronan retained organized movement relative to the beat after these first encounters, including phase changes and persistent changes in the stimulus interval.

The response therefore transferred beyond a repeatedly practiced correction at one fixed perturbation. The animal applied its existing auditory–motor coordination to an unfamiliar mid-trial disturbance, with later sounds continuing to influence the repeating behavioral cycle.

Initial-transfer data and repeated-trial data served different purposes. The former established that correction could appear at first exposure; the latter characterized the consistency and temporal course of correction across presentations, supporting a more detailed model of the behavior’s adjustment dynamics.




Rouse et al. (2016), Fig. 1. First presentations of timing-perturbation conditions.

Phase recovery occurs over successive movements

The researchers aligned trials at the perturbation and compared the relative phase of successive head bobs. Before the change, movement and sound were already coordinated. The sudden phase disturbance created a measurable error that could then be followed over later cycles.

Ronan corrected the disturbance within a few beats. Recovery was a trajectory across successive movements rather than an instantaneous switch from one static state to another. This time course supplied information about the strength and direction of adjustment.

The published model predicted these transient trajectories using coupling between the sound sequence and the behavioral oscillator. **Coupling** means that the timing of one process changes the state of the other; here, the sound events adjusted the timing of the continuing head-bob cycle.




Rouse et al. (2016), Fig. 3. Beat-by-beat recovery after phase changes.

Rate changes produce a lasting timing adjustment

Following a lasting rate change, Ronan's initial movement period and the new stimulus period differed. Head movements therefore had to adjust across several cycles to approach a stable relationship with the new continuing sound sequence.

The measured trajectories recovered after changes in either direction. A faster stimulus required more rapid movement, whereas a slower stimulus required longer intervals between selected movement landmarks. The correction preserved a regular relationship between sound and action despite the disturbance.

The period component of the coupled-oscillator model represented adjustment to this continuing rate mismatch. Combining period adjustment with phase adjustment improved the fit to the measured recovery trajectories, connecting two distinct timing operations to the transient response produced by the animal.




Rouse et al. (2016), Fig. 3. Recovery trajectories following rate shifts.

A double bob exposes an immediate motor adjustment

During some large slowing perturbations, the next sound occurred when Ronan's head was near the high point of a movement rather than its low point. She sometimes made an additional small nod before slowing the broader movement cycle.

The authors called these responses **double bobs** because two identifiable movements occurred around a beat. Their observation connected the timing disturbance with a specific change in movement form, rather than treating all behavior as identical cycles with only altered timestamps.

These unusual nods were excluded from the primary cycle-timing analysis and documented in Supplementary Video 1. Their occurrence is still informative behavioral material: an unexpected sound can modify both the timing of a repeated action and the shape of the immediate motor response.




Rouse et al. (2016), Fig. 3. Behavioral timing after persistent interval changes.

Timing concentration and timing offset are separate

A series of movements can be consistently early, consistently late, or widely scattered around the beat. The average phase describes the center of that relationship, whereas the concentration of phases describes how consistently successive movements occupy that position.

Ronan's post-perturbation phases were tightly concentrated across the tested conditions. Some conditions nevertheless retained systematic offsets. High consistency therefore describes reliable coordination with the sequence and does not automatically mean that the chosen movement landmark coincides exactly with sound onset.

The researchers evaluated these properties separately because different aspects of the timing process influence them. Correcting a disturbance may stabilize a relationship before eliminating its offset, and a repeating behavioral process can remain strongly coordinated while occupying a stable position ahead of or behind the sound.




Rouse et al. (2016), Fig. 2B. Consistency of beat-relative movement timing.

The stable phase varies with the new tempo

Phase changes and tempo changes produced different patterns after recovery. Under phase perturbations, the long-term stimulus rate returned to its original value. Under tempo perturbations, the ongoing sound interval remained altered, creating a new continuing rate relationship.

Ronan's average post-perturbation phase varied systematically with the altered tempo. Slower and faster conditions therefore retained distinct timing positions even when the head movements had regained a consistent relationship with the continuing sequence.

This phase–tempo relationship matched a pattern seen in the earlier transfer experiments. The repeated finding linked stable coordination across unfamiliar tempos with the transient correction of a sudden rate change, giving a common behavioral constraint for the oscillator model to explain.




Rouse et al. (2016), Fig. 2A. Average offsets after phase and tempo perturbations.

An oscillator is a repeating timing process

An **oscillator** is a process that repeatedly passes through a cycle. The model represented Ronan's head-bob timing as an oscillator influenced by the recurring auditory events. Its cycle position and repetition period could both change as sounds arrived.

The hypothesis was that coupled timing processes could account for the observed coordination. A sound arriving at an unexpected cycle position generated a corrective influence, while continuing differences between movement and sound periods produced further adjustment across subsequent cycles.

The published model closely matched the measured recovery trajectories for phase and tempo disturbances. This agreement supplied a mechanistic description of timing behavior, while the experiment itself measured head movement and sound timing rather than recording the neurons that might implement those coupled processes.




Rouse et al. (2016), Fig. 3. Measured recovery and coupled-oscillator predictions.

Strong phase coupling restores the timing position

The model's **phase-coupling parameter** described how strongly a sound event adjusted the position of the behavioral cycle. Stronger coupling produced a larger correction for a given timing error, allowing the movement to regain a stable relationship over fewer cycles.

Across the tested perturbations, phase adjustment was much stronger than period adjustment in the fitted model. Ronan's behavior could therefore be described primarily by rapid correction of cycle position, supplemented by weaker changes in the underlying repetition period.

The strength of phase coupling also increased with the magnitude of the disturbance. Larger timing errors elicited stronger corrections, connecting the severity of the unexpected auditory change with the modeled adjustment rather than assigning a single identical correction to every perturbation.




Rouse et al. (2016), Fig. 4A–C. Fitted phase and period adjustment parameters.

Period coupling adjusts the ongoing cycle duration

The **period** is the duration of one complete movement cycle. Changing the period alters the repetition rate itself. This differs from briefly shifting the cycle's position while leaving the preferred duration of later cycles unchanged.

In the fitted model, period coupling was weaker than phase coupling but improved the account of the observed behavior. Lasting rate changes required ongoing adjustment because correcting the first displaced event alone would leave later movements drifting relative to the sound.

The combination offered a behavioral mechanism with two time scales: relatively strong immediate phase correction and weaker period adaptation. These were model parameters inferred from movement trajectories, not measured concentrations of a transmitter or direct observations of separate neural populations in the sea lion.




Rouse et al. (2016), Fig. 3. Modeled and measured adaptation to new rates.

The same correction transfers to other base rates

The main perturbation series used 85 bpm. Researchers also tested two additional unfamiliar baseline rates, approximately 77.3 and 94.4 bpm, with phase and tempo disturbances. This extended the correction test beyond a single preferred starting rhythm.

Ronan adapted across the additional baseline conditions as well. The ability to correct an ongoing response therefore generalized both across disturbance types and across the initial rate from which the disturbance was introduced.

A timing mechanism that operates across these conditions must use the relationship between the current sound interval and the current movement cycle. A correction learned only for a particular sound timestamp or one fixed movement rate would not provide the same flexibility across different baseline rhythms.




Rouse et al. (2016), Fig. 1. Controlled timing perturbations around a baseline rhythm.

Hama matches human actions beyond specific commands

Hama, a Steller sea lion, had already learned a repertoire of actions through hand signals and verbal commands. Sasaki and colleagues trained a **do-as-I-do** response, in which a human action becomes the model for the animal's action. Demonstrating "stand," a head movement, or a tongue action replaced the usual specific cue.

The human initially paired demonstrations with familiar cues, then reduced those prompts. Hama subsequently matched most tested body actions with both a familiar trainer and an unfamiliar demonstrator. Several actions were familiar to her but had not been taught through the demonstration procedure.

Transfer separated general action matching from simply repeating the three demonstrations used during initial training. Performance also depended on what kind of action was attempted: novel body-oriented responses were easier than novel object manipulations. The acquired relationship linked observed body movement with selection from Hama's own action repertoire.



Sasaki et al. (2025), Fig. 2. Action matching with familiar and unfamiliar human demonstrators.

A Go cue separates remembering from immediate copying

In a second procedure, Hama watched a complete human demonstration while remaining still, then acted upon the verbal cue “Go.” **Response inhibition** is withholding an action that would otherwise occur. Learning to wait changed her earlier history of moving simultaneously with human demonstrations.

The tests included standing, spinning, lifting a hindflipper, and a kissing action. In a control condition, a barrier blocked visual contact with the demonstrator after the demonstration. Hama still produced the corresponding action when the demonstrator delivered the Go cue.

The final cue therefore specified when to respond, while the preceding demonstration specified what to do. Blocking the demonstrator during the response removed ongoing visual guidance as a requirement. This delayed command procedure differs from beat synchronization, where repeated sound events continually regulate the timing of an ongoing movement.




Sasaki et al. (2025), Fig. 5. Visual barriers used after the human demonstration.

Missing beats distinguish prediction from isolated reactions

A reaction to each sound separately need not maintain a rhythm through silence. Cook and colleagues therefore omitted occasional beats from a 120 bpm sequence. Discrete clicks replaced the continuously modulated sound so that the missing event left a genuinely silent interval.

Ronan continued bobbing through the omitted events. Movements before and after the gap retained similar relationships to the expected beat locations. The repeating action continued while the next timing sound was absent instead of restarting only when another click arrived.

Continuation through an omission supports an ongoing timing process beyond isolated sound-triggered reactions. Later sounds can correct the position of the movement within the cycle. This combination of continuation and correction connects predictive timing with the perturbation experiments that independently changed beat position and tempo.




Cook et al. (2013), Fig. 3A–B. Timing before and after omitted clicks.

Sound onsets and movement minima share a time base

A continuous head movement needs a defined event for comparison with a sound onset. Ronan's selected landmark was the lowest point of each head bob. In the human comparison, it was the fullest downward extension of the arm. Using a repeatable landmark keeps the measured timing event consistent.

The interval between successive movement minima supplies the performed tempo. The position of each minimum relative to a sound onset supplies phase. Automated video tracking measured the trajectories while the audio track from the same recording identified sound events.

The 2025 experiment recorded at 240 frames per second, corresponding to roughly 4 ms between frames. This resolution allowed small anticipatory and delayed offsets to be measured within a cycle. A movement rate near the target can coexist with a nonzero offset from each sound, so both quantities characterize the coordination.




Cook et al. (2025), Fig. 1. Movement trajectories aligned with recorded sound onsets.

Precise timing can include an anticipatory offset

A **negative asynchrony** occurs when the selected movement landmark precedes the sound. In Ronan's 2025 snare tests, the average phase at 112 and 120 bpm was negative, while the average at 128 bpm was slightly positive.

This timing pattern is inconsistent with treating every head bob as a delayed reaction initiated only after each sound occurs. A repeating movement can be scheduled in anticipation of the next event while continuing to receive timing information from the current acoustic sequence.

At the same time, anticipation need not place every movement exactly at zero phase. Coordination can be stable and predictive with a systematic offset. Describing the signed relationship preserves that biological timing pattern instead of reducing the response to a simple success-versus-failure label.




Cook et al. (2025), Fig. 3, 120 bpm row. Timing with the familiar-rate snare sequence.

Rate accuracy and response consistency differ

The human comparison separately evaluated performed tempo, variability of movement intervals, and timing relative to sound onset. These dimensions answer different behavioral questions: how fast the repeated action runs, how consistently it repeats, and where it falls within the external cycle.

Ronan’s intervals were more consistent than those of the human participants across the three test rates. Her average tempo or average phase was also closer to the target than that of many participants, but the relative ranking depended on the performance measure.

No human exceeded Ronan on all measured dimensions. The conclusion rests on matched-task performance rather than a claim that the animal is superior at every possible rhythmic activity. Gross movement, stimulus form, and the selected timing landmark define the behavior that was compared.




Cook et al. (2025), Fig. 3, 128 bpm row. Different dimensions of synchronized performance.

Human gross movements vary across individuals

Humans in the comparison were non-musicians performing a large arm movement rather than a familiar tapping task. Their synchronization differed among individuals, with some participants producing less consistent movement intervals and more variable phases than others.

The distribution of performance matters because a comparison with only a hypothetical perfectly regular human would conceal the range of actual behavior. The experiment measured the same sound sequences in both species and retained each participant's timing profile.

The study found no consistent rapid improvement across the brief human testing sequence. Ronan, in contrast, entered with a long history of the head-bob task. The observed comparison therefore describes an experienced sea lion and humans performing the specified large-amplitude coordination task, with their different task histories made explicit.




Cook et al. (2025), Fig. 3. Individual human and sea-lion timing measures.

Long-term performance includes improved phase alignment

At the familiar simple-stimulus rates, Ronan's later phase alignment was much closer to zero than in several early conditions. The large negative offsets recorded at slower rates in 2013 were substantially reduced when those stimuli were retested after longitudinal experience.

Improvement involved more than bringing the average movement rate near the stimulus. The timing landmark moved closer to the sound onset, and successive movements became more consistent. Rate, phase, and consistency therefore changed together in the later performance profile.

The 108 bpm condition already had a phase near zero in the early experiment, leaving less room for improvement on that particular measure. This condition-specific pattern connects the long-term result to the actual starting performance rather than assuming identical improvement at every tested rate.




Cook et al. (2025), Fig. 2. Reduced timing offsets in experienced performance.

The tested skill is flexible beat synchronization

Ronan’s experiments established coordination with regular pulse sequences and with musical recordings that contain a regular pulse. **Beat synchronization** is the alignment of a repeated action with those pulse events, with flexibility across rates and acoustic forms.

More complex musical abilities involve additional structure, such as grouping beats into recurring patterns or coordinating with deliberately irregular sequences. The 2025 paper distinguished the measured beat-keeping skill from these broader questions about musical organization.

That distinction changes the scientific interpretation: flexible auditory–motor timing is experimentally supported in this sea lion, while complete human-like musical competence was not the tested outcome. The behavioral result still provides a substantial comparative example of prediction, repeated movement, transfer, and adjustment to external rhythm.




Cook et al. (2025), Fig. 1. Shared movement and timing definitions.

Rats move near musical beats without task training


Ito and colleagues attached a lightweight wireless accelerometer to rats and measured small head movements while music played. An **accelerometer** measures acceleration along specified axes, allowing movement changes to be quantified without requiring a trained beat-marking action.

The rats listened to Mozart's Sonata for Two Pianos, K.448, at its original rate and at altered playback speeds. Their movements were more strongly aligned with the musical beats near the original tempo, approximately 132 bpm, than at the very fast versions.

This preparation differed from Ronan's explicitly trained head-bob task. It measured spontaneous small movements during listening rather than an operantly reinforced criterion response. The contrast provides two experimental routes to auditory–motor relationships: acquiring a reliable instructed action and measuring naturally occurring timing during sound exposure.



Ito et al. (2022), Fig. 1A–G. Accelerometer measurements during musical playback.




Ito et al. (2022), Movie S1. Rat head movements and simultaneous jerk traces.

Jerk detects rapid changes in head movement


Jerk is the rate of change of acceleration. The rat study used this quantity to identify brief movement changes during musical listening. A small movement with a rapid change in acceleration can therefore contribute strongly to the measured response.

Researchers compared jerk near putative beats with jerk elsewhere in the beat cycle. This beat contrast quantified whether the movement signal was organized around the musical pulse instead of being uniformly distributed through time.

At the original musical tempo, the rats showed beat-related movement contrast. Faster playback weakened that organization. The measured response therefore depended on the sound's temporal scale, connecting the music rate to the timing of small spontaneous movements rather than only to the overall amount of activity.



Ito et al. (2022), Fig. 1C–F. Acceleration, jerk, and beat-relative movement contrast.




Ito et al. (2022), Movie S2. Beat-related movements in a standing rat.

Rat and human movements favor similar tempos

The experiment also measured head movements in humans listening to the same musical excerpt at altered speeds. This provided a comparison in which both species were exposed to the same changing temporal structure, although their absolute movement magnitudes differed.

Beat-related movement organization was strongest near the original musical rate rather than at the greatly accelerated versions. The two species also showed related changes in movement strength across individual beats of the original excerpt.

Inspection of the stimulus and motion timing indicated that many movements occurred predictively near the upcoming beat. In this preparation, beat-related organization emerged during listening without Ronan's explicit synchronization training, broadening the experimental evidence beyond a single kind of instructed motor task.



Ito et al. (2022), Fig. 2A–C. Beat-related movement comparisons across playback speeds.




Ito et al. (2022), Movie S3. Human head movements and marker tracking.

Population activity is tuned to musical event timing

The rat study separately recorded **multiunit activity**, the combined spiking response of several nearby neurons, while presenting music at different playback speeds. Activity following beat events was compared with activity following other events in the same musical sequence.

Neural beat contrast was greatest around the original tempo and weakened at much faster speeds. The response also differed among positions in the repeating beat pattern, connecting neural activity to the temporal organization of the musical excerpt rather than merely its presence.

These recordings and the movement measurements were separate experimental assays. Their shared tempo preference linked a candidate neural timing process with the behavioral preference, while the authors used further click stimuli and a model to examine how adaptation could generate that pattern of sensitivity.




Ito et al. (2022), Fig. 3A–E. Neural responses to musical beats at different rates.

Short-term adaptation favors intermediate rhythms

Short-term adaptation is a change in responsiveness during recent stimulation. Repeated sounds can reduce a neural population's response, with recovery depending on the time since preceding stimulation. This creates a temporal dependence even when each sound is acoustically similar.

Ito and colleagues tested click sequences with three clicks and a rest at 60, 120, 240, and 480 bpm. Recorded responses distinguished beat positions most strongly near an intermediate rate. Very slow sequences and very rapid sequences produced different balances of adaptation and recovery.

The authors' adaptation model reproduced the intermediate-rate preference and related it to the timing structure of musical material. This is a hypothesis about a candidate neural computation; the sea-lion experiments independently established flexible movement correction, rather than directly measuring that rat adaptation process in Ronan.




Ito et al. (2022), Fig. 4A–I. Recorded responses and adaptation-model predictions.

A new sound separates timbre from tempo

Timbre is the acoustic quality that distinguishes sounds with similar timing. Cook and colleagues tested Ronan with a snare-drum sound that differed from the frequency-modulated training stimulus. A learned temporal relationship could therefore be expressed with a new acoustic form.

Before testing, Ronan encountered the snare at 96 bpm during ordinary training. Human participants and Ronan then received matching warm-up presentations, first with the familiar oscillating sound superimposed and then with the snare alone. This preparation reduced disruption by the sound's novelty without practicing the later unfamiliar rates.

The test sequences contained 25 equally spaced snare beats at 112, 120, or 128 bpm. The familiar 120 bpm condition and unfamiliar neighboring rates separated recognition of a practiced speed from flexible adjustment of the ongoing movement to an external pulse.




Cook et al. (2025), Fig. 1. Matched head and arm movement measurements in the UCSC experiment.

The UCSC comparison matches the movement task

The 2025 UCSC study compared Ronan's head bob with a downward arm movement in humans. Both tasks marked the beat with the lowest point of a relatively large movement. Familiar finger tapping would have changed the movement demands as well as the species being compared.

All subjects heard the same ordered sets of snare stimuli. Sound and movement were captured together, and automated video tracking supplied the movement trajectory. The time between consecutive movement minima measured performed tempo; the position of each minimum relative to a sound measured phase.

Ronan's conditioned reinforcer marked the final response and was followed by fish. Humans received a tone marking the trial's end. Reinforcement maintained the trained animal's participation while the matched stimuli and movement landmarks made the temporal measurements comparable across the two preparations.




Cook et al. (2025), Fig. 1. Comparable large-amplitude movements timed to sound onset.

A practiced rhythm supports nearby unfamiliar rates

Ronan’s average performed tempos were 113.1, 121.6, and 129.0 bpm for stimuli at 112, 120, and 128 bpm, respectively. The movement rate therefore changed with the sound sequence, including the two tempos she had not practiced. The behavior extended beyond reproducing the familiar 120 bpm action.

Her average movement phase preceded the sound at 112 and 120 bpm and followed it slightly at 128 bpm. A stable anticipatory movement can preserve an offset from sound onset while remaining coordinated. Rate accuracy and the signed placement of the movement within a cycle describe different features of that coordination.

Her movement intervals were more consistent than those of the human participants at every tested rate. Average rate, average phase, and consistency produced different individual rankings. The experimental result concerns this matched gross-movement task, with Ronan’s extensive practice contributing to the performance being measured.




Cook et al. (2025), Fig. 3. Measured sea-lion and human timing across the three test rates.

Longitudinal practice changes precision within the cycle

Cook and colleagues compared Ronan's early performance at age 3 with later performance at age 15. The same individual and familiar simple sound sequences allowed changes in practiced timing to be examined across a long training history. This is longitudinal evidence rather than a comparison between unrelated animals.

Later performance combined a movement rate near the target with much smaller offsets between movement minima and sound onsets. Slow stimuli that originally elicited markedly anticipatory movements later produced timing closer to the beat. Precision improved within the repeating cycle as well as in the average repetition rate.

The earlier 108 bpm condition already had an offset near zero, so improvement depended on the starting condition. Motor learning can refine different components of an acquired action by different amounts. In Ronan, sustained experience refined an auditory–motor skill that transferred to new acoustic forms and neighboring rates.



Cook et al. (2025), Fig. 2. Early and experienced timing profiles in the same sea lion.

Key takeaways

Sensory organization Facial information enters trigeminal brainstem relays, a large medial thalamic subdivision, and a layer-4-rich somatosensory cortical region.

Active control Sea lions adjust whisker movements to the discrimination task, and reposition their flippers during turns and leaps.

Learned cues A command can initiate exploration or specify when to act; Hama retained a demonstrated action across a delayed Go cue and visual occlusion.

Tempo and phase A matching repetition rate and a stable relationship to individual beats are distinct aspects of synchronization.

Predictive correction Ronan continued through omitted beats and corrected ongoing movement after changes in beat position or tempo.

Experience and comparison An experienced sea lion performed precisely in a matched gross-movement task; rat recordings identify a separate candidate adaptation process.

Sawyer et al. (2016); Milne et al. (2021); Sasaki et al. (2025); Cook et al. (2013, 2025); Ito et al. (2022)