

Ctenophora

Ctenophora (/tɪˈnɒfərə/; singular ctenophore, /ˈtɛnəfɔːr/ or /ˈtiːnəfɔːr/; from the Greek κτείς kteis 'comb' and φέρω pherō 'carry'; commonly known as comb jellies) is a phylum of animals that live in marine waters worldwide. Their most distinctive feature is the ‘combs’ – groups of cilia which they use for swimming – they are the largest animals that swim by means of cilia. Adults of various species range from a few millimeters to 1.5 m (4 ft 11 in) in size. Like cnidarians, their bodies consist of a mass of jelly, with one layer of cells on the outside and another lining the internal cavity. In ctenophores, these layers are two cells deep, while those in cnidarians are only one cell deep. Some authors combined ctenophores and cnidarians in one phylum, Coelenterata, as both groups rely on water flow through the body cavity for both digestion and respiration. Increasing awareness of the differences persuaded more recent authors to classify them as separate phyla.

Almost all ctenophores are predators, taking prey ranging from microscopic larvae and rotifers to the adults of small crustaceans; the exceptions are juveniles of two species, which live as parasites on the salps on which adults of their species feed. In favorable circumstances, ctenophores can eat ten times their own weight in a day. Only 100–150 species have been validated, and possibly another 25 have not been fully described and named. The textbook examples are cydippids with egg-shaped bodies and a pair of retractable tentacles fringed with tentilla ("little tentacles") that are covered with colloblasts, sticky cells that capture prey. The phylum has a wide range of body forms, including the flattened, deep-sea platyctenids, in which the adults of most species lack combs, and the coastal beroids, which lack tentacles and prey on other ctenophores by using huge mouths armed with groups of large, stiffened cilia that act as teeth. These variations enable different species to build huge populations in the same area, because they specialize in different types of prey, which they capture by as wide a range of methods as spiders use.

Most species are hermaphrodites—a single animal can produce both eggs and sperm, meaning it can fertilize its own egg, not needing a mate. Some are simultaneous hermaphrodites, which can produce both eggs and sperm at the same time. Others are sequential hermaphrodites, in which the eggs and sperm mature at different times. Fertilization is generally external, although platyctenids' eggs are fertilized inside their parents' bodies and kept there until they hatch. The young are generally planktonic and in most species look like miniature cydippids, gradually changing into their adult shapes as they grow. The exceptions are the beroids, whose young are miniature beroids with large mouths and no tentacles, and the platyctenids, whose young live as cydippid-like plankton until they reach near-adult size, but then sink to the bottom and rapidly metamorphose into the adult form. In at least some species, juveniles are capable of reproduction before reaching the adult size and shape. The combination of hermaphroditism and early reproduction enables small populations to grow at an explosive rate.

Ctenophores may be abundant during the summer months in some coastal locations, but in other places they are uncommon and difficult to find. In bays where they occur in very high numbers, predation by ctenophores may control the populations of small zooplanktonic organisms such as copepods, which might otherwise wipe out the phytoplankton (planktonic plants), which are a vital part of marine food chains. One ctenophore, *Mnemiopsis*, has accidentally been introduced into the Black Sea, where it is blamed for causing fish stocks to collapse by eating both fish larvae and organisms that would otherwise have fed the fish. The situation was aggravated by other factors, such as over-fishing and long-term environmental changes that promoted the growth of the *Mnemiopsis* population. The later accidental introduction of *Beroe* helped to mitigate the problem, as *Beroe* preys on other ctenophores.

Despite their soft, gelatinous bodies, fossils thought to represent ctenophores, apparently with no tentacles but many more comb-rows than modern forms, have been found in lagerstätten as far back as the early Cambrian, about 515 million years ago. The position of the ctenophores in the evolutionary family tree of animals has long been debated, and the majority view at present, based on molecular phylogenetics, is that cnidarians and bilaterians are more closely related to each other than either is to ctenophores. A recent molecular phylogenetics analysis concluded that the common ancestor of all modern ctenophores was

cydippid-like, and that all the modern groups appeared relatively recently, probably after the Cretaceous–Paleogene extinction event 66 million years ago. Evidence accumulating since the 1980s indicates that the "cydippids" are not monophyletic, in other words do not include all and only the descendants of a single common ancestor, because all the other traditional ctenophore groups are descendants of various cydippids.

Ctenophores form an animal phylum that is more complex than sponges, about as complex as cnidarians (jellyfish, sea anemones, etc.), and less complex than bilaterians (which include almost all other animals). Unlike sponges, both ctenophores and cnidarians have: cells bound by inter-cell connections and carpet-like basement membranes; muscles; nervous systems; and some have sensory organs. Ctenophores are distinguished from all other animals by having colloblasts, which are sticky and adhere to prey, although a few ctenophore species lack them.

Like sponges and cnidarians, ctenophores have two main layers of cells that sandwich a middle layer of jelly-like material, which is called the mesoglea in cnidarians and ctenophores; more complex animals have three main cell layers and no intermediate jelly-like layer. Hence ctenophores and cnidarians have traditionally been labelled diploblastic, along with sponges. Both ctenophores and cnidarians have a type of muscle that, in more complex animals, arises from the middle cell layer, and as a result some recent text books classify ctenophores as triploblastic, while others still regard them as diploblastic.

Ranging from about 1 millimeter (0.039 in) to 1.5 meters (4.9 ft) in size, ctenophores are the largest non-colonial animals that use cilia ("hairs") as their main method of locomotion. Most species have eight strips, called comb rows, that run the length of their bodies and bear comb-like bands of cilia, called "ctenes," stacked along the comb rows so that when the cilia beat, those of each comb touch the comb below. The name "ctenophora" means "comb-bearing", from the Greek κτείς (stem-form κτεν-) meaning "comb" and the Greek suffix -φορος meaning "carrying".

For a phylum with relatively few species, ctenophores have a wide range of body plans. Coastal species need to be tough enough to withstand waves and swirling sediment particles, while some oceanic species are so fragile that it is very difficult to capture them intact for study. In addition oceanic species do not preserve well, and are known mainly from photographs and from observers' notes. Hence most attention has until recently concentrated on three coastal genera – Pleurobrachia, Beroë and Mnemiopsis. At least two textbooks base their descriptions of ctenophores on the cydippid Pleurobrachia.

The internal cavity forms: a mouth that can usually be closed by muscles; a pharynx ("throat"); a wider area in the center that acts as a stomach; and a system of internal canals. These branch through the mesoglea to the most active parts of the animal: the mouth and pharynx; the roots of the tentacles, if present; all along the underside of each comb row; and four branches round the sensory complex at the far end from the mouth – two of these four branches terminate in anal pores. The inner surface of the cavity is lined with an epithelium, the gastrodermis. The mouth and pharynx have both cilia and well-developed muscles. In other parts of the canal system, the gastrodermis is different on the sides nearest to and furthest from the organ that it supplies. The nearer side is composed of tall nutritive cells that store nutrients in vacuoles (internal compartments), germ cells that produce eggs or sperm, and photocytes that produce bioluminescence. The side furthest from the organ is covered with ciliated cells that circulate water through the canals, punctuated by ciliary rosettes, pores that are surrounded by double whorls of cilia and connect to the mesoglea.

The outer surface bears usually eight comb rows, called swimming-plates, which are used for swimming. The rows are oriented to run from near the mouth (the "oral pole") to the opposite end (the "aboral pole"), and are spaced more or less evenly around the body, although spacing patterns vary by species and in most species the comb rows extend only part of the distance from the aboral pole towards the mouth. The "combs" (also called "ctenes" or "comb plates") run across each row, and each consists of thousands of unusually long cilia, up to 2 millimeters (0.079 in). Unlike conventional cilia and flagella, which has a filament structure arranged in a 9 + 2 pattern, these cilia are arranged in a 9 + 3 pattern, where the extra compact filament is suspected to have a supporting function. These normally beat so that the propulsion

stroke is away from the mouth, although they can also reverse direction. Hence ctenophores usually swim in the direction in which the mouth is pointing, unlike jellyfish. When trying to escape predators, one species can accelerate to six times its normal speed; some other species reverse direction as part of their escape behavior, by reversing the power stroke of the comb plate cilia.

It is uncertain how ctenophores control their buoyancy, but experiments have shown that some species rely on osmotic pressure to adapt to water of different densities. Their body fluids are normally as concentrated as seawater. If they enter less dense brackish water, the ciliary rosettes in the body cavity may pump this into the mesoglea to increase its bulk and decrease its density, to avoid sinking. Conversely if they move from brackish to full-strength seawater, the rosettes may pump water out of the mesoglea to reduce its volume and increase its density.

The largest single sensory feature is the aboral organ (at the opposite end from the mouth). Its main component is a statocyst, a balance sensor consisting of a statolith, a solid particle supported on four bundles of cilia, called "balancers", that sense its orientation. The statocyst is protected by a transparent dome made of long, immobile cilia. A ctenophore does not automatically try to keep the statolith resting equally on all the balancers. Instead its response is determined by the animal's "mood", in other words the overall state of the nervous system. For example, if a ctenophore with trailing tentacles captures prey, it will often put some comb rows into reverse, spinning the mouth towards the prey.

Cydropid ctenophores have bodies that are more or less rounded, sometimes nearly spherical and other times more cylindrical or egg-shaped; the common coastal "sea gooseberry," *Pleurobrachia*, sometimes has an egg-shaped body with the mouth at the narrow end, although some individuals are more uniformly round. From opposite sides of the body extends a pair of long, slender tentacles, each housed in a sheath into which it can be withdrawn. Some species of cydropids have bodies that are flattened to various extents, so that they are wider in the plane of the tentacles.

The tentacles of cydropid ctenophores are typically fringed with tentilla ("little tentacles"), although a few genera have simple tentacles without these sidebranches. The tentacles and tentilla are densely covered with microscopic colloblasts that capture prey by sticking to it. Colloblasts are specialized mushroom-shaped cells in the outer layer of the epidermis, and have three main components: a domed head with vesicles (chambers) that contain adhesive; a stalk that anchors the cell in the lower layer of the epidermis or in the mesoglea; and a spiral thread that coils round the stalk and is attached to the head and to the root of the stalk. The function of the spiral thread is uncertain, but it may absorb stress when prey tries to escape, and thus prevent the colloblast from being torn apart. In addition to colloblasts, members of the genus *Haeckelia*, which feed mainly on jellyfish, incorporate their victims' stinging nematocytes into their own tentacles – some cnidaria-eating nudibranchs similarly incorporate nematocytes into their bodies for defense. The tentilla of *Euplokamis* differ significantly from those of other cydropids: they contain striated muscle, a cell type otherwise unknown in the phylum Ctenophora; and they are coiled when relaxed, while the tentilla of all other known ctenophores elongate when relaxed. *Euplokamis*' tentilla have three types of movement that are used in capturing prey: they may flick out very quickly (in 40 to 60 milliseconds); they can wriggle, which may lure prey by behaving like small planktonic worms; and they coil round prey. The unique flicking is an uncoiling movement powered by contraction of the striated muscle. The wriggling motion is produced by smooth muscles, but of a highly specialized type. Coiling around prey is accomplished largely by the return of the tentilla to their inactive state, but the coils may be tightened by smooth muscle.

There are eight rows of combs that run from near the mouth to the opposite end, and are spaced evenly round the body. The "combs" beat in a metachronal rhythm rather like that of a Mexican wave. From each balancer in the statocyst a ciliary groove runs out under the dome and then splits to connect with two adjacent comb rows, and in some species runs all the way along the comb rows. This forms a mechanical system for transmitting the beat rhythm from the combs to the balancers, via water disturbances created by the cilia.

The Lobata have a pair of lobes, which are muscular, cuplike extensions of the body that project beyond

the mouth. Their inconspicuous tentacles originate from the corners of the mouth, running in convoluted grooves and spreading out over the inner surface of the lobes (rather than trailing far behind, as in the Cydippida). Between the lobes on either side of the mouth, many species of lobates have four auricles, gelatinous projections edged with cilia that produce water currents that help direct microscopic prey toward the mouth. This combination of structures enables lobates to feed continuously on suspended planktonic prey.

Lobates have eight comb-rows, originating at the aboral pole and usually not extending beyond the body to the lobes; in species with (four) auricles, the cilia edging the auricles are extensions of cilia in four of the comb rows. Most lobates are quite passive when moving through the water, using the cilia on their comb rows for propulsion, although *Leucothea* has long and active auricles whose movements also contribute to propulsion. Members of the lobate genera *Bathocyroe* and *Ocyropsis* can escape from danger by clapping their lobes, so that the jet of expelled water drives them backwards very quickly. Unlike cydippids, the movements of lobates' combs are coordinated by nerves rather than by water disturbances created by the cilia, yet combs on the same row beat in the same Mexican wave style as the mechanically coordinated comb rows of cydippids and beroids. This may have enabled lobates to grow larger than cydippids and to have shapes that are less egg-like.

The Beroida, also known as Nuda, have no feeding appendages, but their large pharynx, just inside the large mouth and filling most of the saclike body, bears "macrocilia" at the oral end. These fused bundles of several thousand large cilia are able to "bite" off pieces of prey that are too large to swallow whole – almost always other ctenophores. In front of the field of macrocilia, on the mouth "lips" in some species of *Beroe*, is a pair of narrow strips of adhesive epithelial cells on the stomach wall that "zip" the mouth shut when the animal is not feeding, by forming intercellular connections with the opposite adhesive strip. This tight closure streamlines the front of the animal when it is pursuing prey.

The Cestida ("belt animals") are ribbon-shaped planktonic animals, with the mouth and aboral organ aligned in the middle of opposite edges of the ribbon. There is a pair of comb-rows along each aboral edge, and tentilla emerging from a groove all along the oral edge, which stream back across most of the wing-like body surface. Cestids can swim by undulating their bodies as well as by the beating of their comb-rows. There are two known species, with worldwide distribution in warm, and warm-temperate waters: *Cestum veneris* ("Venus' girdle") is among the largest ctenophores – up to 1.5 meters (4.9 ft) long, and can undulate slowly or quite rapidly. *Velamen parallelum*, which is typically less than 20 centimeters (0.66 ft) long, can move much faster in what has been described as a "darting motion".

Most Platyctenida have oval bodies that are flattened in the oral-aboral direction, with a pair of tentilla-bearing tentacles on the aboral surface. They cling to and creep on surfaces by everting the pharynx and using it as a muscular "foot". All but one of the known platyctenid species lack comb-rows. Platyctenids are usually cryptically colored, live on rocks, algae, or the body surfaces of other invertebrates, and are often revealed by their long tentacles with many sidebranches, seen streaming off the back of the ctenophore into the current.

Almost all species are hermaphrodites, in other words they function as both males and females at the same time – except that in two species of the genus *Ocyropsis* individuals remain of the same single sex all their lives. The gonads are located in the parts of the internal canal network under the comb rows, and eggs and sperm are released via pores in the epidermis. Fertilization is external in most species, but platyctenids use internal fertilization and keep the eggs in brood chambers until they hatch. Self-fertilization has occasionally been seen in species of the genus *Mnemiopsis*, and it is thought that most of the hermaphroditic species are self-fertile.

Development of the fertilized eggs is direct, in other words there is no distinctive larval form, and juveniles of all groups generally resemble miniature cydippid adults. In the genus *Beroe* the juveniles, like the adults, lack tentacles and tentacle sheaths. In most species the juveniles gradually develop the body forms of their parents. In some groups, such as the flat, bottom-dwelling platyctenids, the juveniles behave more like true larvae, as they live among the plankton and thus occupy a different ecological niche

from their parents and attain the adult form by a more radical metamorphosis, after dropping to the sea-floor.

When some species, including *Bathyctena chuni*, *Euplokamis stationis* and *Eurhamphaea vexilligera*, are disturbed, they produce secretions (ink) that luminesce at much the same wavelengths as their bodies. Juveniles will luminesce more brightly in relation to their body size than adults, whose luminescence is diffused over their bodies. Detailed statistical investigation has not suggested the function of ctenophores' bioluminescence nor produced any correlation between its exact color and any aspect of the animals' environments, such as depth or whether they live in coastal or mid-ocean waters.

Almost all ctenophores are predators – there are no vegetarians and only one genus that is partly parasitic. If food is plentiful, they can eat 10 times their own weight per day. While *Beroe* preys mainly on other ctenophores, other surface-water species prey on zooplankton (planktonic animals) ranging in size from the microscopic, including mollusc and fish larvae, to small adult crustaceans such as copepods, amphipods, and even krill. Members of the genus *Haeckelia* prey on jellyfish and incorporate their prey's nematocysts (stinging cells) into their own tentacles instead of colloblasts. Ctenophores have been compared to spiders in their wide range of techniques from capturing prey – some hang motionless in the water using their tentacles as "webs", some are ambush predators like Salticid jumping spiders, and some dangle a sticky droplet at the end of a fine thread, as bolas spiders do. This variety explains the wide range of body forms in a phylum with rather few species. The two-tentacled "cydippid" *Lampea* feeds exclusively on salps, close relatives of sea-squirts that form large chain-like floating colonies, and juveniles of *Lampea* attach themselves like parasites to salps that are too large for them to swallow. Members of the cydippid genus *Pleurobrachia* and the lobate *Bolinopsis* often reach high population densities at the same place and time because they specialize in different types of prey: *Pleurobrachia*'s long tentacles mainly capture relatively strong swimmers such as adult copepods, while *Bolinopsis* generally feeds on smaller, weaker swimmers such as rotifers and mollusc and crustacean larvae.

Ctenophores used to be regarded as "dead ends" in marine food chains because it was thought their low ratio of organic matter to salt and water made them a poor diet for other animals. It is also often difficult to identify the remains of ctenophores in the guts of possible predators, although the combs sometimes remain intact long enough to provide a clue. Detailed investigation of chum salmon, *Oncorhynchus keta*, showed that these fish digest ctenophores 20 times as fast as an equal weight of shrimps, and that ctenophores can provide a good diet if there are enough of them around. *Beroids* prey mainly on other ctenophores. Some jellyfish and turtles eat large quantities of ctenophores, and jellyfish may temporarily wipe out ctenophore populations. Since ctenophores and jellyfish often have large seasonal variations in population, most fish that prey on them are generalists, and may have a greater effect on populations than the specialist jelly-eaters. This is underlined by an observation of herbivorous fishes deliberately feeding on gelatinous zooplankton during blooms in the Red Sea. The larvae of some sea anemones are parasites on ctenophores, as are the larvae of some flatworms that parasitize fish when they reach adulthood.

On the other hand, in the late 1980s the Western Atlantic ctenophore *Mnemiopsis leidyi* was accidentally introduced into the Black Sea and Sea of Azov via the ballast tanks of ships, and has been blamed for causing sharp drops in fish catches by eating both fish larvae and small crustaceans that would otherwise feed the adult fish. *Mnemiopsis* is well equipped to invade new territories (although this was not predicted until after it so successfully colonized the Black Sea), as it can breed very rapidly and tolerate a wide range of water temperatures and salinities. The impact was increased by chronic overfishing, and by eutrophication that gave the entire ecosystem a short-term boost, causing the *Mnemiopsis* population to increase even faster than normal – and above all by the absence of efficient predators on these introduced ctenophores. *Mnemiopsis* populations in those areas were eventually brought under control by the accidental introduction of the *Mnemiopsis*-eating North American ctenophore *Beroe ovata*, and by a cooling of the local climate from 1991 to 1993, which significantly slowed the animal's metabolism. However the abundance of plankton in the area seems unlikely to be restored to pre-*Mnemiopsis* levels.

Because of their soft, gelatinous bodies, ctenophores are extremely rare as fossils, and fossils that have been interpreted as ctenophores have been found only in lagerstätten, places where the environment was

exceptionally suited to preservation of soft tissue. Until the mid-1990s only two specimens good enough for analysis were known, both members of the crown group, from the early Devonian (Emsian) period. Three additional putative species were then found in the Burgess Shale and other Canadian rocks of similar age, about 505 million years ago in the mid-Cambrian period. All three apparently lacked tentacles but had between 24 and 80 comb rows, far more than the 8 typical of living species. They also appear to have had internal organ-like structures unlike anything found in living ctenophores. One of the fossil species first reported in 1996 had a large mouth, apparently surrounded by a folded edge that may have been muscular. Evidence from China a year later suggests that such ctenophores were widespread in the Cambrian, but perhaps very different from modern species – for example one fossil's comb-rows were mounted on prominent vanes. The Ediacaran *Eoandromeda* could putatively represent a comb jelly.

The early Cambrian sessile frond-like fossil *Stromatoveris*, from China's Chengjiang lagerstätte and dated to about 515 million years ago, is very similar to *Vendobionta* of the preceding Ediacaran period. De-Gan Shu, Simon Conway Morris et al. found on its branches what they considered rows of cilia, used for filter feeding. They suggested that *Stromatoveris* was an evolutionary "aunt" of ctenophores, and that ctenophores originated from sessile animals whose descendants became swimmers and changed the cilia from a feeding mechanism to a propulsion system.

The relationship of ctenophores to the rest of Metazoa is very important to our understanding of the early evolution of animals and the origin of multicellularity. It has been the focus of debate for many years. Ctenophores have been purported to be the sister lineage to the Bilateria, sister to the Cnidaria, sister to Cnidaria, Placozoa and Bilateria, and sister to all other animal phyla. A series of studies that looked at the presence and absence of members of gene families and signalling pathways (e.g., homeoboxes, nuclear receptors, the Wnt signaling pathway, and sodium channels) showed evidence congruent with the latter two scenarios, that ctenophores are either sister to Cnidaria, Placozoa and Bilateria or sister to all other animal phyla. Several more recent studies comparing complete sequenced genomes of ctenophores with other sequenced animal genomes have also supported ctenophores as the sister lineage to all other animals. This position would suggest that neural and muscle cell types were either lost in major animal lineages (e.g., Porifera) or that they evolved independently in the ctenophore lineage. However, other researchers have argued that the placement of Ctenophora as sister to all other animals is a statistical anomaly caused by the high rate of evolution in ctenophore genomes, and that Porifera (sponges) is the earliest-diverging animal phylum instead. Ctenophores and sponges are also the only known animal phyla that lack any true *hox* genes.

Since all modern ctenophores except the beroids have cydippid-like larvae, it has widely been assumed that their last common ancestor also resembled cydippids, having an egg-shaped body and a pair of retractable tentacles. Richard Harbison's purely morphological analysis in 1985 concluded that the cydippids are not monophyletic, in other words do not contain all and only the descendants of a single common ancestor that was itself a cydippid. Instead he found that various cydippid families were more similar to members of other ctenophore orders than to other cydippids. He also suggested that the last common ancestor of modern ctenophores was either cydippid-like or beroid-like. A molecular phylogeny analysis in 2001, using 26 species, including 4 recently discovered ones, confirmed that the cydippids are not monophyletic and concluded that the last common ancestor of modern ctenophores was cydippid-like. It also found that the genetic differences between these species were very small – so small that the relationships between the Lobata, Cestida and Thalassocalycida remained uncertain. This suggests that the last common ancestor of modern ctenophores was relatively recent, and perhaps was lucky enough to survive the Cretaceous–Paleogene extinction event 65.5 million years ago while other lineages perished. When the analysis was broadened to include representatives of other phyla, it concluded that cnidarians are probably more closely related to bilaterians than either group is to ctenophores but that this diagnosis is uncertain.

What is a ctenophora

phylum of animals that live in marine waters

What does the ctenophora use to swim

‘combs’ – groups of cilia

What does ctenophora use for digestion and respiration

water flow through the body cavity

How large can ctenophora grow

1.5 m (4 ft 11 in)

What is the most distinctive feature of ctenophora

‘combs’ – groups of cilia

What are ctenophora commonly known as

comb jellies

How big can ctenophora grow

1.5 m (4 ft 11 in)

What does ctenophora rely on for digestion and respiration

water flow through the body cavity

What does ctenophora mean in Greek

κτείς kteis 'comb' and φέρω pherō 'carry'

Where do ctenophora live

marine waters

How many species of Ctenophora have been validated

100–150 species

What are the little tentacles that cydippids have called

tentilla

How much food does a ctenophora eat in a day

ten times their own weight

What do coastal beroids not have that other ctenophora have

tentacles

What does coastal beroids use as teeth

groups of large, stiffened cilia

How much do ctenophore eat in a day

ten times their own weight

What are the small tentacles on Cydippids called

tentilla

What do Beroids use as teeth

groups of large, stiffened cilia

What do Cydippids use to capture their prey

colloblasts

How many different species of ctenophore are there

100–150 species

What is unique about a hermaphrodite

can produce both eggs and sperm, meaning it can fertilize its own egg

What can a simultaneous hermaphrodite do

can produce both eggs and sperm at the same time

What type of hermaphrodite produces egg and sperm at different times

sequential

Which species eggs are fertilized and kept inside the parents body until hatched

platyctenids

What causes the population of ctenophora to grow at an explosive rate

hermaphroditism and early reproduction

What is a hermaphrodite

a single animal can produce both eggs and sperm

What is unique about simultaneous hermaphrodites

can produce both eggs and sperm at the same time.

What is a trait of sequential hermaphrodites

the eggs and sperm mature at different times

Which group keep the eggs are fertilized and kept inside the parent's body until they hatch

platyctenids

Which group has young that are born with no tentacles and a large mouth

beroids

What does the Beroe eat

other ctenophores

What was introduced into the Black Sea

Mnemiopsis

What does mnemiopsis eat

fish larvae and organisms

Where do ctenophores be found in large numbers

In bays

Where can ctenophores be found in large amounts

In bays

What are phytoplankton

planktonic plants

What ctenophore was accidentally introduced into The Black Sea

Mnemiopsis

What event was blamed on the introduction of mnemiopsis into The Black Sea

causing fish stocks to collapse

What was done to counteract the overpopulation of mnemiopsis in The Black Sea

introduction of Beroe

When did the Cretaceous-Paleogene extinction happen

66 million years ago

Evidence indicates that Cydippids are not what

monophyletic

How old are the fossils found that represent ctenophores

515 million years

What were the fossils that were found to represent ctenophores missing that current ctenophora have

tentacles

Fossils found that were believed to be ctenophores were how old

515 million years

What event happened 66 million years ago

Cretaceous–Paleogene extinction

Cypiddids are not what

monophyletic

What do current ctenophores have that fossils found did not have

tentacles

Jellyfish ans sea anemones belong to what phylum

cnidarians

What makes ctenophores different from all other animals

by having colloblasts

Ctenophora are less complex than which other phylum

bilaterians

Which phylum is more complex than sponges

Ctenophores

What does ctenophore use to capture prey

colloblasts

Jellyfish and sea anemones belong to which group/

cnidarians

What do ctenophores have that no other animals have

colloblasts

What do ctenophore use to capture their prey

colloblasts

Which two groups have cells bound by inter-cell connections and membranes, muscles, a nervous system and sensory organs

ctenophores and cnidarians

Ctenophores are less complex than what other group

bilaterians

What is the jelly-like substance called

mesoglea

Ctenophores and cnidarians are classified as what

diploblastic

Which group has two layers of cells with a middle layer of mesoglea

sponges and cnidarians, ctenophores

Ctenophores, cnidarians and what other group are labelled diploblastic

sponges

What are the hairs on ctenophores called

cilia

What are cilia used for

method of locomotion

Comb like bands of cilia are called what

ctenes

What does ctenophore mean in Greek

comb-bearing

Which cidippid is used as a description on ctenophores in most textbooks

Pleurobrachia

Which group of ctenophore are are hardest to study

oceanic species

Why are coastal species tough

to withstand waves and swirling sediment particles

Which ctenophora have been studies the most

Pleurobrachia, Beroe and Mnemiopsis

The inside of a ctenophore is lined with what

epithelium

What do photocytes produce

bioluminescence

What is the throat called

pharynx

What does the internal cavity contain

a mouth that can usually be closed by muscles; a pharynx ("throat"); a wider area in the center that acts as a stomach; and a system of internal canals.

What are the most active parts of ctenophora

the mouth and pharynx;

What are the eight comb rows on the outer surface called

swimming-plates

Combs are called what

also called "ctenes" or "comb plates

What does the 9 +3 pattern of cilia thought to do

supporting function

What direction do ctenophore swim

in the direction in which the mouth is pointing,

Cilia can grow up too what length

2 millimeters (0.079 in)

How do ctenophores control buoyancy

osmotic pressure

Ciliary rosettes pump water into what to control buoyancy

the mesoglea

What does pumping water into the mesoglea do

increase its bulk and decrease its density

What does the ciliary rosettes do to decrease bulk and increase density

pump water out of the mesoglea

What is the largest sensory feature of the ctenophora

aboral organ

Where is the aboral organ located

at the opposite end from the mouth

What protects the statocyst

a transparent dome made of long, immobile cilia

What is the main component of the aboral organ

a statocyst

What is a statocyst

a balance sensor

What is the common coastal pleurobrachia called

sea gooseberry

What does the pleurobrachia have on opposite sides of its body

a pair of long, slender tentacles

Cydippid are typically what shape

more or less rounded

The long tentacles on the pleurbrachia are protected by what

a sheath

Where is the mouth located on the pleuobrachia located

at the narrow end

What are the tentacles of cydipped ctenophores are usually fringed with

tentilla

What are colloblasts

specialized mushroom-shaped cells in the outer layer of the epidermis

What makes the tentilla of euplokamis different from other cysippids

they contain striated muscle,

How many types of movements do euplokamis tentilla have

three types of movement

What does the euplokamis use the three types of movement for

capturing prey

How many rows of combs are there

eight rows

Where are the rows of combs located

from near the mouth to the opposite end

How are the combs spaced

evenly round the body

What runs from the balancer in the statocyst to the comb rows

ciliary groove

What does the lobata have pair of

lobes

What are auricles

gelatinous projections edged with cilia that produce water currents

How many auricles do most species have

four

What do the auricles do

help direct microscopic prey toward the mouth

What do lobates feed on

suspended planktonic prey

What does the bathocyroe and ocyropsis do to escape danger

by clapping their lobes

What happens when bathocyroe and ocyropsis clap their lobes together

jet of expelled water drives them backwards very quickly.

The movements of the lobates combs are controlled by what

nerves

Cydippids combs are controlled by what

water disturbances created by the cilia

Beroida are known by what other name

Nuda

What group has no feeding appendages

The Beroida

Some species of beroe have a pair of strips of adhesive cells on the stomach wall. What does it do

zip" the mouth shut when the animal is not feeding,

What does the beroe do when pursuing prey

"zip" the mouth shut

What does the beroida have instead of feeding appendages

large pharynx

Which species are ribbon-shaped planktonic animals

The Cestida

What are cestida called

belt animals

How do cestids swim

by undulating their bodies as well as by the beating of their comb-rows.

What is the largest ctenophore

Cestum veneris

Which species moves by a darting motion

Velamen parallelum

What do most platyctenida have on their aboral surface

a pair of tentilla-bearing tentacles

What do platyctenida use their pharynx for

cling to and creep on surfaces

What do all but one platyctenida species lack

comb-rows

Where do platyctenida live

on rocks, algae, or the body surfaces of other invertebrates

How are eggs and sperm released

via pores in the epidermis

How do platyctenids reproduce

internal fertilization and keep the eggs in brood chambers until they hatch.

What genus has self-fertilization been seen

Mnemiopsis

Where are the gonads located

in the parts of the internal canal network under the comb rows

Is fertilization internal or external in most species

external

Juvenile and adult beroe lack what

tentacles and tentacle sheaths

Where do juvenile platyctenids live

among the plankton

When do juvenile develop into adults

after dropping to the sea-floor

Juvenile platyctenids act like what

more like true larvae

Which genus lack tentacles and sheaths

Beroe

What do bathyctena chuni, euplokamis stationis and eurhamphaea vexilligera have in common

they produce secretions (ink) that luminesce

When do bathyctena chuni, euplokamis stationis and eurhamphaea vexilligera excrete secretions

are disturbed,

What are the secretions commonly called

ink

Do adults or juveniles secretions luminesce brighter

Juveniles will luminesce more brightly

Are ctenophores predators, vegetarian or parasitic

Almost all ctenophores are predators

Haeckelia prey mostly on what animal

jellyfish

What happens to the jellyfish nematocysts when they are eaten by the haeckelia

incorporate their prey's nematocysts (stinging cells) into their own tentacles instead of colloblasts

What does the bolinopsis generally eat

smaller, weaker swimmers such as rotifers and mollusc and crustacean larvae.

What is the name of the two-tentacled cydippid that feeds entirely on salps called

Lampea

Why was it thought that ctenophores were a poor diet for other animals

their low ratio of organic matter to salt and water

What is the oncorhynchus also called

chum salmon

What do beroids typically eat

ctenophores

Where have herbivorous fishes been seen feeding on gelatinous zooplankton

the Red Sea

Turtles and jellyfish can eat large quantities of what

ctenophores,

What is blamed for causing the sharp drop in fish catches in the Black Sea and the Sea of Azov

ctenophore Mnemiopsis leidyi

How was the ctenophore mnemiopsis leidyi introduced into The Black Sea and the Sea of Azov

via the ballast tanks of ships

When was ctenophore mnemiopsis leidyi introduced into The Black Sea and the Sea of Azov

in the late 1980s

How was the population of mnemiopsis in The black Sea and the Sea of Azov brought under control

by the accidental introduction of the Mnemiopsis-eating North American ctenophore Beroe ovata,

How does cooling of the local environment affect the mnemiopsis

significantly slowed the animal's metabolism

Why are ctenophores extremely rare as fossils

Because of their soft, gelatinous bodies

Ediacaran eoandromeda can be regarded to represent what

comb jelly.

What was the period called that was 505 million years ago

Cambrian period.

How many species were found in the Burgess Shale

Three additional putative species

What did the fossils found in the Burgess Shale

lack

lacked tentacles

How old were the fossils found in China

515 million years

What type of fossils were found in China

Cambrian sessile frond-like fossil *Stromatoveris*

Which genus is considered the "aunt" of ctenophores

Stromatoveris

Stromatoveris is similar to which genus

Vendobionta

Vendobionta lived during which period

Ediacaran period

Recent studies believe that ctenophores are the sister lineage to what

all other animals

What do some researchers believe is the earliest-diverging animal phylum is

Porifera

Which genus of ctenophores does not have cydippid-like larvae

beroids

A molecular phylogeny analysis confirmed that cydippid are not what

monophyletic

When was the Cretaceous-Paleogen extinction occur

65.5 million years ago

Who did a morphologically analysis in 1985 that concluded cydippids are not monophyletic

Richard Harbison