


## RESEARCH ARTICLE

## Association of brain microbleeds with risk factors, cognition, and MRI markers in MESA

Paul N. Jensen<sup>1,2</sup> | Tanweer Rashid<sup>3</sup> | Jeffrey B. Ware<sup>4</sup> | Yuhua Cui<sup>5</sup> |  
 Colleen M. Sitlani<sup>1,2</sup> | Thomas R. Austin<sup>1,6</sup> | W. T. Longstreth Jr.<sup>6,7</sup> | Alain G. Bertoni<sup>8</sup> |  
 Elizabeth Mamourian<sup>4,5</sup> | R. Nick Bryan<sup>4</sup> | Ilya M. Nasrallah<sup>4,5</sup> | Mohamad Habes<sup>3,5</sup> |  
 Susan R. Heckbert<sup>1,6</sup> 

<sup>1</sup>Cardiovascular Health Research Unit,  
University of Washington, Seattle,  
Washington, USA

<sup>2</sup>Department of Medicine, University of  
Washington, Seattle, Washington, USA

<sup>3</sup>Neuroimage Analytics Laboratory and Biggs  
Institute Neuroimaging Core, Glenn Biggs  
Institute for Alzheimer's and  
Neurodegenerative Diseases, University of  
Texas Health Science Center San Antonio, San  
Antonio, Texas, USA

<sup>4</sup>Department of Radiology, Perelman School of  
Medicine, University of Pennsylvania,  
Philadelphia, Pennsylvania, USA

<sup>5</sup>Center for AI and Data Science for Integrated  
Diagnostics and Center for Biomedical Image  
Computing and Analytics, University of  
Pennsylvania, Philadelphia, Pennsylvania, USA

<sup>6</sup>Department of Epidemiology, University of  
Washington, Seattle, Washington, USA

<sup>7</sup>Department of Neurology, University of  
Washington, Seattle, Washington, USA

<sup>8</sup>Department of Internal Medicine, Wake  
Forest School of Medicine, Winston-Salem,  
North Carolina, USA

## Correspondence

Susan R. Heckbert, Department of  
Epidemiology, Cardiovascular Health Research  
Unit, University of Washington, 4333 Brooklyn  
Ave NE, Box 359458, Seattle, WA  
98195-9458, USA.  
Email: [heckbert@uw.edu](mailto:heckbert@uw.edu)

## Abstract

**INTRODUCTION:** Little is known about the epidemiology of brain microbleeds in racially/ethnically diverse populations.

**METHODS:** In the Multi-Ethnic Study of Atherosclerosis, brain microbleeds were identified from 3T magnetic resonance imaging susceptibility-weighted imaging sequences using deep learning models followed by radiologist review.

**RESULTS:** Among 1016 participants without prior stroke (25% Black, 15% Chinese, 19% Hispanic, 41% White, mean age 72), microbleed prevalence was 20% at age 60 to 64.9 and 45% at  $\geq 85$  years. Deep microbleeds were associated with older age, hypertension, higher body mass index, and atrial fibrillation, and lobar microbleeds with male sex and atrial fibrillation. Overall, microbleeds were associated with greater white matter hyperintensity volume and lower total white matter fractional anisotropy.

**DISCUSSION:** Results suggest differing associations for lobar versus deep locations. Sensitive microbleed quantification will facilitate future longitudinal studies of their potential role as an early indicator of vascular pathology.

## KEYWORDS

atrial fibrillation, brain microbleeds, brain magnetic resonance imaging, cognition, deep learning, hypertension, Multi-Ethnic Study of Atherosclerosis, race and ethnicity, small vessel disease, white matter fractional anisotropy, white matter hyperintensity

## 1 | BACKGROUND

Brain microbleeds, small hemorrhages less than 10 mm in diameter seen on brain magnetic resonance imaging (MRI), are often manifesta-

tions of small vessel disease and amyloid pathology.<sup>1,2</sup> High microbleed number has been associated with impaired cognitive function and dementia.<sup>3</sup> However, limited information is available about the epidemiology of microbleeds in the general population of older individuals for race or ethnic groups other than White. Previous studies of the epidemiology of microbleeds have relied on manual identification of

Mohamad Habes and Susan R. Heckbert are equally contributing senior authors.

microbleeds,<sup>2,4–6</sup> but recently developed deep learning models using the more sensitive susceptibility-weighted imaging (SWI) sequences at 3-Tesla, rather than gradient-echo imaging,<sup>7</sup> offer the possibility of automated, accurate, and more complete ascertainment of microbleeds.<sup>8,9</sup> To understand the epidemiology of brain microbleeds using these recently developed tools, we analyzed data from the Multi-Ethnic Study of Atherosclerosis (MESA) to examine cross-sectional associations of sociodemographic characteristics and cardiovascular risk factors with microbleed presence, number, and location. We further examined cross-sectional associations of microbleeds with cognitive test scores, physical function, and other aging-related markers on brain MRI.

## 2 | METHODS

### 2.1 | Multi-Ethnic Study of Atherosclerosis

MESA is a community-based cohort study of subclinical cardiovascular disease that enrolled 6814 American men and women 45 to 84 years of age who self-identified with one of four groups: Black, Chinese, Hispanic, or White. Participants were free of clinically recognized cardiovascular disease at baseline in 2000 to 2002 and were recruited at six US field centers (Baltimore, Maryland; Chicago, Illinois; Forsyth County, North Carolina; Los Angeles County, California; New York, New York; and St. Paul, Minnesota).<sup>10</sup> Five follow-up study visits have been conducted, including Exam 6 between 2016 and 2018. During 2018 through 2019, a median of 18 months after Exam 6, consenting participants in the MESA Atrial Fibrillation ancillary study<sup>11</sup> completed a brain MRI.<sup>12</sup> Each field center obtained institutional review board approval, and all participants provided written informed consent.

### 2.2 | Participant characteristics

Participant characteristics obtained at study baseline included self-reported sex, race and ethnicity, and educational attainment. Additional study measures were derived from the 2016 through 2018 (Exam 6) study visit, including age, weight, height, waist circumference, systolic and diastolic blood pressure, cigarette smoking status, medication use, and history of diabetes. Age was also ascertained at the brain MRI visit. Fasting blood specimens from Exam 6 were used to measure total and high-density lipoprotein (HDL) cholesterol; low-density lipoprotein (LDL) cholesterol was calculated using the Friedewald equation. At telephone contacts every 9 to 12 months throughout MESA follow-up, participants were asked to identify new hospitalizations and diagnoses, and medical records were obtained. During follow-up to the date of the brain MRI, clinically recognized myocardial infarction, heart failure,<sup>13</sup> and stroke<sup>14</sup> were adjudicated. Clinically recognized atrial fibrillation was identified from hospital discharge diagnoses and Medicare claims<sup>15</sup> and sub-clinical atrial fibrillation by 14-day ambulatory electrocardiographic monitoring at Exam 6.<sup>11</sup> Apolipoprotein E (APOE) isoforms were

## RESEARCH IN CONTEXT

- 1. Systematic Review:** We searched PubMed for community-based studies of the epidemiology of brain microbleeds. Existing studies included mostly White participants and manual identification of microbleeds; few used 3-Tesla (3T) magnetic resonance imaging (MRI).
- 2. Interpretation:** In the Multi-Ethnic Study of Atherosclerosis, we identified microbleeds from 3T MRI susceptibility-weighted imaging sequences using deep learning models followed by radiologist review. Overall, microbleeds were independently associated with older age and atrial fibrillation and with greater white matter hyperintensity volume and lower white matter fractional anisotropy on MRI. Associations differed for microbleeds in lobar, deep, and infratentorial locations.
- 3. Future Directions:** Longitudinal research is needed to determine the temporal sequence of atrial fibrillation, microbleed development, and the development of other brain MRI markers of small vessel disease.

estimated from single nucleotide polymorphisms rs429358 and rs7412.<sup>16</sup>

### 2.3 | Cognitive test scores and physical function

Four cognitive tests were administered at Exam 6: the Cognitive Abilities Screening Instrument (CASI, version 2), the Digit Symbol Coding Test, and the Digit Span Forward and Backward tests. The CASI was developed for cross-cultural use<sup>17</sup> and includes items representing nine cognitive domains: attention, concentration, orientation, short-term memory, long-term memory, language, visual construction, verbal fluency, and abstraction/judgment; the maximum score is 100. The Digit Symbol Coding Test is a subtest of the Wechsler Adult Intelligence Scale-III<sup>18</sup> that measures processing speed and working memory, with a maximum score of 133. The Digit Span Forward and Backward tests, also subtests of the Wechsler Adult Intelligence Scale-III,<sup>18</sup> assess working memory, with maximum scores of 16 (Forward) and 14 (Backward). Physical function was assessed using the Six-Minute Walk Test (6MWT).<sup>19</sup> The score is expressed in meters walked on a flat surface over 6 minutes.

### 2.4 | Brain MRI and microbleed quantification


Brain MRI was conducted on 3-Tesla scanners as previously described,<sup>12</sup> including 3D T1-weighted, T2-weighted, and fluid attenuated inversion recovery (FLAIR) structural imaging; axial 2D echo-planar diffusion-tensor imaging; and SWI. MRI scanner

parameters are in the Supplemental Methods in supporting information.<sup>20</sup> Microbleeds were initially identified by a deep learning-based method that used T2-weighted quantitative susceptibility mapping and SWI images to segment the lesions and differentiate microbleeds from iron deposits (Figure 1).<sup>8</sup> Identified lesions were then reviewed by a radiologist (JBW) who made the final classification, as previously described.<sup>20</sup> Microbleed location was classified by mapping the 146 Multi-atlas region Segmentation Using Ensembles (MUSE)-based regions of interest<sup>21</sup> to the Microbleed Anatomical Rating Scale (MARS) regions of interest<sup>22</sup> and grouping them into three categories: lobar (frontal, parietal, temporal, occipital, and insula), deep (basal ganglia, thalamus, internal capsule, corpus callosum, and deep and periventricular white matter [WM]), and infratentorial (brainstem and cerebellum). More details on mapping the regions of interest from MUSE to MARS definitions are in the Supplemental Methods and Table S1 in supporting information. Automated measurement of three aging-related MRI markers were made as previously described:<sup>12</sup> total gray matter volume as a measure of atrophy, and WM hyperintensity (WMH) volume and total WM fractional anisotropy as measures of WM injury. Fractional anisotropy, a scalar ranging from 0 to 1, reflects the degree to which water diffusion is limited to a single dimension; lower values are interpreted as indicating reduced WM microstructural integrity, a feature of small vessel disease.


## 2.5 | Statistical analysis

In our primary analyses, we compared participants with at least one microbleed to those without, both overall and in each of the lobar, deep, and infratentorial location categories. We used multivariable Poisson relative risk regression with robust standard errors to examine cross-sectional associations of participant characteristics and cardiovascular risk factors with the presence of microbleeds.<sup>23</sup> This analysis included age, sex, race and ethnicity, field center, educational attainment, body mass index, systolic blood pressure, diastolic blood pressure, use of antihypertensive medication, HDL and LDL cholesterol levels, APOE genotype, cigarette smoking status, diabetes mellitus, history of atrial fibrillation, and history of myocardial infarction or heart failure as covariates in a single model. Next, we used multivariable linear regression to examine cross-sectional associations of microbleed presence, both overall and by location, with the outcomes of cognitive test scores, 6MWT, and three brain MRI measures: total gray matter volume, WMH volume, and total WM fractional anisotropy. Models for total gray matter volume and WMH volume included additional adjustment for total intracranial volume. WMH volume had a right-skewed distribution and was natural log transformed. Differences in test scores, 6MWT distance, and brain MRI findings were expressed in standard deviation units.

In secondary analyses limited to participants with at least one microbleed, we analyzed participant characteristics and cardiovascular risk factors in relation to the number of microbleeds, both overall and by location, using linear regression with robust standard errors. Six participants had more than 12 microbleeds (81% of which were lobar);



**FIGURE 1** Examples of microbleeds detected in lobar, deep, and infratentorial regions segmented by our in-house-developed deep learning model trained with susceptibility weighted, T2-weighted, and quantitative susceptibility mapping images. The unmagnified views show the microbleed segmentation masks (in red, 2–10 voxels in volume). The magnified views show the same microbleeds indicated by red arrows and without the segmentation mask. CMB, cerebral microbleed.



**FIGURE 2** Study inclusion diagram. MESA, Multi-Ethnic Study of Atherosclerosis; MRI, magnetic resonance imaging, SWI, susceptibility-weighted imaging sequence.

in analyses of microbleed counts these values were winsorized at 12 microbleeds. The number of microbleeds was log transformed, and results of multivariable analyses are presented as the geometric mean ratio, which provides the percentage difference in microbleed number per increment of the characteristic. For example, a geometric mean ratio of 1.07 for the association with age indicates a 7% higher number of microbleeds per 10-year increment of age. Next, we used linear regression to examine associations of log base 2-transformed microbleed number with cognitive test scores, 6MWT, and aging-related findings on brain MRI, expressed in standard deviation units.

Multiple imputation with chained equations was used to impute missing values of education, systolic and diastolic blood pressure, treated hypertension, HDL and LDL cholesterol, smoking, prevalent diabetes, and APOE genotype (all < 5% missing) using information on age, sex, race, body mass index, and history of myocardial infarction, heart failure, or atrial fibrillation. In sensitivity analyses, we examined associations with microbleeds for our main models after excluding participants who were using anticoagulant medications at Exam 6.

### 3 | RESULTS

Among the 3303 participants who attended Exam 6 (2016–2018), 1062 had a brain MRI (Figure 2). SWI sequence data were unavailable for 4, MRI images did not meet quality control criteria for 16, and we excluded 26 with a history of clinically recognized stroke before the MRI, leaving 1016 in the analysis. Participants self-reported Black (25%), Chinese (15%), Hispanic (19%), or White (41%) race and

ethnicity; the average (standard deviation [SD]) age at the time of MRI was 72 (8) years and 47% were men. Compared to Exam 6 participants not included in the analysis, those included were on average younger, a smaller proportion used antihypertensive medication, and a smaller proportion had experienced a myocardial infarction or heart failure before the brain MRI (Table S2 in supporting information).

On brain MRI, a total of 339 (33%) participants had at least one microbleed, of which 188 (55%) had a single microbleed, 68 (20%) had 2, 76 (22%) had 3 to 11, and one each had 12, 17, 20, 21, 25, 54, and 112 microbleeds (Figure 3A). Microbleeds were widely distributed in the brain (Figure 3B, Video S1 in supporting information, and Table S1); of the total of 947 microbleeds detected, 67% were lobar, 22% deep, and 12% infratentorial. The prevalence of microbleeds increased with age (Figure 4) from 20% of participants aged 60 to 64.9 years to 45% of those over age 85. Compared to those with no microbleeds, participants with microbleeds were slightly older and had higher systolic blood pressure but lower LDL cholesterol, and a larger proportion were men, had treated hypertension, and had a history of atrial fibrillation (Table 1).



#### 3.1 | Multivariable associations of sociodemographic characteristics and cardiovascular risk factors with microbleeds

In a multivariable regression model, only older age and a history of atrial fibrillation were independently associated with microbleed presence overall (Figure 5 and Table S3 in supporting information). There was no association of use of anticoagulant (prevalence ratio = 0.95; 95% confidence interval [CI] 0.63–1.45) or antiplatelet agents (prevalence ratio = 0.99; 95% CI 0.82–1.18) with overall microbleed presence when added to this multivariable model. Lobar, deep, and infratentorial locations showed differing associations with risk factors. Older age was associated with both deep and infratentorial microbleeds, but not lobar microbleeds. Male sex was associated with lobar microbleeds but not with microbleeds at deep or infratentorial sites. Higher systolic blood pressure and treated hypertension were uniquely associated with deep microbleeds. A history of atrial fibrillation was associated with microbleeds in both lobar and deep locations. No associations were detected, either overall or by location, with diastolic blood pressure, diabetes, or the APOE ε4 allele.

#### 3.2 | Multivariable associations of microbleeds with cognitive test performance, 6MWT, and brain MRI measures

In multivariable regression analyses, microbleed presence versus absence was not associated with lower scores on any of the four cognitive tests or with lower 6MWT distance (Figure 6 and Table S4 in supporting information). However, microbleed presence overall and in each of the three investigated location categories was associated with greater WMH volume, and total and deep microbleeds were





**FIGURE 3** A, By location, number of microbleeds per person among 339 participants with at least one microbleed. B, Three-dimensional rendering of all microbleeds observed in the Multi-Ethnic Study of Atherosclerosis sample; orange color indicates anatomical locations where two microbleeds were observed; brighter yellow areas are more superficial while indistinct areas are deeper. Also see [Video S1](#) in supporting information.

associated with lower total WM fractional anisotropy (Figure 6 and Table S5 in supporting information). The associations with WM fractional anisotropy were little changed by additional adjustment for WMH volume. No association was detected between microbleed presence and total gray matter volume.

### 3.3 | Among participants with at least one microbleed, multivariable associations with number of microbleeds


In multivariable analyses among those with at least one microbleed, associations of sociodemographic characteristics and cardiovascular risk factors with greater microbleed number were similar in direction

to the associations for presence versus absence of microbleeds (Table S6 in supporting information). Greater microbleed number both overall and in lobar locations was associated with greater WMH volume and lower total WM fractional anisotropy (Table S5). No association was detected between microbleed number and cognitive test scores, 6MWT distance, or total gray matter volume.

In sensitivity analyses, exclusion of participants using anticoagulant medication did not materially change the results.

## 4 | DISCUSSION

In this analysis of a racially and ethnically diverse older community-based cohort with mean (SD) age of 72 (8) years and without prior



**FIGURE 4** Microbleed prevalence by age group among 1016 Multi-Ethnic Study of Atherosclerosis participants, 2018–2019. Bars indicate 95% confidence intervals.

clinically recognized stroke, brain microbleeds were present in 33% of participants overall and in 45% of participants aged  $\geq 85$  years. In cross-sectional multivariable analysis, advanced age, higher systolic blood pressure and body mass index, treated hypertension, and a history of atrial fibrillation were independently associated with deep microbleeds. Male sex and a history of atrial fibrillation were independently associated with lobar microbleeds. Both overall and in all three locations, microbleeds were strongly and independently associated with one or both brain MRI markers of WM injury: greater WMH volume and lower total WM fractional anisotropy.

Several strengths and limitations of our analysis should be noted. Strengths include the diverse sample of older participants and the assessment of both clinically recognized and subclinical atrial fibrillation. The brain MRI used high field strength (3-Tesla) and microbleeds were detected from SWI sequences using an automated and validated method followed by radiologist review,<sup>8</sup> and automated methods were used for WMH measurement.<sup>12</sup> We were able to examine associations with microbleeds by location within the brain. However, because our analyses were cross-sectional, we were unable to assess temporal relationships between study measures and microbleed development. Also, brain MRI readings of subclinical or covert infarcts, another important direct measure of cerebrovascular disease, were not available in this MESA dataset. Our analysis of associations with cognitive performance, physical function, and brain MRI markers of small vessel disease included several tests for significance, and type 1 error is possible as a result. However, correction for multiple testing would be overly conservative because the various measures are highly correlated with one another. We have presented point estimates and 95% CIs without focusing exclusively on statistical significance. Participants able and willing to complete the brain MRI had fewer cardiovascular risk factors than those who did not; this bias would likely lead to underestimates for the prevalence of microbleeds at each age in the population at large.

Nonetheless, the measured prevalence of microbleeds was higher at comparable ages in MESA than in the UK Biobank, the only other large community-based study that used 3-Tesla field strength and SWI sequence data to identify microbleeds. Microbleed prevalence was 7%

at a mean (SD) age of 62 (7) years in the UK Biobank,<sup>24</sup> but 20% among those aged 60 to 64.9 years in MESA. The higher microbleed prevalence at comparable ages in MESA may be due to greater sensitivity of the deep learning method for microbleed detection as opposed to manual detection the UK Biobank. The higher prevalence may also be related to differences in cohort characteristics including the higher prevalence of diabetes and higher average body mass index in MESA participants. As expected, the use of 3-Tesla field strength and SWI sequence data<sup>25</sup> in MESA yielded higher age-specific microbleed prevalence than in community-based studies that did not use these techniques, including the Framingham Heart Study;<sup>4</sup> Rotterdam Scan Study;<sup>2</sup> Age, Gene/Environment Susceptibility Study;<sup>5</sup> and Atherosclerosis Risk in Communities (ARIC) study.<sup>6</sup>

We found that a history of atrial fibrillation was strongly associated with microbleed prevalence both overall and in deep and lobar locations. This association has not been examined to date in most large community-based studies, but the Gothenburg H70 Birth Cohort study of 776 individuals examined at age 70 reported an association of atrial fibrillation with frontal microbleeds after adjustment for other cardiovascular risk factors.<sup>26</sup> A clinical study reported higher microbleed prevalence in patients with atrial fibrillation than in those without, but the analyses were not adjusted for other microbleed risk factors.<sup>27</sup> We previously showed that greater left atrial volume index was associated with microbleeds in MESA participants.<sup>20</sup> Taken together, these findings support the involvement of atrial fibrillation in the pathophysiology of microbleeds, but do not provide insight as to whether atrial fibrillation is causally related to microbleeds or is a manifestation of underlying vascular pathology or shared risk factors.


In multiply adjusted analyses in the multi-ethnic MESA population, race and ethnicity were not associated with microbleed presence, either overall or by location. However, our finding of a prevalence ratio of 1.62 for deep microbleeds in Chinese versus White participants is intriguing given previous findings of a higher prevalence of deep and/or infratentorial or mixed microbleeds in Eastern (Japanese and Chinese) than in White Western participants.<sup>28</sup> Additional investigation will be needed to elucidate a possible relation to the 2-fold higher observed

**TABLE 1** Characteristics of 1016 MESA participants in 2016–2018 (Exam 6) by presence and number of microbleeds on brain MRI.

	No microbleeds (N = 677)		1+ microbleed (N = 339)		1 microbleed (N = 188)		2 microbleeds (N = 68)		≥3 microbleeds (N = 83)	
	Mean or %	SD	Mean or %	SD	Mean or %	SD	Mean or %	SD	Mean or %	SD
Age, years	71	8	74	8	74	8	74	8	76	9
Sex, % male	45%		50%		47%		47%		60%	
Race, %										
Black	23%		27%		28%		29%		25%	
Chinese	14%		17%		16%		18%		18%	
Hispanic	22%		15%		12%		21%		19%	
White	41%		40%		44%		32%		37%	
Education, %										
<High school	11%		13%		14%		12%		12%	
High school graduate	17%		13%		11%		16%		14%	
Some college	32%		25%		26%		21%		27%	
College graduate	41%		49%		49%		51%		47%	
Body mass index, kg/m <sup>2</sup>	28	6	28	5	27	5	29	4	29	5
Height, cm	165	10	165	10	165	10	165	10	165	9
Weight, kg	77	17	77	15	75	16	78	14	79	15
Waist circumference, cm	98	13	99	12	97	13	100	12	102	10
Systolic BP, mmHg	125	19	129	22	128	22	127	20	134	22
Diastolic BP, mmHg	69	10	69	10	69	11	68	10	70	9
Treated hypertension, %	54%		66%		62%		64%		77%	
Smoking history, %										
Current	6%		5%		4%		6%		7%	
Past	45%		49%		49%		49%		49%	
HDL cholesterol, mg/dL	60	19	61	18	63	20	58	16	57	13
LDL cholesterol, mg/dL	111	35	102	34	103	35	99	34	101	31
Anticoagulation use, %	3%		5%		4%		4%		8%	
Antiplatelet* use, %	43%		48%		46%		51%		51%	
APOE, ≥1 ε4 allele, %	28%		27%		24%		24%		35%	
Diabetes mellitus, %	20%		25%		24%		28%		26%	
History of AF, %	10%		19%		15%		22%		28%	
History of MI or heart failure, %	3%		4%		3%		3%		7%	
CASI score	90	7	90	7	90	7	90	6	89	8
Digit Symbol Coding score	53	18	50	18	52	19	48	16	48	16
Digit Span Forward score	10	3	10	3	10	3	10	3	10	3
Digit Span Backward score	6	2	6	2	6	2	6	2	5	2
Six-Minute Walk Test, m	426	95	419	93	427	88	420	109	397	89
Total brain volume, mL	1094	111	1089	118	1083	118	1095	125	1095	112
Total gray matter volume, mL	600	63	591	67	589	66	596	75	593	64
WMH volume, mL	5.6	9.0	9.2	12.5	7.6	11.1	7.4	8.3	14.3	16.4
Fractional anisotropy	0.40	0.02	0.40	0.03	0.39	0.03	0.39	0.03	0.38	0.03

Abbreviations: AF, atrial fibrillation; APOE, apolipoprotein E; BP, blood pressure; CASI, Cognitive Abilities Screening Instrument; HDL, high-density lipoprotein cholesterol; LDL, low-density lipoprotein cholesterol; MESA, Multi-Ethnic Study of Atherosclerosis; MI, myocardial infarction; MRI, magnetic resonance imaging; SD, standard deviation; WMH, white matter hyperintensity.

\*Antiplatelet agents included regular aspirin (at least 3 days per week) or adenosine diphosphate receptor inhibitors.



**FIGURE 5** Multivariable\* association of sociodemographic characteristics and cardiovascular risk factors with the presence of brain microbleeds, overall and by microbleed location. \* Also adjusted for field center, educational attainment, smoking (never, former, current), and HDL cholesterol. APOE, apolipoprotein E; BMI, body mass index; BP, blood pressure; HDL, high-density lipoprotein cholesterol; LDL, low-density lipoprotein cholesterol; MI, myocardial infarction.

incidence of intracerebral hemorrhage in East and Southeast Asian compared with White populations.<sup>29</sup>


Our findings that older age, higher systolic blood pressure, and treated hypertension were associated with deep microbleeds are in accord with findings from other community-based studies,<sup>2,24</sup> as is the lack of association with diabetes or hemoglobin A1c. The strong association of deep microbleeds with high blood pressure and with MRI markers of small vessel disease support the hypothesis that deep microbleeds are mainly of vascular origin in distinction to lobar microbleeds, which may reflect amyloid-related pathologies.<sup>30</sup>

We did not detect an association of lobar microbleeds with APOE ε4 genotype, the strongest genetic marker of Alzheimer's disease; such an association has been reported in studies of largely White populations.<sup>2,24,31</sup> The multi-ethnic character of MESA may explain this discrepancy, in that the contribution of APOE ε4 allele to alterations in cognition and brain health may vary by race and ethnicity.<sup>32,33</sup> Despite the higher proportion reporting use of anticoagulants among

participants with more microbleeds, after adjustment for sociodemographic characteristics and cardiovascular risk factors, we found no association of anticoagulant use with microbleeds, in agreement with the Rotterdam and ARIC studies.<sup>34,35</sup> In addition, we report that use of antiplatelet agents was not associated with microbleeds in adjusted analyses. Associations of microbleeds with other risk factors were not affected by exclusion of participants who used anticoagulant medication.

Previous studies of clinical and community-based samples have described associations of microbleeds with cognitive impairment,<sup>36,37</sup> cognitive decline,<sup>3</sup> and dementia<sup>3,5,38</sup> and with elevation in cerebrospinal fluid amyloid beta and phosphorylated tau 181 protein levels.<sup>36</sup> In cross-sectional analyses in MESA, we did not detect an association of microbleeds with cognitive test scores. However, the cognitive tests administered in MESA were limited and did not provide a comprehensive assessment of all cognitive domains. Thus, based on findings in previous studies, microbleeds should not be considered





**FIGURE 6** Adjusted associations\* of microbleed presence with cognitive test performance, Six-Minute Walk Test distance, and brain MRI measures, overall and by microbleed location. \*All models adjusted for age, sex, race and ethnicity, field center, educational attainment, body mass index, systolic blood pressure, diastolic blood pressure, use of antihypertensive medication, HDL and LDL cholesterol, cigarette smoking status, diabetes mellitus, history of atrial fibrillation, and history of myocardial infarction or heart failure. Total gray matter volume and WM hyperintensity volume models are also adjusted for total intracranial volume. CASI, Cognitive Abilities Screening Instrument; HDL, high-density lipoprotein cholesterol; LDL, low-density lipoprotein cholesterol; MRI, magnetic resonance imaging; SD, standard deviation; WM, white matter.

benign with respect to cognition. Future analysis in MESA of change in cognitive test scores over time and adjudicated cognitive status will provide a more sensitive assessment of associations of microbleeds with cognition.

In cross-sectional but not in longitudinal analyses in the ARIC study, greater microbleed number was associated with slower gait speed.<sup>39</sup> In our cross-sectional analysis, we did not detect an association with distance walked. Both the present study and the UK Biobank found strong associations of microbleed presence with greater WMH volume and total WM fractional anisotropy but not with gray matter volume.<sup>24</sup> Longitudinal analysis of repeated brain MRIs will be needed to elucidate the temporal sequence of development of microbleeds, reduction in WM fractional anisotropy, and increase in WMH volume.

## 5 | CONCLUSION

The results of our analysis of brain microbleeds in a diverse community-based cohort of older individuals without prior stroke suggests that microbleeds are common and are associated with older age, atrial fibrillation, and MRI measures of WM injury. Our findings are consistent with the existing view that deep microbleeds reflect vascular brain injury. Sensitive quantification of microbleeds will facilitate future longitudinal studies of the potential role of microbleeds as an early indicator of vascular pathology.

## ACKNOWLEDGMENTS

The authors thank the other investigators, the staff, and the participants of the MESA study for their valuable contributions. A full list of participating MESA investigators and institutions can be found at <http://www.mesa-nhlbi.org>. This MESA research was supported by contracts 75N92020D00001, HHSN268201500003I, N01-HC-

95159, 75N92020D00005, N01-HC-95160, 75N92020D00002, N01-HC-95161, 75N92020D00003, N01-HC-95162, 75N92020D00006, N01-HC-95163, 75N92020D00004, N01-HC-95164, 75N92020D00007, N01-HC-95165, N01-HC-95166, N01-HC-95167, N01-HC-95168, and N01-HC-95169 and grants R01 HL127659 and R01 HL127028 from the National Heart, Lung, and Blood Institute, and by grants UL1-TR-000040, UL1-TR-001079, and UL1-TR-001420 from the National Center for Advancing Translational Sciences (NCATS). Additional support was provided by the National Institute on Aging and grant R01 AG080821. The funding sources had no role in the study design, the analysis and interpretation of data, the writing of the report, or the decision to submit the article for publication.

## CONFLICT OF INTEREST STATEMENT

The authors declare no conflicts of interest. Author disclosures are available in the [supporting information](#).

## CONSENT STATEMENT

All Multi-Ethnic Study of Atherosclerosis participants provided written informed consent.

## ORCID

Susan R. Heckbert <https://orcid.org/0000-0002-7100-512X>

## REFERENCES

- Gouw AA, Seewann A, van der Flier WM, et al. Heterogeneity of small vessel disease: a systematic review of MRI and histopathology correlations. *Journal of neurology, neurosurgery, and psychiatry*. 2011;82:126-135. doi:10.1136/jnnp.2009.204685
- Vernooij MW, van der Lugt A, Ikram MA, et al. Prevalence and risk factors of cerebral microbleeds: the Rotterdam Scan Study. *Neurology*. 2008;70:1208-1214. doi:10.1212/01.wnl.0000307750.41970.d9

3. Akoudad S, Wolters FJ, Viswanathan A, et al. Association of cerebral microbleeds with cognitive decline and dementia. *JAMA Neurol.* 2016;73:934-943. doi:10.1001/jamaneurol.2016.1017
4. Romero JR, Preis SR, Beiser A, et al. Risk factors, stroke prevention treatments, and prevalence of cerebral microbleeds in the Framingham Heart Study. *Stroke.* 2014;45:1492-1494. doi:10.1161/STROKEAHA.114.004130
5. Ding J, Sigurdsson S, Jonsson PV, et al. Space and location of cerebral microbleeds, cognitive decline, and dementia in the community. *Neurology.* 2017;88:2089-2097. doi:10.1212/WNL.0000000000003983
6. Graff-Radford J, Simino J, Kantarci K, et al. Neuroimaging correlates of cerebral microbleeds: the ARIC Study (Atherosclerosis Risk in Communities). *Stroke.* 2017;48:2964-2972. doi:10.1161/STROKEAHA.117.018336
7. Haller S, Vernooij MW, Kuijper JPA, Larsson EM, Jager HR, Barkhof F. Cerebral microbleeds: imaging and clinical significance. *Radiology.* 2018;287:11-28. doi:10.1148/radiol.2018170803
8. Rashid T, Abdulkadir A, Nasrallah IM, et al. DEEPNIR: a deep neural network for differential detection of cerebral microbleeds and iron deposits in MRI. *Sci Rep.* 2021;11:14124. doi:10.1038/s41598-021-93427-x
9. Sundaresan V, Arthofer C, Zamboni G, et al. Automated detection of candidate subjects with cerebral microbleeds using machine learning. *Front Neuroinform.* 2021;15:777828. doi:10.3389/fninf.2021.777828
10. Bild DE, Bluemke DA, Burke GL, et al. Multi-Ethnic Study of Atherosclerosis: objectives and design. *Am J Epidemiol.* 2002;156:871-881.
11. Heckbert SR, Austin TR, Jensen PN, et al. Differences by race/ethnicity in the prevalence of clinically detected and monitor-detected atrial fibrillation: mESA. *Circulation Arrhythmia and electrophysiology.* 2020;13:e007698. doi:10.1161/CIRCEP.119.007698
12. Austin TR, Nasrallah IM, Erus G, et al. Association of brain volumes and white matter injury with race, ethnicity, and cardiovascular risk factors: the multi-ethnic study of atherosclerosis. *Journal of the American Heart Association.* 2022;11:e023159. doi:10.1161/JAHA.121.023159
13. Bluemke DA, Kronmal RA, Lima JA, et al. The relationship of left ventricular mass and geometry to incident cardiovascular events: the MESA (Multi-Ethnic Study of Atherosclerosis) study. *J Am Coll Cardiol.* 2008;52:2148-2155. doi:10.1016/j.jacc.2008.09.014
14. Longstreth WT Jr, Gasca NC, Gottesman RF, Pearce JB, Sacco RL. Adjudication of transient ischemic attack and stroke in the multi-ethnic study of atherosclerosis. *Neuroepidemiology.* 2018;50:23-28. doi:10.1159/000486174
15. Austin TR, Wiggins KL, Blackshear C, et al. Atrial fibrillation in an African-American cohort: the Jackson Heart Study. *Clin Cardiol.* 2018;41:1049-1054. doi:10.1002/clc.23020
16. Fitzpatrick AL, Rapp SR, Luchsinger J, et al. Sociodemographic correlates of cognition in the Multi-Ethnic Study of Atherosclerosis (MESA). *The American journal of geriatric psychiatry : official journal of the American Association for Geriatric Psychiatry.* 2015;23:684-697. doi:10.1016/j.jagp.2015.01.003
17. Teng EL, Hasegawa K, Homma A, et al. The Cognitive Abilities Screening Instrument (CASI): a practical test for cross-cultural epidemiological studies of dementia. *Int Psychogeriatr.* 1994;6:45-58.
18. Wechsler D. *Wechsler Adult Intelligence Scale-III (WAIS-III).* Psychological Corporation/Harcourt, Inc; 1996.
19. ATS. Committee on proficiency standards for clinical pulmonary function laboratories. ATS statement: guidelines for the six-minute walk test. *Am J Respir Crit Care Med.* 2002;166:111-117. doi:10.1164/ajrccm.166.1.at1102
20. Austin TR, Jensen PN, Nasrallah IM, et al. Left atrial function and arrhythmias in relation to small vessel disease on brain MRI: the Multi-Ethnic Study of Atherosclerosis. *Journal of the American Heart Association.* 2022;11:e026460. doi:10.1161/JAHA.122.026460
21. Doshi J, Erus G, Ou Y, et al. Alzheimer's Neuroimaging Initiative. MUSE: mUlti-atlas region Segmentation utilizing Ensembles of registration algorithms and parameters, and locally optimal atlas selection. *Neuroimage.* 2016;127:186-195. doi:10.1016/j.neuroimage.2015.11.073
22. Gregoire SM, Chaudhary UJ, Brown MM, et al. The Microbleed Anatomical Rating Scale (MARS): reliability of a tool to map brain microbleeds. *Neurology.* 2009;73:1759-1766. doi:10.1212/WNL.0b013e3181c34a7d
23. Zou G. A modified Poisson regression approach to prospective studies with binary data. *Am J Epidemiol.* 2004;159:702-706. doi:10.1093/aje/kwh090
24. Lu D, Liu J, MacKinnon AD, Tozer DJ, Markus HS. Prevalence and risk factors of cerebral microbleeds: an analysis from the UK Biobank. *Neurology.* 2021. doi:10.1212/WNL.0000000000012673
25. Cheng AL, Batool S, McCreary CR, et al. Susceptibility-weighted imaging is more reliable than T2\*-weighted gradient-recalled echo MRI for detecting microbleeds. *Stroke.* 2013;44:2782-2786. doi:10.1161/STROKEAHA.113.002267
26. Ryden L, Sacuiu S, Wetterberg H, et al. Atrial fibrillation, stroke, and silent cerebrovascular disease: a population-based MRI study. *Neurology.* 2021;97:e1608-e1619. doi:10.1212/WNL.0000000000012675
27. Saito T, Kawamura Y, Tanabe Y, et al. Cerebral microbleeds and asymptomatic cerebral infarctions in patients with atrial fibrillation. *Journal of stroke and cerebrovascular diseases : the official journal of National Stroke Association.* 2014;23:1616-1622. doi:10.1016/j.jstrokecerebrovasdis.2014.01.005
28. Yakushiji Y, Wilson D, Ambler G, et al. Distribution of cerebral microbleeds in the East and West: individual participant meta-analysis. *Neurology.* 2019;92:e1086-e1097. doi:10.1212/WNL.0000000000007039
29. van Asch CJ, Luitse MJ, Rinkel GJ, van der Tweel I, Algra A, Klijn CJ. Incidence, case fatality, and functional outcome of intracerebral haemorrhage over time, according to age, sex, and ethnic origin: a systematic review and meta-analysis. *Lancet neurology.* 2010;9:167-176. doi:10.1016/S1474-4422(09)70340-0
30. Park JH, Seo SW, Kim C, et al. Pathogenesis of cerebral microbleeds: in vivo imaging of amyloid and subcortical ischemic small vessel disease in 226 individuals with cognitive impairment. *Annals of neurology.* 2013;73:584-593. doi:10.1002/ana.23845
31. Ingala S, Mazzai L, Sudre CH, et al. The relation between APOE genotype and cerebral microbleeds in cognitively unimpaired middle- and old-aged individuals. *Neurobiol Aging.* 2020;95:104-114. doi:10.1016/j.neurobiolaging.2020.06.015
32. Mayeux R, Apolipoprotein E. Alzheimer disease, and African Americans. *Arch Neurol.* 2003;60:161-163. doi:10.1001/archneur.60.2.161
33. Le Guen Y, Raulin AC, Logue MW, et al. Association of African Ancestry-Specific APOE Missense Variant R145C With Risk of Alzheimer Disease. *JAMA.* 2023;329:551-560. doi:10.1001/jama.2023.0268
34. Vernooij MW, Haag MD, van der Lugt A, et al. Use of antithrombotic drugs and the presence of cerebral microbleeds: the Rotterdam Scan Study. *Arch Neurol.* 2009;66:714-720. doi:10.1001/archneur.2009.42
35. Sharma R, Matsushita K, Wu A, et al. Common Medications and Intracerebral Hemorrhage: the ARIC Study. *Journal of the American Heart Association.* 2021;10:e014270. doi:10.1161/JAHA.120.014270
36. Sparacia G, Agnello F, La Tona G, Iaia A, Midiri F, Sparacia B. Assessment of cerebral microbleeds by susceptibility-weighted imaging in Alzheimer's disease patients: a neuroimaging biomarker of the disease. *Neuroradiol J.* 2017;30:330-335. doi:10.1177/1971400916689483

37. Paradise M, Seruga A, Crawford JD, et al. The relationship of cerebral microbleeds to cognition and incident dementia in non-demented older individuals. *Brain Imaging Behav.* 2019;13:750-761. doi:[10.1007/s11682-018-9883-3](https://doi.org/10.1007/s11682-018-9883-3)
38. Miwa K, Tanaka M, Okazaki S, et al. Multiple or mixed cerebral microbleeds and dementia in patients with vascular risk factors. *Neurology.* 2014;83:646-653. doi:[10.1212/WNL.0000000000000692](https://doi.org/10.1212/WNL.0000000000000692)
39. Sullivan KJ, Ranadive R, Su D, et al. Imaging-based indices of neuropathology and gait speed decline in older adults: the Atherosclerosis Risk in Communities study. *Brain Imaging Behav.* 2021;15:2387-2396. doi:[10.1007/s11682-020-00435-y](https://doi.org/10.1007/s11682-020-00435-y)

## SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

**How to cite this article:** Jensen PN, Rashid T, Ware JB, et al. Association of brain microbleeds with risk factors, cognition, and MRI markers in MESA. *Alzheimer's Dement.* 2023;19:4139-4149. <https://doi.org/10.1002/alz.13346>