

1. In each question part below, I will list a set S and a rule that defines a relation R on S as follows: $(m, n) \in R$ if m and n satisfy the given rule. For each set and rule, do the following five things:

- i List the order pairs in the relation.
- ii Draw the associated arrow diagram.
- iii Give the matrix representation of the relation. Be sure to label the rows and columns with the elements of S .
- iv Identify which properties the relation has: reflexive, antireflexive, or neither; symmetric, antisymmetric, or neither; transitive or not transitive. Be sure to explain your answers.
- v Decide whether the relation is an equivalence relation. If so, identify the equivalence classes.

(a) $S = \{4, 9, 17\}; m \geq n$.


(b) $S = \{0, 2, 5\}; mn = 0$.

(c) $S = \{1, 2, 6, 7, 11\}; m \equiv n \pmod{3}$

Answers for (a):

- $R = \{(m, n) : m \geq n \text{ and } m, n \in S\}$
 $\Rightarrow \{(17, 4), (9, 4), (17, 9), (4, 4), (9, 9), (17, 17)\}$

- Diagram:




- Matrix:

$$\begin{matrix} & \begin{matrix} 4 & 9 & 17 \end{matrix} \\ \begin{matrix} 4 \\ 9 \\ 17 \end{matrix} & \begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 1 & 1 & 1 \end{bmatrix}
 \end{matrix}
 \text{ Matrix} = \begin{cases} 1 : (m, n) \in R \\ 0 : \text{otherwise} \end{cases}$$
- - R is reflexive as $(m, n) \in R \forall m \in S$
 - R is anti symmetric. As if $(a, b) \in R$ and $(b, a) \in R$ then $a = b$ for:
 - $a = b = 4$
 - $a = b = 9$
 - $a = b = 17$
 - R is transitive. As if $(a, b) \in R$ and $(b, c) \in R$ then $(a, c) \in R$ holds. $\therefore a, b, c \in S$
- R is not an equivalence relation based on the reasons mentioned above.

Answers for (b):

- $R = \{(m, n) : mn = 0 \text{ and } m, n \in S\}$
 $\Rightarrow \{(0,0), (0,2), (0,5), (2,0), (5,0)\}$

• Diagram:



• Matrix:


$$\begin{matrix} & 0 & 2 & 5 \\ \begin{matrix} 0 \\ 2 \\ 5 \end{matrix} & \begin{bmatrix} 1 & 1 & 1 \\ 1 & 0 & 0 \\ 1 & 0 & 0 \end{bmatrix} \end{matrix} \text{ Matrix} = \begin{cases} 1 : mn = 0 \\ 0 : otherwise \end{cases}$$

- R is not reflexive as $(2, 2) \notin R$
- R is symmetric as $(0, 2) \in R$ and $(2, 0) \in R$
- R is not transitive. Because $(2, 0)$ and $(0, 5) \in R$ but $(2, 5) \notin R$
- R is not an equivalence relation based on the reasons mentioned above.

Answers for (c):

- $R = \{(m, n) : m \equiv n \pmod{3} \text{ and } m, n \in S\}$
 $\Rightarrow \{(1,1), (2,2), (6,6), (7,7), (11,11), (1,7), (7,1), (2,11), (11,2)\}$

• Diagram:



• Matrix:

$$\begin{matrix} & 1 & 2 & 6 & 7 & 11 \\ \begin{matrix} 1 \\ 2 \\ 6 \\ 7 \\ 11 \end{matrix} & \begin{bmatrix} 1 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 1 \end{bmatrix} \end{matrix} \text{ Matrix} = \begin{cases} 1 : m \equiv n \pmod{3} \\ 0 : otherwise \end{cases}$$

- R is reflexive as the diagonal of the matrix consist on only 1's.
- R is symmetric as since $\forall m$ and n , (m, n) and (n, m) are 1's.
- R is transitive as $\forall m$ and $n \in R$, if mRn and nRm then mRm and nRn .
- R is an equivalence relation since it is reflexive, symmetric, and transitive for the reasons mentioned above. The equivalence class for R is

2. Suppose X is the set $\{a, b, c, d, e\}$ and $P(X)$ is the power set of X . The domain of each of the relations below is $P(X)$. For each relation, describe the relation in words. Then determine whether it is reflexive, antireflexive, or neither; symmetric, anti-symmetric, or neither; and transitive or not transitive.

- (a) A is related to B if $|A - B| = 1$. The vertical bars mean the cardinality of $A - B$.
 (b) A is related to B if $A \subset B$. Note that this notation means proper subset.

- i. Let $A \in P(X)$:

\Rightarrow Now, $A - A = \emptyset$

$\Rightarrow |A - A| = 0 \neq 1$

$\Rightarrow A \not\sim A$.

Since the choice of set A is completely arbitrary, we can say that set A is **anit-reflexive**.

- ii. Let $A = \{a\}$, $B = \emptyset$ and $A, B \in P(X)$:

$\Rightarrow A - B = \{a\}$ and $B - A = \emptyset$

$\Rightarrow |A - B| = 1$ and $|B - A| = 0$

\Rightarrow From this we can say that $A \sim B$ but $B \not\sim A$

This results in the conclusion that the relation is **not symmetric**.

- iii. Let $A = \{a, b\}$, $B = \{b, a\}$ and $A, B \in P(X)$:

$\Rightarrow A - B = \{a\}$ and $B - A = \{c\}$

$\Rightarrow |A - B| = 1$ and $|B - A| = 1$

\Rightarrow From this we can say that $A \sim B$ and $B \sim A$ but $A \neq B$

We can say that the relation is not antisymmetric. \therefore The relation is

neiter symmetric nor antisymmetric.

- iv. Let $A = \{a, b\}$, $B = \{b, c\}$ and $A, B \in P(X)$:

$\Rightarrow A - B = \{a\}$ and $B - A = \{c\}$

$\Rightarrow |A - B| = 1$ and $|B - A| = 1$

\Rightarrow From this we can say that $A \sim B$ and $B \sim A$

But from the (i) we said that $A \not\sim A$,

\therefore The relation is **not transitive**.

In short, the relation is :

- **antireflexive**
- **neither symmetric and anti-symmetric**
- **not transitive**

- i. Since every element of a set A is contained in itself, $A \sim A \forall A$ and it makes the relation **reflexive**

- ii. Let $A \sim B$ and $B \sim A$. Here as per definition, every element of $A \in B$ and $B \in A$ also however $A \neq B$.

\therefore The relation is **antisymmetric**.

- iii. Let $A \sim B$ and $B \sim C$, then it means that $A \in B$ and $B \in C$. That automatically puts every element of A inside C . So $A \sim C$ and hence the relation is **transitive**.

In short, the relation is :

- **reflexive**
- **antisymmetric**
- **transitive**

3. Determine whether the following relations are equivalence relations. If so, show that the relation fulfills the requirements for an equivalence relation, and describe the equivalence classes. If not, show how the relation fails to fulfill the requirements.

(a) The relation \mathcal{C} is defined by $(x, y) \in \mathcal{C}$ iff $\cos(x) = \cos(y)$, where $x, y \in \mathbb{R}$.

- Reflexive:

Let $y = x$. From this, $(x, x) \in \mathcal{C}$.

$\Rightarrow \therefore \cos(x) = \cos(x)$ holds.

Similarly, let $x = y$. From this (y, y)

$\Rightarrow \therefore \cos(y) = \cos(y)$ also holds.

Reflexivity holds.

- Symmetric:

If $x \sim y$, then it implies that $y \sim x$.

Assume that $(x, y) \in \mathcal{C}$ then $\cos(x) = \cos(y)$ can be rewritten as $\cos(y) = \cos(x)$.

This implies that $(y, x) \in \mathcal{C}$.

Symmetry holds.

- Transitivity:

If $(x, y) \in \mathcal{C}$ and $(y, z) \in \mathcal{C}$, then $(x, z) \in \mathcal{C}$.

Assume that $(x, y) \in \mathcal{C}$, $(y, z) \in \mathcal{C}$.

$\cos(x) = \cos(y)$ and $\cos(y) = \cos(z)$

This implies that $\cos(x) = \cos(z)$ and $(x, z) \in \mathcal{C}$.

Transitivity holds.

Therefore, it is an equivalence relation. The equivalence classes are:

\rightarrow If $a \in \mathbb{R}$, the equivalence classes will be $[a] = \{x \in \mathbb{R} : \cos(a) = \cos(x)\}$

(b) The relation α on the domain $\mathbb{Z}^+ \times \mathbb{Z}^+$ is defined by $(m, n) \alpha (p, q)$ iff $mq = np$.

- Reflexive:

Let $n = m$ and $q = p$. Then $(m, m) \in \mathbb{Z}^+ \times \mathbb{Z}^+$ iff $mp = mp$

\therefore **Reflexivity holds.**

- Symmetric:

Let (m, n) and (p, q) are related then it implies (p, q) and (m, n) are related.

Assume that $(m, n) \alpha (p, q)$, this says that $mq = np$ and can be rewritten as $qm = pn$.

Based on comunitive properties, $qm = pn$ can be rewritten as well as $pn = mq$ which shows that **symmetry holds.**

- Transitive:

Let $(a, b), (c, d) \in \mathbb{Z}^+ \times \mathbb{Z}^+$ and $(c, d), (e, f) \in \mathbb{Z}^+ \times \mathbb{Z}^+$. This says that:

$(a, b), (c, d)$ is $(ad) = (bc)$ and $(c, d), (e, f)$ is $(cf) = (de)$

Multiplying the equations, we get: $(ad)(cf) = (bc)(de)$

Rearranging the values via comunitive properties: $(af)(cd) = (be)(cd)$

Divide by (cd) , we obtain $(af) = (be)$ and since $(a, b), (e, f) \in \mathbb{Z}^+ \times \mathbb{Z}^+$, this proves that the relation is **transitive.**

Therefore, it is an equivalence relation. The equivalence classes are:

\rightarrow If $a \in \mathbb{R}$, the equivalence classes will be $[m, n] = \{(p, q) \in \mathbb{Z}^+ \times \mathbb{Z}^+ : (mq) = (np)\}$