

Quantum Monte-Carlo Studies of Generalized Many-body Systems

Jørgen Høgberget

June 17, 2013

Contents

- 1 The basics of Quantum Monte-Carlo
- 2 Live Demo
- 3 Results and Discussions

Using the *projection operator* $\hat{\mathbf{P}}(\tau)$ on an arbitrary state $|\Psi_\tau\rangle$ projects the state onto the ground state of $\hat{\mathbf{H}}$ as $\tau \rightarrow \infty$.

Using the *projection operator* $\hat{\mathbf{P}}(\tau)$ on an arbitrary state $|\Psi_T\rangle$ projects the state onto the ground state of $\hat{\mathbf{H}}$ as $\tau \rightarrow \infty$.

$$\hat{\mathbf{P}}(\tau) |\Psi_T\rangle = \exp(-(\hat{\mathbf{H}} - E_0)\tau) |\Psi_T\rangle$$

Using the *projection operator* $\hat{\mathbf{P}}(\tau)$ on an arbitrary state $|\Psi_T\rangle$ projects the state onto the ground state of $\hat{\mathbf{H}}$ as $\tau \rightarrow \infty$.

$$\begin{aligned}\hat{\mathbf{P}}(\tau) |\Psi_T\rangle &= \exp(-(\hat{\mathbf{H}} - E_0)\tau) |\Psi_T\rangle \\ &= \exp(-(\hat{\mathbf{H}} - E_0)\tau) \sum_k C_k |\Psi_k\rangle\end{aligned}$$

Using the *projection operator* $\hat{\mathbf{P}}(\tau)$ on an arbitrary state $|\Psi_T\rangle$ projects the state onto the ground state of $\hat{\mathbf{H}}$ as $\tau \rightarrow \infty$.

$$\begin{aligned}\hat{\mathbf{P}}(\tau) |\Psi_T\rangle &= \exp(-(\hat{\mathbf{H}} - E_0)\tau) |\Psi_T\rangle \\ &= \exp(-(\hat{\mathbf{H}} - E_0)\tau) \sum_k C_k |\Psi_k\rangle \\ &= \sum_k C_k \exp(-(E_k - E_0)\tau) |\Psi_k\rangle\end{aligned}$$

Using the *projection operator* $\hat{\mathbf{P}}(\tau)$ on an arbitrary state $|\Psi_T\rangle$ projects the state onto the ground state of $\hat{\mathbf{H}}$ as $\tau \rightarrow \infty$.

$$\begin{aligned}\hat{\mathbf{P}}(\tau) |\Psi_T\rangle &= \exp(-(\hat{\mathbf{H}} - E_0)\tau) |\Psi_T\rangle \\ &= \exp(-(\hat{\mathbf{H}} - E_0)\tau) \sum_k C_k |\Psi_k\rangle \\ &= \sum_k C_k \exp(-(E_k - E_0)\tau) |\Psi_k\rangle \\ &= C_0 |\Psi_0\rangle + \sum_{k=1} C_k \exp(-\Delta E_k \tau) |\Psi_k\rangle ,\end{aligned}$$

where $\Delta E_k > 0$ and $C_k = \langle \Psi_k | \Psi_T \rangle$.

In other words

$$\lim_{\tau \rightarrow \infty} \langle \mathbf{r} | \hat{\mathbf{P}}(\tau) | \psi_T \rangle = \langle \psi_0 | \psi_T \rangle \psi_0(\mathbf{r}). \quad (1)$$

In other words

$$\lim_{\tau \rightarrow \infty} \langle \mathbf{r} | \hat{\mathbf{P}}(\tau) | \Psi_T \rangle = \langle \Psi_0 | \Psi_T \rangle \Psi_0(\mathbf{r}). \quad (1)$$

Idea I: Use an arbitrary *trial wave function* $\Psi_T(\mathbf{r})$ and perform the projection.

In other words

$$\lim_{\tau \rightarrow \infty} \langle \mathbf{r} | \hat{\mathbf{P}}(\tau) | \Psi_T \rangle = \langle \Psi_0 | \Psi_T \rangle \Psi_0(\mathbf{r}). \quad (1)$$

Idea I: Use an arbitrary *trial wave function* $\Psi_T(\mathbf{r})$ and perform the projection.

Problem: Requires a priori knowledge of the exact ground state energy E_0 .

In other words

$$\lim_{\tau \rightarrow \infty} \langle \mathbf{r} | \hat{\mathbf{P}}(\tau) | \Psi_T \rangle = \langle \Psi_0 | \Psi_T \rangle \Psi_0(\mathbf{r}). \quad (1)$$

Idea I: Use an arbitrary *trial wave function* $\Psi_T(\mathbf{r})$ and perform the projection.

Problem: Requires a priori knowledge of the exact ground state energy E_0 .

Solution: Introduce a *trial energy* $E_T(\tau)$. Will work as long as the trial energy drops below E_1 at a certain stage (and stays there).

In order to relate the projection process to a *Markov chain* Monte-Carlo process, τ needs to be split into sequential steps $\delta\tau$.

In order to relate the projection process to a *Markov chain* Monte-Carlo process, τ needs to be split into sequential steps $\delta\tau$.

This is achieved by using the following property of the projection operator

$$\begin{aligned}\hat{\mathbf{P}}(\tau + \delta\tau) &= \exp(-(\hat{\mathbf{H}} - E_T(\tau + \delta\tau))(\tau + \delta\tau)) \\ &= \exp(-(\hat{\mathbf{H}} - E_T(\tau + \delta\tau))\delta\tau) \\ &\quad \times \exp(-(\hat{\mathbf{H}} - E_T(\tau + \delta\tau))\tau) \\ &\simeq \exp(-(\hat{\mathbf{H}} - E_T(\tau + \delta\tau))\delta\tau)\hat{\mathbf{P}}(\tau),\end{aligned}$$

where the relation is approximate due to E_T not being constant.

Let $|\Phi(\tau)\rangle$ represent $\hat{\mathbf{P}}(\tau) |\Psi_T\rangle$. Using the relation from the previous slide reveals

Let $|\Phi(\tau)\rangle$ represent $\hat{\mathbf{P}}(\tau) |\Psi_T\rangle$. Using the relation from the previous slide reveals

$$\begin{aligned}\Phi(\mathbf{r}, \tau + \delta\tau) &= \langle \mathbf{r} | \Phi(\tau + \delta\tau) \rangle \\ &= \langle \mathbf{r} | \hat{\mathbf{P}}(\tau + \delta\tau) | \Psi_T \rangle\end{aligned}$$

Let $|\Phi(\tau)\rangle$ represent $\hat{\mathbf{P}}(\tau) |\Psi_T\rangle$. Using the relation from the previous slide reveals

$$\begin{aligned}\Phi(\mathbf{r}, \tau + \delta\tau) &= \langle \mathbf{r} | \Phi(\tau + \delta\tau) \rangle \\ &= \langle \mathbf{r} | \hat{\mathbf{P}}(\tau + \delta\tau) | \Psi_T \rangle \\ &\simeq \langle \mathbf{r} | \exp(-(\hat{\mathbf{H}} - E_T)\delta\tau) \hat{\mathbf{P}}(\tau) | \Psi_T \rangle\end{aligned}$$

Let $|\Phi(\tau)\rangle$ represent $\hat{\mathbf{P}}(\tau) |\Psi_T\rangle$. Using the relation from the previous slide reveals

$$\begin{aligned}\Phi(\mathbf{r}, \tau + \delta\tau) &= \langle \mathbf{r} | \Phi(\tau + \delta\tau) \rangle \\ &= \langle \mathbf{r} | \hat{\mathbf{P}}(\tau + \delta\tau) | \Psi_T \rangle \\ &\simeq \langle \mathbf{r} | \exp(-(\hat{\mathbf{H}} - E_T)\delta\tau) \hat{\mathbf{P}}(\tau) | \Psi_T \rangle \\ &= \langle \mathbf{r} | \exp(-(\hat{\mathbf{H}} - E_T)\delta\tau) | \Phi(\tau) \rangle\end{aligned}$$

Let $|\Phi(\tau)\rangle$ represent $\hat{\mathbf{P}}(\tau) |\Psi_T\rangle$. Using the relation from the previous slide reveals

$$\begin{aligned}\Phi(\mathbf{r}, \tau + \delta\tau) &= \langle \mathbf{r} | \Phi(\tau + \delta\tau) \rangle \\ &= \langle \mathbf{r} | \hat{\mathbf{P}}(\tau + \delta\tau) | \Psi_T \rangle \\ &\simeq \langle \mathbf{r} | \exp(-(\hat{\mathbf{H}} - E_T)\delta\tau) \hat{\mathbf{P}}(\tau) | \Psi_T \rangle \\ &= \langle \mathbf{r} | \exp(-(\hat{\mathbf{H}} - E_T)\delta\tau) | \Phi(\tau) \rangle \\ &= \int_{\mathbf{r}'} \langle \mathbf{r} | \exp(-(\hat{\mathbf{H}} - E_T)\delta\tau) | \mathbf{r}' \rangle \Phi(\mathbf{r}', \tau) d\mathbf{r}',\end{aligned}$$

Let $|\Phi(\tau)\rangle$ represent $\hat{\mathbf{P}}(\tau) |\Psi_T\rangle$. Using the relation from the previous slide reveals

$$\begin{aligned}\Phi(\mathbf{r}, \tau + \delta\tau) &= \langle \mathbf{r} | \Phi(\tau + \delta\tau) \rangle \\ &= \langle \mathbf{r} | \hat{\mathbf{P}}(\tau + \delta\tau) | \Psi_T \rangle \\ &\simeq \langle \mathbf{r} | \exp(-(\hat{\mathbf{H}} - E_T)\delta\tau) \hat{\mathbf{P}}(\tau) | \Psi_T \rangle \\ &= \langle \mathbf{r} | \exp(-(\hat{\mathbf{H}} - E_T)\delta\tau) | \Phi(\tau) \rangle \\ &= \int_{\mathbf{r}'} \langle \mathbf{r} | \exp(-(\hat{\mathbf{H}} - E_T)\delta\tau) | \mathbf{r}' \rangle \Phi(\mathbf{r}', \tau) d\mathbf{r}',\end{aligned}$$

where $\langle \mathbf{r} | \exp(-(\hat{\mathbf{H}} - E_T)\delta\tau) | \mathbf{r}' \rangle \equiv G(\mathbf{r}', \mathbf{r}; \delta\tau)$ is a *Green's function* interpreted as the transition probability between \mathbf{r} and \mathbf{r}' .

Idea: In order to relate the Green's function to well known Markov processes, the exponential is split

$$\begin{aligned}\exp(-(\hat{\mathbf{H}} - E_T)\delta\tau) &= \exp\left(\frac{1}{2}\nabla^2\delta\tau - (\hat{\mathbf{V}} - E_T)\delta\tau\right) \\ &= \exp\left(\frac{1}{2}\nabla^2\delta\tau\right) \exp(-(\hat{\mathbf{V}} - E_T)\delta\tau) \\ &\quad + \mathcal{O}(\delta\tau^2),\end{aligned}$$

where the kinetic part describes a diffusion process with diffusion constant $D = \frac{1}{2}$, and the potential part describes a weighting (linear in position space). This is referred to as the *short time approximation*.

Problem: The Green's function has singular points in the Coulomb interaction.

Problem: The Green's function has singular points in the Coulomb interaction.

Solution: By evolving $f(\mathbf{r}, \tau) = \Phi(\mathbf{r}, \tau)\Psi_T(\mathbf{r})$ instead of $\Phi(\mathbf{r}, \tau)$ alone, the singularities are *implicitly* taken care of.

Originally:

$$\frac{\partial \Phi(\mathbf{r}, \tau)}{\partial \tau} = \left[\frac{1}{2} \nabla^2 - \left(\hat{\mathbf{V}} - E_T \right) \right] \Phi(\mathbf{r}, \tau).$$

Originally:

$$\frac{\partial \Phi(\mathbf{r}, \tau)}{\partial \tau} = \left[\frac{1}{2} \nabla^2 - (\hat{\mathbf{V}} - E_T) \right] \Phi(\mathbf{r}, \tau).$$

Importance sampled:

$$\frac{\partial f(\mathbf{r}, \tau)}{\partial \tau} = \left[\frac{1}{2} \nabla \cdot (\nabla - \mathbf{F}(\mathbf{r})) - (E_L(\mathbf{r}) - E_T) \right] f(\mathbf{r}, \tau), \quad (2)$$

where

$$\mathbf{F}(\mathbf{r}) = 2\Psi_T(\mathbf{r})^{-1} \nabla \Psi_T(\mathbf{r}) \quad (3)$$

is the *quantum force* and

$$E_L(\mathbf{r}) = \Psi_T(\mathbf{r})^{-1} \hat{\mathbf{H}} \Psi_T(\mathbf{r}) \quad (4)$$

is the *local energy*.

The Green's functions have closed form solutions on the form

$$G_{\text{Diff}}(\mathbf{r}', \mathbf{r}; \delta\tau) \propto \exp\left(-|\mathbf{r} - \mathbf{r}' - D\delta\tau\mathbf{F}(\mathbf{r})|^2 / 4D\delta\tau\right),$$
$$G_{\text{B}}(\mathbf{r}', \mathbf{r}; \delta\tau) \propto \exp\left(-\left(\frac{1}{2}[E_L(\mathbf{r}) + E_L(\mathbf{r}')] - E_T\right)\delta\tau\right),$$

where B denotes *branching*.

The process of projection is modelled by letting an ensemble of walkers span $f(\mathbf{r}, \tau)$. An iteration involves diffusing all the walkers and distributing new weights.

Branching

Idea: The weights are modelled by spawning and killing walkers with a rate equal to G_B .

Branching

Idea: The weights are modelled by spawning and killing walkers with a rate equal to G_B .

Problem: G_B is not necessarily an integer.

Branching

Idea: The weights are modelled by spawning and killing walkers with a rate equal to G_B .

Problem: G_B is not necessarily an integer.

Solution: Create $\text{floor}(G_B)$ copies with a $G_B - \text{floor}(G_B)$ chance of creating an additional copy.

Branching

Idea: The weights are modelled by spawning and killing walkers with a rate equal to G_B .

Problem: G_B is not necessarily an integer.

Solution: Create $\text{floor}(G_B)$ copies with a $G_B - \text{floor}(G_B)$ chance of creating an additional copy.

Or more compact: Create

$$\overline{G}_B = \text{floor}(G_B + a) \quad (5)$$

copies, where $a \in [0, 1)$ is a uniformly distributed random number. If the value is zero, the walker dies.

Branching




Figure: Branching illustrated. The integer values represent \bar{G}_B .

Diffusion

The diffusion equation representing G_{Diff} is the so-called *Fokker-Planck equation*.

Diffusion

The diffusion equation representing G_{Diff} is the so-called *Fokker-Planch equation*.

Anisotropic diffusion process. The quantum force pushes the walkers into regions of higher probability.

Diffusion

The diffusion equation representing G_{Diff} is the so-called *Fokker-Planch equation*.

Anisotropic diffusion process. The quantum force pushes the walkers into regions of higher probability.

According to the Fokker-Planch formalism, a new position \mathbf{r}_{i+1} is calculated from the old one, \mathbf{r}_i , as follows

$$\mathbf{r}_{i+1} = \mathbf{r}_i + D\delta\tau\mathbf{F}(\mathbf{r}_i) + \xi, \quad (6)$$

where ξ is a vector of normal distributed random numbers with variance $\sqrt{2D\delta\tau}$.

Diffusion

Problem: Due to a finite time step, the previous equations do not guarantee convergence.

Diffusion

Problem: Due to a finite time step, the previous equations do not guarantee convergence.

Solution: The *Metropolis algorithm* will correct this bias:

$$A(i \rightarrow j) = \min\{R_G(i \rightarrow j)R_\psi(i \rightarrow j)^2, 1\}, \quad (7)$$

where $i \rightarrow j$ denotes a transition from state i to state j , A is the probability of accepting the transition,

$$R_G(i \rightarrow j) = G_{\text{diff}}(\mathbf{r}_i, \mathbf{r}_j; \delta\tau) / G_{\text{diff}}(\mathbf{r}_j, \mathbf{r}_i; \delta\tau),$$

and

$$R_\psi(i \rightarrow j) = |\Psi_T(\mathbf{r}_i) / \Psi_T(\mathbf{r}_j)|.$$

End of section.

hei

hei