

Given a 6x6 2D Array, arr :

```
1 1 1 0 0 0
0 1 0 0 0 0
1 1 1 0 0 0
0 0 0 0 0 0
0 0 0 0 0 0
0 0 0 0 0 0
```

We define an hourglass in A to be a subset of values with indices falling in this pattern in 's graphical representation:

```
a b c
  d
e f g
```

There are 16 hourglasses in arr, and an hourglass sum is the sum of an hourglass' values. Calculate the hourglass sum for every hourglass in arr, then print the maximum hourglass sum.

For example, given the 2D array:

```
-9 -9 -9 1 1 1
0 -9 0 4 3 2
-9 -9 -9 1 2 3
0 0 8 6 6 0
0 0 0 -2 0 0
0 0 1 2 4 0
```

We calculate the following 16 hourglass values:

```
-63, -34, -9, 12,
-10, 0, 28, 23,
-27, -11, -2, 10,
9, 17, 25, 18
```

Our highest hourglass value is 28 from the hourglass:

```
0 4 3
  1
8 6 6
```

Note: If you have already solved the Java domain's Java 2D Array challenge, you may wish to skip this challenge.

Function Description

Complete the function `hourglassSum` in the editor below. It should return an integer, the maximum hourglass sum in the array.

`hourglassSum` has the following parameter(s):

- `arr`: an array of integers

Input Format

Each of the lines of inputs contains space-separated integers .

Constraints

- $-9 \leq arr[i][j] \leq 9$
- $0 \leq i, j \leq 5$

Output Format

Print the largest (maximum) hourglass sum found in .

Sample Input

```
1 1 1 0 0 0
0 1 0 0 0 0
1 1 1 0 0 0
0 0 2 4 4 0
0 0 0 2 0 0
0 0 1 2 4 0
```

Sample Output

19

Explanation

arr contains the following hourglasses:

```
1 1 1 1 1 0 1 0 0 0 0 0
 1      0      0      0
1 1 1 1 1 0 1 0 0 0 0 0

0 1 0 1 0 0 0 0 0 0 0 0
 1      1      0      0
0 0 2 0 2 4 2 4 4 4 4 0

1 1 1 1 1 0 1 0 0 0 0 0
 0      2      4      4
0 0 0 0 0 2 0 2 0 2 0 0

0 0 2 0 2 4 2 4 4 4 4 0
 0      0      2      0
0 0 1 0 1 2 1 2 4 2 4 0
```

The hourglass with the maximum sum (19) is:

2 4 4

2

1 2 4