


112 families were paired-matched; 114 children

56 pairs
+ 1 twin
+ 1 sibling



Abandoned study=4
Died before age 5=2
Developmentally delayed=2

Died before age 5=2
Withdrew before age 5=1

Replacement=7

Crossover=1
Non-randomly reassigned=2

Replacement=1

[R=1;D=0]=7
[R=1;D=1]=50

Non-compliance=3

[R=0;D=1]=3
[R=0;D=0]=51

Participated in ABC Program [D=1]
N=56 children

Not participated in ABC Program [D=0]
N=54 children

Not=3
Followed

Not=3
Followed

School-Age Randomization [SR]

[R=1;D=1;SR=1]=25
[R=1;D=1;SR=No]=1

[R=0;D=0;SR=1]=24

[R=0;D=0;SR=0]=23

[R=0;D=0;SR=No]=4

Temporary Attrited=5

Temporary Attrited=4

School-Age Treatment Group [SR=1]
N=49 children

School-Age Control Group [SR=0]
N=47 children

No Show=3

Participated
N=46 children

Not Participated
N=47 children

Attrited=1

Adult Follow-Up (Age 21)

[R=1; SR=1]=25

[R=0; SR=1]=24

[R=1; SR=0]=19

[R=0; SR=0]=27

[R=1; SR=No]=4