

§7.7: Approximate Integration

Ch 7: Techniques of Integration Math 5B: Calculus II

Dr. Jorge Eduardo Basilio

Department of Mathematics & Computer Science
Pasadena City College

Class #12 Notes

April 2, 2019
Spring 2019

- 1 Guiding Questions
- 2 Review of definitions of Integration
- 3 Endpoint Approximations
- 4 Midpoint Approximations
- 5 Trapezoid Approximations

Guiding Question(s)

- 1 What are some common techniques to **approximate integrals numerically**?

Guiding Question(s)

- ① What are some common techniques to **approximate integrals numerically**?
- ② What are **error bounds** for the midpoint and trapezoid rules

Guiding Question(s)

- ① What are some common techniques to **approximate integrals numerically**?
- ② What are **error bounds** for the midpoint and trapezoid rules
- ③ How can we use them to anticipate the number of sums needed to achieve a certain accuracy?

- Definition of **anti-derivative**. Given a function $f(x)$, we say $F(x)$ is an **anti-derivative** of $f(x)$ if

$$F'(x) = f(x). \quad \text{Notation: } F(x) + C = \int f(x) dx.$$

- Definition of **anti-derivative**. Given a function $f(x)$, we say $F(x)$ is an **anti-derivative** of $f(x)$ if

$$F'(x) = f(x). \quad \text{Notation: } F(x) + C = \int f(x) dx.$$

so we could find anti-derivatives $F(x)$ using our knowledge of derivatives and “guessing” $F(x)$.

Review Integration

- Definition of **anti-derivative**. Given a function $f(x)$, we say $F(x)$ is an **anti-derivative** of $f(x)$ if

$$F'(x) = f(x). \quad \text{Notation: } F(x) + C = \int f(x) dx.$$

so we could find anti-derivatives $F(x)$ using our knowledge of derivatives and “guessing” $F(x)$.

- Definition of **definite integral**.

This is way more complicated. Notation: $\int_a^b f(x) dx$.

Review Integration

- Definition of **anti-derivative**. Given a function $f(x)$, we say $F(x)$ is an **anti-derivative** of $f(x)$ if

$$F'(x) = f(x). \quad \text{Notation: } F(x) + C = \int f(x) dx.$$

so we could find anti-derivatives $F(x)$ using our knowledge of derivatives and “guessing” $F(x)$.

- Definition of **definite integral**.

This is way more complicated. Notation: $\int_a^b f(x) dx$.

- **Important point**: though the notation is similar, they are very different.

$$\int f(x) dx = \text{infinitely many functions} \quad \text{and} \quad \int_a^b f(x) dx = \text{a single number}$$

- Definition of **definite integral**. This is way more complicated.

- Definition of **definite integral**. This is way more complicated.

First, we cut up the interval $[a, b]$ into n pieces and denote the endpoints of the subintervals $[x_{i-1}, x_i]$ with a **partition**

$$a = x_0 < x_1 < x_2 < \cdots < x_{i-1} < x_i < x_{i+1} < \cdots < x_n = b.$$

- Definition of **definite integral**. This is way more complicated.

First, we cut up the interval $[a, b]$ into n pieces and denote the endpoints of the subintervals $[x_{i-1}, x_i]$ with a **partition**

$$a = x_0 < x_1 < x_2 < \cdots < x_{i-1} < x_i < x_{i+1} < \cdots < x_n = b.$$

Pick a random sample of $c_i \in [x_{i-1}, x_i]$ and compute the area of the i th sub-rectangle: Area (one rectangle) = $f(c_i)\Delta x_i$ where $\Delta x_i = x_i - x_{i-1}$

- Definition of **definite integral**. This is way more complicated.

First, we cut up the interval $[a, b]$ into n pieces and denote the endpoints of the subintervals $[x_{i-1}, x_i]$ with a **partition**

$$a = x_0 < x_1 < x_2 < \cdots < x_{i-1} < x_i < x_{i+1} < \cdots < x_n = b.$$

Pick a random sample of $c_i \in [x_{i-1}, x_i]$ and compute the area of the i th sub-rectangle: Area (one rectangle) = $f(c_i)\Delta x_i$ where $\Delta x_i = x_i - x_{i-1}$

Approximate area under the graph with n rectangles:

$$S_n = \sum_{i=1}^n f(c_i) \Delta x_i$$

- Definition of **definite integral**. This is way more complicated.

First, we cut up the interval $[a, b]$ into n pieces and denote the endpoints of the subintervals $[x_{i-1}, x_i]$ with a **partition**

$$a = x_0 < x_1 < x_2 < \cdots < x_{i-1} < x_i < x_{i+1} < \cdots < x_n = b.$$

Pick a random sample of $c_i \in [x_{i-1}, x_i]$ and compute the area of the i th sub-rectangle: Area (one rectangle) = $f(c_i)\Delta x_i$ where $\Delta x_i = x_i - x_{i-1}$

Approximate area under the graph with n rectangles:

$$S_n = \sum_{i=1}^n f(c_i) \Delta x_i$$

To get the **exact** area we take a limit:

- Definition of **definite integral**. This is way more complicated.

First, we cut up the interval $[a, b]$ into n pieces and denote the endpoints of the subintervals $[x_{i-1}, x_i]$ with a **partition**

$$a = x_0 < x_1 < x_2 < \cdots < x_{i-1} < x_i < x_{i+1} < \cdots < x_n = b.$$

Pick a random sample of $c_i \in [x_{i-1}, x_i]$ and compute the area of the i th sub-rectangle: Area (one rectangle) = $f(c_i)\Delta x_i$ where $\Delta x_i = x_i - x_{i-1}$

Approximate area under the graph with n rectangles:

$$S_n = \sum_{i=1}^n f(x_i) \Delta x_i$$

To get the **exact** area we take a limit:

$$\int_a^b f(x) dx = \lim_{n \rightarrow \infty} S_n = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i) \Delta x_i$$

Review Integration

- This is hard and alot of work! But don't ever forget that **definite integrals are (infinite) sums which we can always approximate**

$$\int_a^b f(x) dx \approx S_n.$$

Review Integration

- This is hard and a lot of work! But don't ever forget that **definite integrals are (infinite) sums which we can always approximate**

$$\int_a^b f(x) dx \approx S_n.$$

- The **Fundamental Theorem of Calculus** is a huge short-cut!

$$\int_a^b f(x) dx = F(x)|_a^b = F(b) - F(a)$$

provided we can find an anti-derivative $F(x)$.

Review Integration

- This is hard and a lot of work! But don't ever forget that **definite integrals** are (infinite) sums which we can always approximate

$$\int_a^b f(x) dx \approx S_n.$$

- The **Fundamental Theorem of Calculus** is a huge short-cut!

$$\int_a^b f(x) dx = F(x)|_a^b = F(b) - F(a)$$

provided we can find an anti-derivative $F(x)$.

- At this point, you now have many tools in your **integration toolbox** to compute a lot of integrals. But...

Review Integration

- This is hard and a lot of work! But don't ever forget that **definite integrals are (infinite) sums which we can always approximate**

$$\int_a^b f(x) dx \approx S_n.$$

- The **Fundamental Theorem of Calculus** is a huge short-cut!

$$\int_a^b f(x) dx = F(x)|_a^b = F(b) - F(a)$$

provided we can find an anti-derivative $F(x)$.

- At this point, you now have many tools in your **integration toolbox** to compute a lot of integrals. But...
- There's still many, many integrals we can't find anti-derivatives of!

Review Integration

- This is hard and a lot of work! But don't ever forget that **definite integrals are (infinite) sums which we can always approximate**

$$\int_a^b f(x) dx \approx S_n.$$

- The **Fundamental Theorem of Calculus** is a huge short-cut!

$$\int_a^b f(x) dx = F(x)|_a^b = F(b) - F(a)$$

provided we can find an anti-derivative $F(x)$.

- At this point, you now have many tools in your **integration toolbox** to compute a lot of integrals. But...
- There's still many, many integrals we can't find anti-derivatives of!
- So we will go back to the definition of the definite integral and settle for approximation.

Approximate Integration Techniques

Essentially, we have many choices for how to pick $c_i \in [x_{i-1}, x_i]$
(Note: the book uses x_i^* instead of c_i).

- **Left-endpoint approximation (LEA):** $c_i = x_{i-1}$.

Ex: when f is increasing, this **underestimates** the exact area.

- **Right-endpoint approximation (REA):** $c_i = x_i$.

Ex: when f is increasing, this **overestimates** the exact area.

- **Midpoint approximation (MPA):** $c_i = \frac{x_{i-1} + x_i}{2}$.

this is much better than the LEA and REA, in general, since it averages the them!

In my opinion, it's a bit silly to write out the full formulas for these approximations. If you know the idea, then you can just find these values by hand with patience.

Endpoint Approximations

Set up for all approximations: n a positive integer. $a, b \in \mathbb{R}$. for $i = 0, 1, 2, \dots, n$:

$$\Delta x = \frac{b-a}{n}$$

and

$$x_i = a + i \cdot \Delta x$$

and

subintervals $[x_{i-1}, x_i]$

Definition 1: Endpoint Approximations

- Left Endpoint approximation (LEA):

$$\int_a^b f(x) dx \approx \Delta x [f(x_0) + f(x_1) + f(x_2) + \cdots + f(x_{n-1})] = S_n(LEA)$$

- Right Endpoint approximation (REA):

$$\int_a^b f(x) dx \approx \Delta x [f(x_1) + f(x_2) + f(x_3) + \cdots + f(x_n)] = S_n(REA)$$

Midpoint Approximations

Set up for all approximations: n a positive integer. $a, b \in \mathbb{R}$. for $i = 0, 1, 2, \dots, n$:

$$\Delta x = \frac{b - a}{n}$$

and

$$x_i = a + i \cdot \Delta x$$

and

subintervals $[x_{i-1}, x_i]$

Definition 2: Midpoint Approximations

- **Midpoint approximation (MPA):** if we average the endpoints of the subintervals, we get the midpoints, given by

$$\bar{x}_i = \frac{x_{i-1} + x_i}{2} = a + i \cdot (\Delta x / 2). \text{ Then}$$

$$\int_a^b f(x) dx \approx \Delta x [f(\bar{x}_1) + f(\bar{x}_2) + f(\bar{x}_3) + \cdots + f(\bar{x}_n)] = S_n(MPA)$$

Trapezoid Approximations

Set up for all approximations: n a positive integer. $a, b \in \mathbb{R}$. for $i = 0, 1, 2, \dots, n$:

$$\Delta x = \frac{b - a}{n}$$

and

$$x_i = a + i \cdot \Delta x$$

and

subintervals $[x_{i-1}, x_i]$

Definition 3: Trapezoid Approximations

- **Midpoint approximation (MPA):** if we average the **height** of the function at the endpoints of the subintervals, then we get trapezoids.

$$\int_a^b f(x) dx \approx \Delta x [f(x_0) + 2f(x_1) + \dots + 2f(x_{n-1}) + f(x_n)] = S_n(\text{TrapA})$$

Note: $\text{Areas}(\text{Trap}) = \frac{f(x_{i-1}) + f(x_i)}{2} \Delta x$