

## §7.8: Improper Integrals

### Ch 7: Techniques of Integration

### Math 5B: Calculus II

Dr. Jorge Eduardo Basilio

Department of Mathematics & Computer Science  
Pasadena City College

**Class #14 Notes**

April 9, 2019  
Spring 2019

- 1 Guiding Questions
- 2 Introduction
- 3 Improper Integrals: Type I
- 4  $p$ -Test Type I
- 5 Improper Integrals: Type II
- 6  $p$ -Test Type II
- 7 Improper Integrals: Comparison Test

# Guiding Questions for §7.8

## Guiding Question(s)

- 1 Can we extend definite integrals,  $\int_a^b f(x) dx$ , to more general intervals?
- 2 How can we deal with integrals with an **infinite interval of integration**?
- 3 How can we deal with integrals where  $f(x)$  has an **infinite discontinuity** inside the interval of integration?


- What's wrong with the following argument?

$$\begin{aligned}\int_{-1}^1 \frac{1}{x^2} dx &= \left[ \frac{x^{-1}}{-1} \right]_{-1}^1 \\ &= -1 + -1 \\ &= -2\end{aligned}$$


But, we know that  $f(x) = \frac{1}{x^2}$  is always positive, and the integral is the area under the curve. What went wrong?

- What about:

$$\begin{aligned}\int_1^\infty \frac{1}{x^2} dx &= \left[ \frac{x^{-1}}{-1} \right]_1^\infty \\ &= \frac{-1}{\infty} + 1 \\ &= 1.\end{aligned}$$



Graph of  $f(x) = \frac{1}{x^2}$  over  $[-1, 1]$



Graph of  $f(x) = \frac{1}{x^2}$  over  $[1, 10]$

- The integral value  $\int_1^\infty \frac{1}{x^2} dx = 1$  seems reasonable.
- We can make it rigorous by using limits and setting:

$$\int_1^\infty \frac{1}{x^2} dx = \lim_{R \rightarrow \infty} \int_1^R \frac{1}{x^2} dx$$

- Since we can evaluate as before:

$$\begin{aligned} &= \lim_{R \rightarrow \infty} \left[ \frac{x^{-1}}{-1} \right]_1^R \\ &= \lim_{R \rightarrow \infty} \left[ \frac{-1}{R} + 1 \right] \\ &= 1 \end{aligned}$$

# Improper Integrals: Type I

## Definition 1: Improper Integrals: Type I

- Assume  $\int_a^b f(x) dx$  exists for all  $b \geq a$ . We define  $\int_a^\infty f(x) dx$  to be the limit (when it exists):

$$\int_a^\infty f(x) dx = \lim_{R \rightarrow \infty} \int_a^R f(x) dx$$

- Similarly, assume  $\int_a^b f(x) dx$  exists for all  $a \leq b$ . We define  $\int_{-\infty}^b f(x) dx$  to be the limit (when it exists):

$$\int_{-\infty}^b f(x) dx = \lim_{R \rightarrow -\infty} \int_R^b f(x) dx$$

- These are called **improper integrals of type I**.

# Improper Integrals: Type I

## Definition 2: Improper Integrals: Type I

Given improper integrals:  $\int_a^\infty f(x) dx$  or  $\int_{-\infty}^b f(x) dx$

- **Converge:** if the limit exists & is a finite number. We say that the improper integral **converges**.
- **Diverge:** if the limit does not exist OR is not a finite number. We say that the improper integral **diverges**.

When both  $\int_a^\infty f(x) dx$  and  $\int_{-\infty}^b f(x) dx$  converge,

- We define  $\int_{-\infty}^\infty f(x) dx$  to be:

$$\int_{-\infty}^\infty f(x) dx = \int_{-\infty}^c f(x) dx + \int_c^\infty f(x) dx$$

Where  $c$  can be any real number. Usually, we pick  $c = 0$  out of convenience.



# Improper Integrals: Type I

## Activity 1:

C or D? That is, do the following improper integrals converge or diverge? If they converge, find their value.

(a)  $\int_1^{\infty} \frac{1}{x} dx$

(b)  $\int_{-\infty}^1 xe^x dx$

(c)  $\int_{-\infty}^{\infty} \frac{1}{1+x^2} dx$

[Outline](#)

[Guiding Questions](#)

[Intro](#)

[Type I](#)

[p-Test Type I](#)

[Type II](#)

[p-Test Type II](#)

[Comparison Test](#)

# Improper Integrals: Type I

§7.8

Dr. Basilio

[Outline](#)

[Guiding  
Questions](#)

[Intro](#)

[Type I](#)

[p-Test Type I](#)

[Type II](#)

[p-Test Type II](#)

[Comparison  
Test](#)

# Improper Integrals: Type I

§7.8

Dr. Basilio

[Outline](#)

[Guiding  
Questions](#)

[Intro](#)

[Type I](#)

[p-Test Type I](#)

[Type II](#)

[p-Test Type II](#)

[Comparison  
Test](#)

# Improper Integrals: Type I

## Activity 2:

Investigate numerically using Sage whether  $\int_1^{\infty} e^{-x^2} dx$  converges or diverges. If it appears to converge, estimate its value.

An important test for convergence and divergence is the following:

## Theorem 1: $p$ -test Type I

- $$\int_1^{\infty} \frac{1}{x^p} dx = \begin{cases} \text{converges,} & \text{if } p > 1 \\ \text{diverges,} & \text{if } p \leq 1 \end{cases}$$
- In fact, when  $p > 1$ ,  $\int_1^{\infty} \frac{1}{x^p} dx = \frac{1}{p-1}$

# Improper Integrals: $p$ -Test

## Proof:

Proof breaks into several steps:

- $p = 1$  we showed earlier that it diverges
- Compute  $\int_1^R x^{-p} dx$  for any  $R \geq 1$ .
- Case:  $p > 1$
- Case  $p < 1$

# Improper Integrals: $p$ -Test

§7.8

Dr. Basilio

[Outline](#)

[Guiding  
Questions](#)

[Intro](#)

[Type I](#)

[\$p\$ -Test Type I](#)

[Type II](#)

[\$p\$ -Test Type II](#)

[Comparison  
Test](#)

# Improper Integrals: Type II

## Definition 3: Improper Integrals: Type II

Assume  $f$  has a **vertical asymptote** at  $x = c$ . We sometimes call this a “infinite discontinuity” at  $x = c$  but otherwise is continuous on the interval  $(a, b)$  with  $a < c < b$ . We define:

- $$\int_a^c f(x) dx = \lim_{b \rightarrow c^-} \int_a^b f(x) dx$$

- $$\int_c^b f(x) dx = \lim_{a \rightarrow c^+} \int_a^b f(x) dx$$

- When both integrals above converge, we define

$$\int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx$$

- These are called **improper integrals of type II**.

[Outline](#)

[Guiding Questions](#)

[Intro](#)

[Type I](#)

[p-Test Type I](#)

[Type II](#)

[p-Test Type II](#)

[Comparison Test](#)



## Activity 3:

C or D? That is, do the following improper integrals converge or diverge? If they converge, find their value.

(a)  $\int_0^9 \frac{1}{\sqrt{x}} dx$

(b)  $\int_0^1 \frac{1}{x} dx$

(c)  $\int_0^2 \frac{1}{\sqrt[3]{(x-1)^2}} dx$

[Outline](#)

[Guiding Questions](#)

[Intro](#)

[Type I](#)

[p-Test Type I](#)

[Type II](#)

[p-Test Type II](#)

[Comparison Test](#)

# Improper Integrals: Type II

## Activity 4:

*C* or *D*? That is, do the following improper integrals converge or diverge? If they converge, find their value.

(a)  $\int_0^1 \frac{1}{\sqrt{1-x^2}} dx$

(b)  $\int_0^{\pi/2} \sec(x) dx$

# Improper Integrals: Type II

# Improper Integrals: $p$ -Test Type II

An important test for convergence and divergence is the following:

## Theorem 2: $p$ -test Type II

- $$\int_0^1 \frac{1}{x^p} dx = \begin{cases} \text{diverges,} & \text{if } p \geq 1 \\ \text{converges,} & \text{if } p < 1 \end{cases}$$
- In fact, when  $p < 1$ ,  $\int_0^1 \frac{1}{x^p} dx = \frac{1}{p-1}$

**NOTICE:** that this is similar to the  $p$ -test for type I integrals but the roles switch!

# Improper Integrals: Comparison Test

Many times the integral is either impossible to evaluate directly or too difficult. The following test gives us an easy way to determine convergence or divergence. Note it does not give us a value of the integral if it converges (but it does give us an estimate, however).

## Theorem 3: Comparison Test

Assume that  $f$  and  $g$  are continuous on  $[a, \infty)$ .

Assume that  $f(x) \geq g(x) \geq 0$ . Then

- (a) If  $\int_a^\infty f(x) dx$  converges, THEN  $\int_a^\infty g(x) dx$  also converges.
- (b) If  $\int_a^\infty g(x) dx$  diverges, THEN  $\int_a^\infty f(x) dx$  also diverges.

# Improper Integrals: Comparison Test

## Activity 5:

Use the comparison test to show that  $\int_{-\infty}^{\infty} e^{-x^2} dx$  converges.

(Hint: split into three integrals:  $\int_{-\infty}^{-1} + \int_{-1}^1 + \int_1^{\infty}$ . Use the comparison test with  $e^{-x^2} \leq e^{-x}$ .)

# Improper Integrals: Comparison Test

## Activity 6:

Use the comparison test to determine whether the integral converges or diverges.

(a)  $\int_1^{\infty} \frac{1}{\sqrt{x^3 + 1}} dx$

(b)  $\int_1^{\infty} \frac{1}{e^{3x} + \sqrt{x}} dx$

(c)  $\int_1^{\infty} \frac{1 + \sin^2(x)}{\sqrt{x}} dx$

[Outline](#)

[Guiding Questions](#)

[Intro](#)

[Type I](#)

[p-Test Type I](#)

[Type II](#)

[p-Test Type II](#)

[Comparison Test](#)



# Improper Integrals: Comparison Test

§7.8

Dr. Basilio

[Outline](#)

[Guiding  
Questions](#)

[Intro](#)

[Type I](#)

[p-Test Type I](#)

[Type II](#)

[p-Test Type II](#)

[Comparison  
Test](#)