

# Solving the Many-Electron Schrödinger Equation

## With a Transformer Architecture

# Outline

- 1 The problem
- 2 Finding the solution
- 3 Challenges and opportunities

## Setup: distances, spins, coordinates

- System:  $N$  electrons, nuclei with charge  $Z_I$  at fixed  $\mathbf{R}_I$ .
- Electron coordinates  $\mathbf{r}_i \in \mathbb{R}^3$ , spin  $\sigma_i \in \{\uparrow, \downarrow\}$ .
- Atomic units (distances in Bohr, energies in Hartree).
- Distances:  $r_{ij} = |\mathbf{r}_i - \mathbf{r}_j|$ ,  $r_{iI} = |\mathbf{r}_i - \mathbf{R}_I|$ .

# The equation (TISE Hamiltonian)

- Kinetic energy:  $T = -\frac{1}{2} \sum_{i=1}^N \nabla_i^2$ .
- Electron-nuclear attraction:  $V_{en} = - \sum_{i,I} \frac{Z_I}{r_{iI}}$ .
- Electron-electron repulsion:  $V_{ee} = \sum_{i<j} \frac{1}{r_{ij}}$ .
- Hamiltonian:  $H = T + V_{en} + V_{ee}$ .
- Solve the time-independent Schrödinger equation:  $H \Psi = E \Psi$ .

# Fermi–Dirac statistics & antisymmetry

- Electrons are indistinguishable fermions.
- Exchanging two electrons flips the wavefunction's sign:  
$$\Psi(\dots i, j \dots) = -\Psi(\dots j, i \dots).$$
- Pauli exclusion: no two electrons can occupy the same state.
- Use a Slater determinant to enforce an antisymmetric  $\Psi$ .

# Exponential decay of wavefunction

- Bound-state wavefunctions decay exponentially as  $r \rightarrow \infty$ .
- Example: hydrogen ground state  $\psi(r) \sim e^{-r}$  (in a.u.).
- Ansätze include an exponential envelope so that  $\Psi \rightarrow 0$  as  $r \rightarrow \infty$ .

# Kato cusp conditions

- Coulomb potentials cause a sharp cusp in  $\Psi$  when particles coalesce.
- Electron–nucleus cusp:  $\frac{\partial \Psi}{\partial r_{iI}} \Big|_{r_{iI}=0} = -Z_I \Psi(0).$
- Electron–electron cusp:  $\frac{\partial \Psi}{\partial r_{ij}} \Big|_{r_{ij}=0} = \frac{1}{2} \Psi(0).$

# The Psiformer Ansatz (overview)

- Neural network trial wavefunction (“Wavefunction Transformer”).
- Uses self-attention layers to model electron correlations.
- Permutation equivariant (independent of electron ordering).
- Greatly improves accuracy, especially for larger systems.
- Solves Schrödinger equation from first principles (no external data).



# Dissecting the Ansatz

- Slater–Jastrow form:  $\Psi = \left( \sum_d c_d \det[\phi_k^d(\mathbf{r}_i)] \right) \exp(J)$ .
- Slater determinant part ensures the correct antisymmetric exchange.
- One-electron orbitals  $\phi_k^d$  are learned functions of all electron coordinates.
- Jastrow factor  $J$  enforces electron-electron cusp conditions.
- Additional envelope functions ensure exponential decay at long range.

# Jastrow factor form (parallel/antiparallel spins)

- Jastrow factor:  $J = \sum_{i < j} u_{\sigma_i, \sigma_j}(r_{ij})$ .
- Use separate  $u(r)$  for same-spin vs opposite-spin pairs.
- Example choice:  $u_{\uparrow\downarrow}(r) = \frac{\alpha r}{1 + \beta r}$  (two parameters).
- Satisfies cusp:  $u(r) \approx \frac{1}{2}r$  as  $r \rightarrow 0$ .
- Jastrow adds correlation beyond antisymmetry (lowers the energy).

# Loss: Rayleigh quotient

- Variational principle uses the energy expectation as loss.
- $E[\Psi] = \frac{\langle \Psi | H | \Psi \rangle}{\langle \Psi | \Psi \rangle} \geq E_0$  (Rayleigh quotient).
- Minimizing  $E[\Psi]$  drives the ansatz toward the ground state.
- $E_0$  is the true ground energy (minimum possible).

# Metropolis–Hastings sampling

- Markov Chain Monte Carlo algorithm to sample  $|\Psi(\mathbf{r})|^2$ .
- Propose random moves for electron coordinates.
- Accept move with probability  $A = \min\left(1, \frac{|\Psi_{\text{new}}|^2}{|\Psi_{\text{old}}|^2}\right)$ .
- After thermalization, obtained samples follow the target distribution  $|\Psi|^2$ .

# Variational Monte Carlo loop

- Initialize neural network parameters (trial  $\Psi$ ).
- Sample electron configurations via Metropolis–Hastings.
- Evaluate energy and gradient from the sampled configurations.
- Update parameters to lower the energy (gradient descent).
- Iterate until convergence to the minimum energy (ground state).

# Optimizer: KFAC (natural gradient)

- Kronecker-Factored Approximate Curvature (KFAC) optimizer.
- Approximates the natural gradient using a factored curvature matrix.
- Speeds up training and improves stability for large networks.
- Used to efficiently optimize FermiNet and Psiformer wavefunctions.

# FermiNet baseline

- FermiNet: first deep-neural-network wavefunction ansatz (Pfau et al., 2020).
- Architecture: multiple dense layers with electron-wise feature streams.
- Outputs single-electron orbitals feeding into a Slater determinant.
- Achieved high accuracy on small molecules (near chemical accuracy).
- Faced scaling limits: accuracy and efficiency drop for larger systems.

# Psiformer vs FermiNet: Architecture & Attention

- FermiNet layers mix electron features via fixed functions; Psiformer uses self-attention.
- Self-attention: each electron attends to all others (learns interactions).
- Both ansatzes enforce antisymmetry via Slater determinants.
- Psiformer captures correlations more effectively with fewer parameters.
- Attention mechanism is permutation-invariant and scales to complex interactions.



# Computational power & scaling

- Solving many-electron systems is computationally intensive.
- Computational cost grows steeply with electron number  $N$ .
- Wavefunction evaluation involves expensive operations (e.g. determinants,  $O(N^3)$ ).
- Requires significant computing resources (GPUs/TPUs, parallelization).

# Numerical stability

- Exponential and cusp factors can cause numerical overflow/underflow if not handled carefully.
- Ensuring stable Monte Carlo estimates (variance reduction techniques).
- Feature scaling and proper initialization improve training stability.
- Jastrow factor helps manage large energy fluctuations (by satisfying cusps).

# Scaling with electron count

- Wavefunction resides in a  $3N$ -dimensional configuration space.
- Monte Carlo integration becomes harder as  $N$  increases (high dimensionality).
- More electrons often require deeper or wider neural networks.
- Ongoing research into architectures that scale better or transfer to larger  $N$ .

# Thanks

Thank you for your attention!