

A Course in Applied Econometrics

Lecture 3: Linear Panel Data Models, I

Jeff Wooldridge
IRP Lectures, UW Madison, August 2008

1. Overview of the Basic Model
2. New Insights Into Old Estimators
3. Behavior of Estimators without Strict Exogeneity
4. IV Estimation under Sequential Exogeneity

1

1. Overview of the Basic Model

- Unless stated otherwise, the methods discussed in these slides are for the case with a large cross section and small time series, although some approximations are based on T increasing.
- For a generic i in the population,

$$y_{it} = \eta_t + \mathbf{x}_{it}\boldsymbol{\beta} + c_i + u_{it}, \quad t = 1, \dots, T, \quad (1)$$

where η_t is a separate time period intercept (parameters that can be estimated), \mathbf{x}_{it} is a $1 \times K$ vector of explanatory variables, c_i is the time-constant unobserved effect, and the $\{u_{it} : t = 1, \dots, T\}$ are idiosyncratic errors. We view the c_i as random draws along with the observed variables.

2

- An attractive assumption (but, of course, not universally applicable) is *contemporaneous exogeneity conditional on c_i* :

$$E(u_{it}|\mathbf{x}_{it}, c_i) = 0, \quad t = 1, \dots, T. \quad (2)$$

This equation defines $\boldsymbol{\beta}$ in the sense that under (1) and (2),

$$E(y_{it}|\mathbf{x}_{it}, c_i) = \eta_t + \mathbf{x}_{it}\boldsymbol{\beta} + c_i, \quad (3)$$

so the β_j are partial effects holding c_i fixed.

- Unfortunately, $\boldsymbol{\beta}$ is not identified only under (2). If we add the strong assumption $Cov(\mathbf{x}_{it}, c_i) = \mathbf{0}$, then $\boldsymbol{\beta}$ is identified.

3

- Allow any correlation between \mathbf{x}_{it} and c_i by assuming *strict exogeneity conditional on c_i* :

$$E(u_{it}|\mathbf{x}_{i1}, \mathbf{x}_{i2}, \dots, \mathbf{x}_{iT}, c_i) = 0, \quad t = 1, \dots, T, \quad (4)$$

which can be expressed as

$$E(y_{it}|\mathbf{x}_i, c_i) = E(y_{it}|\mathbf{x}_{it}, c_i) = \eta_t + \mathbf{x}_{it}\boldsymbol{\beta} + c_i. \quad (5)$$

If $\{\mathbf{x}_{it} : t = 1, \dots, T\}$ has suitable time variation, $\boldsymbol{\beta}$ can be consistently estimated by fixed effects (FE) or first differencing (FD), or generalized least squares (GLS) or generalized method of moments (GMM) versions of them.

4

- Make inference fully robust to heteroskedasticity and serial dependence, even if use GLS. With large N and small T , there is little excuse not to compute “cluster” standard errors.
- Violation of strict exogeneity: always if \mathbf{x}_{it} contains lagged dependent variables, but also if changes in u_{it} cause changes in $\mathbf{x}_{i,t+1}$ (“feedback effect”).
- *Sequential exogeneity condition on c_i :*

$$E(u_{it}|\mathbf{x}_{i1}, \mathbf{x}_{i2}, \dots, \mathbf{x}_{it}, c_i) = 0, t = 1, \dots, T \quad (6)$$

or, maintaining the linear model,

$$E(y_{it}|\mathbf{x}_{i1}, \dots, \mathbf{x}_{it}, c_i) = E(y_{it}|\mathbf{x}_{it}, c_i). \quad (7)$$

5

Allows for lagged dependent variables and other feedback. Generally, β is identified under sequential exogeneity. (More later.)

- The key “random effects” assumption is

$$E(c_i|\mathbf{x}_i) = E(c_i). \quad (8)$$

Pooled OLS or any GLS procedure, including the RE estimator, are consistent. Fully robust inference is available for both.

- It is useful to define two *correlated random effects* assumptions. The first just defines a linear projection:

$$L(c_i|\mathbf{x}_i) = \psi + \mathbf{x}_i\xi, \quad (9)$$

Called the *Chamberlain device*, after Chamberlain (1982).

6

- Mundlak (1978) used a restricted version

$$E(c_i|\mathbf{x}_i) = \psi + \bar{\mathbf{x}}_i\xi, \quad (10)$$

where $\bar{\mathbf{x}}_i = T^{-1} \sum_{t=1}^T \mathbf{x}_{it}$. Then

$$y_{it} = \eta_t + \mathbf{x}_{it}\beta + \bar{\mathbf{x}}_i\xi + a_i + u_{it}, \quad (11)$$

and we can apply pooled OLS or RE because $E(a_i + u_{it}|\mathbf{x}_i) = 0$. Both equal the FE estimator of β .

- (10) is in the spirit of approaches for nonlinear models, where often an entire conditional distribution is specified.

7

- Equation (11) makes it easy to compute a fully robust Hausman test comparing RE and FE. Separate covariates into aggregate time effects, time-constant variables, and variables that change across i and t :

$$y_{it} = \mathbf{g}_t\eta + \mathbf{z}_t\gamma + \mathbf{w}_{it}\delta + c_i + u_{it}. \quad (12)$$

We cannot estimate γ by FE, so it is not part of the Hausman test comparing RE and FE. Less clear is that coefficients on the time dummies, η , cannot be included, either. (RE and FE estimation only with aggregate time effects are identical.) We can only compare $\hat{\delta}_{FE}$ and $\hat{\delta}_{RE}$ (M parameters).

8

- Convenient test:

$$y_{it} \text{ on } \mathbf{g}_t, \mathbf{z}_i, \mathbf{w}_{it}, \bar{\mathbf{w}}_i, t = 1, \dots, T; i = 1, \dots, N, \quad (13)$$

which makes it clear there are M restrictions to test. Pooled OLS or RE, fully robust!

- Regression (13) can also be used to estimate coefficients on \mathbf{z}_i while allowing correlation between c_i and $\bar{\mathbf{w}}_i$. For these estimated coefficients to be consistent for $\boldsymbol{\gamma}$, we would assume

$$E(c_i|\mathbf{z}_i, \bar{\mathbf{w}}_i) = E(c_i|\bar{\mathbf{w}}_i) = \psi + \bar{\mathbf{w}}_i \boldsymbol{\xi} \text{ (or the linear projection version).}$$

- Be cautious using canned procedures: the df are often wrong (the aggregate time variables are included) and the tests are nonrobust.

2. New Insights Into Old Estimators

- Consider an extension of the usual model to allow for unit-specific slopes,

$$y_{it} = c_i + \mathbf{x}_{it} \mathbf{b}_i + u_{it} \quad (14)$$

$$E(u_{it}|\mathbf{x}_i, c_i, \mathbf{b}_i) = 0, t = 1, \dots, T, \quad (15)$$

where \mathbf{b}_i is $K \times 1$. We act as if \mathbf{b}_i is constant for all i but think c_i might be correlated with \mathbf{x}_{it} ; we apply usual FE estimator. When does the usual FE estimator consistently estimate the population average effect, $\boldsymbol{\beta} = E(\mathbf{b}_i)$?

- A sufficient condition for consistency of the FE estimator, along with along with (15) and the usual rank condition, is

$$E(\mathbf{b}_i|\bar{\mathbf{x}}_{it}) = E(\mathbf{b}_i) = \boldsymbol{\beta}, \quad t = 1, \dots, T \quad (16)$$

where $\bar{\mathbf{x}}_{it}$ are the time-demeaned covariates. Allows the slopes, \mathbf{b}_i , to be correlated with the regressors \mathbf{x}_{it} through permanent components. For example, if $\mathbf{x}_{it} = \mathbf{f}_i + \mathbf{r}_{it}, t = 1, \dots, T$. Then (16) holds if

$$E(\mathbf{b}_i|\mathbf{r}_{i1}, \mathbf{r}_{i2}, \dots, \mathbf{r}_{iT}) = E(\mathbf{b}_i).$$

- Extends to a more general class of estimators. Write

$$y_{it} = \mathbf{w}_t \mathbf{a}_i + \mathbf{x}_{it} \mathbf{b}_i + u_{it}, \quad t = 1, \dots, T \quad (17)$$

where \mathbf{w}_t is a set of deterministic functions of time. FE now sweeps away \mathbf{a}_i by netting out \mathbf{w}_t from \mathbf{x}_{it} .

- In the random trend model, $\mathbf{w}_t = (1, t)$. If $\mathbf{x}_{it} = \mathbf{f}_i + \mathbf{h}_i t + \mathbf{r}_{it}$, then \mathbf{b}_i can be arbitrarily correlated with $(\mathbf{f}_i, \mathbf{h}_i)$.
- Generally, need $\dim(\mathbf{w}_t) < T$.

- Can apply to models with time-varying factor loads, η_t :

$$y_{it} = \mathbf{x}_{it}\boldsymbol{\beta} + \eta_t c_i + u_{it}, t = 1, \dots, T. \quad (18)$$

Sufficient for consistency of FE estimator that ignores the η_t is

$$\text{Cov}(\tilde{\mathbf{x}}_{it}, c_i) = \mathbf{0}, t = 1, \dots, T. \quad (19)$$

- Now let some elements of \mathbf{x}_{it} be correlated with $\{u_{ir} : r = 1, \dots, T\}$, but with strictly exogenous instruments (conditional on c_i). Assume

$$E(u_{it} | \mathbf{z}_i, \mathbf{a}_i, \mathbf{b}_i) = 0. \quad (20)$$

Also, replace (16) with

$$E(\mathbf{b}_i | \tilde{\mathbf{z}}_{it}) = E(\mathbf{b}_i) = \boldsymbol{\beta}, \quad t = 1, \dots, T. \quad (21)$$

Still not enough. A sufficient condition is

$$\text{Cov}(\tilde{\mathbf{x}}_{it}, \mathbf{b}_i | \tilde{\mathbf{z}}_{it}) = \text{Cov}(\tilde{\mathbf{x}}_{it}, \mathbf{b}_i), t = 1, \dots, T. \quad (22)$$

$\text{Cov}(\tilde{\mathbf{x}}_{it}, \mathbf{b}_i)$, a $K \times K$ matrix, need not be zero, or even constant across time. The *conditional* covariance cannot depend on the time-demeaned instruments. Then, FEIV is consistent for $\boldsymbol{\beta} = E(\mathbf{b}_i)$ provided a full set of time dummies is included.

- Assumption (22) cannot be expected to hold when endogenous elements of \mathbf{x}_{it} are discrete.

3. Behavior of Estimators without Strict Exogeneity

- Both the FE and FD estimators are inconsistent (with fixed $T, N \rightarrow \infty$) without the strict exogeneity assumption. But inconsistencies (as function of T) can differ.
- If we maintain $E(u_{it} | \mathbf{x}_{it}, c_i) = 0$ and assume $\{(\mathbf{x}_{it}, u_{it}) : t = 1, \dots, T\}$ is “weakly dependent”, can show

$$\text{plim}_{N \rightarrow \infty} \hat{\boldsymbol{\beta}}_{FE} = \boldsymbol{\beta} + O(T^{-1}) \quad (23)$$

$$\text{plim}_{N \rightarrow \infty} \hat{\boldsymbol{\beta}}_{FD} = \boldsymbol{\beta} + O(1). \quad (24)$$

- Interestingly, still holds if $\{\mathbf{x}_{it} : t = 1, \dots, T\}$ has unit roots as long as $\{u_{it}\}$ is $I(0)$ and contemporaneous exogeneity holds.

- Catch: if $\{u_{it}\}$ is $I(1)$ – so that the time series “model” is a spurious regression (y_{it} and \mathbf{x}_{it} are not *cointegrated*), then (23) is no longer true. FD eliminates any unit roots.
- Same conclusions hold for IV versions: FE has bias of order T^{-1} if $\{u_{it}\}$ is weakly dependent.
- Simple test for lack of strict exogeneity in covariates:

$$y_{it} = \eta_t + \mathbf{x}_{it}\boldsymbol{\beta} + \mathbf{w}_{i,t+1}\boldsymbol{\delta} + c_i + e_{it} \quad (25)$$

Estimate the equation by fixed effects and test $H_0 : \boldsymbol{\delta} = \mathbf{0}$.

- Test contemporaneous endogeneity of a subset of certain regressors.

$$y_{it1} = \mathbf{z}_{it1}\boldsymbol{\delta}_1 + \mathbf{y}_{it2}\boldsymbol{\alpha}_1 + \mathbf{y}_{it3}\boldsymbol{\gamma}_1 + c_{i1} + u_{it1}, \quad (26)$$

where, in an FE environment, we want to test $H_0 : E(\mathbf{y}_{it3}'u_{it1}) = \mathbf{0}$.

- Reduced form for \mathbf{y}_{it3} :

$$\mathbf{y}_{it3} = \mathbf{z}_{it}\boldsymbol{\Pi}_3 + \mathbf{c}_{i3} + \mathbf{v}_{it3}. \quad (27)$$

- Obtain FE residuals, $\hat{\mathbf{v}}_{it3} = \mathbf{y}_{it3} - \mathbf{z}_{it}\hat{\boldsymbol{\Pi}}_3$ ($\hat{\boldsymbol{\Pi}}_3$ FE estimates). Estimate

$$y_{it1} = \mathbf{z}_{it1}\boldsymbol{\delta}_1 + \mathbf{y}_{it2}\boldsymbol{\alpha}_1 + \mathbf{y}_{it3}\boldsymbol{\gamma}_1 + \hat{\mathbf{v}}_{it3}\boldsymbol{\rho}_1 + error_{it1} \quad (28)$$

by FEIV, using instruments $(\mathbf{z}_{it}, \mathbf{y}_{it3}, \hat{\mathbf{v}}_{it3})$. \mathbf{y}_{it3} exogenous: use (robust) test that $\boldsymbol{\rho}_1 = \mathbf{0}$; need not adjust for the first-step estimation.

4. IV Estimation under Sequential Exogeneity

We now consider IV estimation of the model

$$y_{it} = \mathbf{x}_{it}\boldsymbol{\beta} + c_i + u_{it}, \quad t = 1, \dots, T, \quad (29)$$

under sequential exogeneity assumptions; the weakest form is

$$Cov(\mathbf{x}_{is}, u_{it}) = 0, \text{ all } s \leq t.$$

This leads to simple moment conditions after first differencing:

$$E(\mathbf{x}_{is}'\Delta u_{it}) = \mathbf{0}, \quad s = 1, \dots, t-1; \quad t = 2, \dots, T. \quad (30)$$

Therefore, at time t , the available instruments in the FD equation are in the vector $\mathbf{x}_{it}^o \equiv (\mathbf{x}_{i1}, \mathbf{x}_{i2}, \dots, \mathbf{x}_{it})$. The matrix of instruments is

$$\mathbf{W}_i = \text{diag}(\mathbf{x}_{i1}^o, \mathbf{x}_{i2}^o, \dots, \mathbf{x}_{iT-1}^o), \quad (31)$$

which has $T-1$ rows. Routine to apply GMM estimation.

- Simple strategy: estimate a reduced form for $\Delta \mathbf{x}_{it}$ separately for each t . So, at time t , run the regression $\Delta \mathbf{x}_{it}$ on $\mathbf{x}_{i,t-1}^o$, $i = 1, \dots, N$, and obtain the fitted values, $\widehat{\Delta \mathbf{x}_{it}}$. Then, estimate the FD equation

$$\Delta y_{it} = \Delta \mathbf{x}_{it}\boldsymbol{\beta} + \Delta u_{it}, \quad t = 2, \dots, T \quad (32)$$

by pooled IV using instruments (not regressors) $\widehat{\Delta \mathbf{x}_{it}}$.

- Can suffer from a weak instrument problem when $\Delta \mathbf{x}_{it}$ has little correlation with $\mathbf{x}_{i,t-1}^o$.

- If we assume

$$E(u_{it} | \mathbf{x}_{it}, y_{i,t-1}, \mathbf{x}_{i,t-1}, \dots, y_{i1}, \mathbf{x}_{i1}, c_i) = 0, \quad (33)$$

many more moment conditions are available. Using linear functions only, for $t = 3, \dots, T$,

$$E[(\Delta y_{i,t-1} - \Delta \mathbf{x}_{i,t-1}\boldsymbol{\beta})'(y_{it} - \mathbf{x}_{it}\boldsymbol{\beta})] = \mathbf{0}. \quad (34)$$

- Drawback: we often do not want to assume (30). Plus, the conditions in (31) are nonlinear in parameters.

- Arellano and Bover (1995) suggested instead the restrictions

$$Cov(\Delta \mathbf{x}_{it}', c_i) = 0, \quad t = 2, \dots, T, \quad (35)$$

which imply linear moment conditions in the levels equation,

$$E[\Delta \mathbf{x}_{it}'(y_{it} - \alpha - \mathbf{x}_{it}\boldsymbol{\beta})] = \mathbf{0}, \quad t = 2, \dots, T. \quad (36)$$

- Simple AR(1) model:

$$y_{it} = \rho y_{i,t-1} + c_i + u_{it}, \quad t = 1, \dots, T. \quad (37)$$

- Typically, the minimal assumptions imposed are

$$E(y_{is}u_{it}) = 0, \quad s = 0, \dots, t-1, \quad t = 1, \dots, T, \quad (38)$$

so for $t = 2, \dots, T$,

$$E[y_{is}(\Delta y_{it} - \rho \Delta y_{i,t-1})] = 0, \quad s \leq t-2. \quad (39)$$

Again, can suffer from weak instruments when ρ is close to unity.

Blundell and Bond (1998) showed that if the condition

$$Cov(\Delta y_{i1}, c_i) = Cov(y_{i1} - y_{i0}, c_i) = 0 \quad (40)$$

is added to $E(u_{it}|y_{i,t-1}, \dots, y_{i0}, c_i) = 0$ then

$$E[\Delta y_{i,t-1}(y_{it} - \alpha - \rho y_{i,t-1})] = 0 \quad (41)$$

which can be added to the usual moment conditions (38). We have two sets of moments linear in the parameters.

- Condition (40) can be interpreted as a restriction on the initial condition, y_{i0} . Write y_{i0} as a deviation from its steady state, $c_i/(1 - \rho)$ (obtained for $|\rho| < 1$ by recursive substitution and then taking the limit), as $y_{i0} = c_i/(1 - \rho) + r_{i0}$. Then $(1 - \rho)y_{i0} + c_i = (1 - \rho)r_{i0}$, and so (40) reduces to

$$Cov(r_{i0}, c_i) = 0. \quad (42)$$

The deviation of y_{i0} from its SS is uncorrelated with the SS.

- Extensions of the AR(1) model, such as

$$y_{it} = \rho y_{i,t-1} + \mathbf{z}_{it}\boldsymbol{\gamma} + c_i + u_{it}, \quad t = 1, \dots, T \quad (43)$$

and use FD:

$$\Delta y_{it} = \rho \Delta y_{i,t-1} + \Delta \mathbf{z}_{it}\boldsymbol{\gamma} + \Delta u_{it}, \quad t = 1, \dots, T. \quad (44)$$

- Airfare example in notes. The data set can be obtained from the web site for Wooldridge (2002), and is called AIRFARE.RAW:

Dependent Variable:	<i>lfare</i>		
	(1)	(2)	(3)
Explanatory Variable	Pooled OLS	Pooled IV	Arellano-Bond
<i>lfare</i> ₋₁	-.126	.219	.333
	(.027)	(.062)	(.055)
<i>concen</i>	.076	.126	.152
	(.053)	(.056)	(.040)
<i>N</i>	1,149	1,149	1,149

- Arellano and Alvarez (2003) show that the GMM estimator that accounts for the MA(1) serial correlation in the FD errors has better properties when T and N are both large.