

Interstellar Interceptors

Mission design for rendezvous with objects in hyperbolic orbits

Jorge Martínez

Supervised by:

Josep M. Trigo-Rodríguez (ICE-CSIC/IEEC)
Eloy Peña-Asensio (Politecnico di Milano)


Universidad Internacional de Valencia

May 22, 2024

What are interstellar objects?

Definition

Interstellar objects (ISOs) are asteroids, comets or planetary bodies moving through interstellar medium (ISM) without being gravitationally bound to a star.



ISOs follow hyperbolic orbits

Why are interstellar objects important?

They present a unique opportunity to study extraterrestrial bodies that have traversed vast cosmic distances.

Why are interstellar objects important?

They present a unique opportunity to study extraterrestrial bodies that have traversed vast cosmic distances.

Their study can unleash information about:

- Better understanding the formation of planetary systems
- Exploring their physical and chemical composition
- Technological motivation

Why are interstellar objects important?

They present a unique opportunity to study extraterrestrial bodies that have traversed vast cosmic distances.

Their study can unleash information about:

- Better understanding the formation of planetary systems
- Exploring their physical and chemical composition
- Technological motivation

Motivation of this work

Design orbits for rendezvous with ISOs to study their physical properties.

Discovered interstellar objects

There are two confirmed ISOs to this day:



1I/'Oumuamua

ESO's VLT and GST Telescopes




2I/Borisov

NASA Hubble Space Telescope

Discovered interstellar objects

There are two confirmed ISOs to this day:



1I/'Oumuamua

ESO's VLT and GST Telescopes




2I/Borisov

NASA Hubble Space Telescope


These interlopers present the following orbit attributes:

- Hyperbolic orbits
- High relative velocity w.r.t. the Sun
- Random inclination
- Discovered close to the direction of the Solar Apex


Orbits of 1I/'Oumuamua and 2I/Borisov




1I/'Oumuamua orbit top view



2I/Borisov orbit top view




1I/'Oumuamua orbit side view



2I/Borisov orbit side view

Navigating through space: the Lambert's problem

Lambert's problem is the Boundary Value Problem (BVP) in the context of the restricted two-body problem dynamics.




$$\ddot{\vec{r}} = -\frac{\mu}{r^3}\vec{r} \quad \left\{ \begin{array}{l} \vec{r}(t_1) = \vec{r}_1 \\ \vec{r}(t_2) = \vec{r}_2 \end{array} \right.$$

Geometry of the Lambert's problem

Navigating through space: the Lambert's problem

Lambert's problem is the Boundary Value Problem (BVP) in the context of the restricted two-body problem dynamics.



$$\ddot{\vec{r}} = -\frac{\mu}{r^3}\vec{r} \quad \left\{ \begin{array}{l} \vec{r}(t_1) = \vec{r}_1 \\ \vec{r}(t_2) = \vec{r}_2 \end{array} \right.$$

Solve for the orbit which passes through \vec{r}_1 and \vec{r}_2 over a finite amount of time $\Delta t = t_2 - t_1$.

Geometry of the Lambert's problem

Estimating the cost of the maneuver using the C_3 energy

Lambert's problem computes the initial velocity \vec{v}_1 and final velocity \vec{v}_2 of the orbit.

- First impulse: $\Delta v_1 = \|\vec{v}_1 - \vec{v}_{\text{origin}}\|$
- Last impulse: $\Delta v_2 = \|\vec{v}_2 - \vec{v}_{\text{iso}}\|$

Estimating the cost of the maneuver using the C_3 energy

Lambert's problem computes the initial velocity \vec{v}_1 and final velocity \vec{v}_2 of the orbit.

- First impulse: $\Delta v_1 = \|v_1 - v_{\text{origin}}\|$
- Last impulse: $\Delta v_2 = \|v_2 - v_{\text{iso}}\|$

The total cost of the maneuver is $\Delta v = \Delta v_1 + \Delta v_2$. This relates with the fuel mass via the Tsiolkovsky rocket equation:

$$\Delta v = v_e \ln \left(\frac{m_0}{m_f} \right)$$

Estimating the cost of the maneuver using the C_3 energy

Lambert's problem computes the initial velocity \vec{v}_1 and final velocity \vec{v}_2 of the orbit.

- First impulse: $\Delta v_1 = \|v_1 - v_{\text{origin}}\|$
- Last impulse: $\Delta v_2 = \|v_2 - v_{\text{iso}}\|$


The total cost of the maneuver is $\Delta v = \Delta v_1 + \Delta v_2$. This relates with the fuel mass via the Tsiolkovsky rocket equation:

$$\Delta v = v_e \ln \left(\frac{m_0}{m_f} \right)$$

The characteristic energy for hyperbolic orbits C_3 is defined as:

$$C_3 = v_\infty^2$$

Modern launching technologies




Maximum payload mass vs. C_3 energy for different launchers

Minimizing the cost of the maneuver

Porkchop plots are used to find the optimal launch and arrival dates by solving Lambert's problem for a variety of trajectories.

Minimizing the cost of the maneuver

Porkchop plots are used to find the optimal launch and arrival dates by solving Lambert's problem for a variety of trajectories.



Porkchop plot between Earth and Mars

Analyzed scenarios


The analyzed scenarios in this work for each discovered ISO include:

- Direct transfer between the Earth and the ISO
- Direct transfer between the L2 point and the ISO

Analyzed scenarios


The analyzed scenarios in this work for each discovered ISO include:

- Direct transfer between the Earth and the ISO
- Direct transfer between the L2 point and the ISO




Lagrange points for the Sun Earth-Moon system

11/'Oumuamua: direct prograde transfer from Earth



1I/'Oumuamua: direct prograde transfer from L2

Detailed launch energy C_3 and time of flight
L2 - 1I/'Oumuamua direct and prograde transfers between 2016 and 2019



Direct transfer from L2

1I/'Oumuamua: summary of results

Δv launch Earth [km/s]	Δv launch L2 [km/s]	Reduction [%]
13.85	3.80	72.56

Launch energy comparison


ΔV arrival Earth [km/s]	ΔV arrival L2 [km/s]	Reduction [%]
62.33	61.46	1.40

Arrival velocity comparison

C_3 launch Earth [km ² /s ²]	C_3 launch L2 [km ² /s ²]	Reduction [%]
192.00	14.41	92.51

Characteristic energy comparison


2I/Borisov: direct prograde transfer from Earth



Direct transfer from Earth

2I/Borisov: direct prograde transfer from L2

Detailed launch energy C_3 and time of flight
L2 - 2I/Borisov direct and prograde optimum transfer



Direct transfer from Earth

21/Borisov: summary of results

Δv launch Earth [km/s]	Δv launch L2 [km/s]	Reduction [%]
16.90	5.85	65.38

Launch energy comparison

ΔV arrival Earth [km/s]	ΔV arrival L2 [km/s]	Reduction [%]
33.00	33.02	-0.06

Arrival velocity comparison

C_3 launch Earth [km ² /s ²]	C_3 launch L2 [km ² /s ²]	Reduction [%]
286.00	34.30	88.08

Characteristic energy comparison

Conclusions

L2 as the optimal launching point

Launching an intercepting spacecraft from L2 is more fuel efficient. These results agree with the ones presented by the Comet Interceptor mission.

Conclusions

L2 as the optimal launching point

Launching an intercepting spacecraft from L2 is more fuel efficient. These results agree with the ones presented by the Comet Interceptor mission.

A direct transfer is possible

Existing propulsive technologies allow for a direct transfer from Earth to an ISO with similar characteristics to 1I/'Oumuamua and 2I/Borisov.

Conclusions

L2 as the optimal launching point

Launching an intercepting spacecraft from L2 is more fuel efficient. These results agree with the ones presented by the Comet Interceptor mission.

A direct transfer is possible

Existing propulsive technologies allow for a direct transfer from Earth to an ISO with similar characteristics to 1I/'Oumuamua and 2I/Borisov.

Need of increased vigilance in the search for ISOs

Both optimum transfer dates take place prior the discovery of 1I/'Oumuamua and 2I/Borisov. Surveillance of the sky is crucial to detect future ISOs.

Future work

Generating a synthetic population of ISOs

Synthetic orbits of ISOs can be generated by using Monte-Carlo simulations. This could allow to predict future ISO encounters.

Future work

Generating a synthetic population of ISOs

Synthetic orbits of ISOs can be generated by using Monte-Carlo simulations. This could allow to predict future ISO encounters.

Drafting optimum transfers

Optimum transfers can be solved for previous synthetic data. Solutions would be used by future missions parked at L2.

Future work

Generating a synthetic population of ISOs

Synthetic orbits of ISOs can be generated by using Monte-Carlo simulations. This could allow to predict future ISO encounters.

Drafting optimum transfers


Optimum transfers can be solved for previous synthetic data. Solutions would be used by future missions parked at L2.

Trajectory optimization

A combination of multiple gravity assists and deep space maneuvers may allow for a more efficient transfer to an ISO, depending on the scenario.

References

All this work is reproducible, including figures and results.



Document and figures generated via LaTeX and Python

Official repository

All source code is hosted in <https://github.com/jorgepiloto/tfm>


Additional materials: 1I/'Oumuamua shape



1I/'Oumuamua shape


Illustration by William Hartmann/Michael Belton/AP

Additional materials: ISEE-3 mission




ISEE-3 mission
Illustration by NASA

Additional materials: NEO surveyor



NEO surveyor
Illustration by NASA

Additional materials: Large Synoptic Survey Telescope



LSST

Illustration by LSST Project Office