

A NEW STRONG INHIBITOR OF BETA-MANNOSIDASE

H. THERISOD[#], Y. LETOURNEUX and M. THERISOD^{*}

*S.E.S.N.A.B., Pole Sciences et Technologie, Université de La Rochelle,
F-17042 La Rochelle cedex 1*

*[#]Laboratoire des Endotoxines (ERS 571) Bat. 432, Université Paris-Sud,
F-91405 Orsay cedex*

Received 28 October 1997; accepted 13 January 1998


Abstract: N-phenyl-carbamate of D-mannonohydroxymolactone (**I**) was synthesized from mannose and was shown to be the best competitive inhibitor of beta-mannosidase so far reported ($K_i = 25$ nM).

© 1998 Elsevier Science Ltd. All rights reserved.

Natural as well as synthetic inhibitors of α -mannosidases have been the focus of much interest, since some of them (such as swainsonine) have been shown to have potential therapeutic applications as anticancer drugs¹. On the other hand, only a few inhibitors of β -mannosidases have been described. Moreover, most of the numerous known inhibitors of α -mannosidases have poor, if any, action on β -mannosidases.

β -mannosidases are involved in number of biological processes like degradation of glycoproteins, germination of seeds and sporangia². It is one of the numerous glycosidases acting on hemicelluloses present in wood degrading organisms³ (fungi, termites, bacteria). In all these cases, inhibitors acting at very low concentration would be of interest in theoretical studies as well as in practical applications.

The syntheses of several derivatives of glyconolactone-oximes which are transition-state analogue inhibitors of glucosidases^{4,5} and N-acetylglucosaminidase⁶ have been published. We report here the synthesis of two strong inhibitors of β -mannosidase belonging to this class of products.



Compounds **I** and **II** were synthesized in a moderate yield of 50 %, following the general method of Vasella, *via* the known mannonohydroximolactone peracetate^{4,7} and as previously reported for derivatives of several glyconolactone-oximes^{4,5}. **I** and **II** were tested for their inhibitory activities on α -mannosidase from Jack beans and β -mannosidase from snails, using p-nitrophenylmannopyranosides as substrates ($K_M = 1.2$ and 0.57 mM respectively). Both compounds **I** and **II** were found to be competitive inhibitors of the two enzymes with comparable, modest activities on α -mannosidase (K_i 80 and 50 μ M respectively). **II** was found to be a good inhibitor of β -mannosidase ($K_i = 10$ μ M). Compound **I** strongly inhibited β -mannosidase, with $K_i = 25$ nM (Fig. 1). Thus, compound **I** is the best inhibitor of β -mannosidase so far reported.

^{*} Present adress: LCBB, Bat. 420, Université Paris-Sud, F-91405 Orsay cedex. Email: therisod@icmo.u-psud.fr




Fig. 1: Lineweaver-Burk plots of inhibition of β -mannosidase by **I** (left) and **II** (right). Conditions: p-nitrophenyl- β -mannoside as substrate, acetate buffer pH 4.5, 34°C; reaction stopped after 2–4 min by addition of sodium borate, then lecture of absorbance at 410 nm. Concentrations of inhibitor: 40 and 200 nM of **I** (left); 40 and 100 μM of **II** (right).

The *gluco*-analogue of **I** was previously reported to be a good inhibitor of β -glucosidase, and a modest inhibitor of α -glucosidase ($K_i = 2.5$ and $75 \mu\text{M}$ respectively)^{4,5}. It is a better inhibitor of β -glucosidase than gluconolactone ($K_i = 30 \mu\text{M}$). Mannonolactone itself is a good inhibitor of β -mannosidase ($K_i = 17 \mu\text{M}$). Thus, the behavior of **I** on β -mannosidase is quite parallel to that previously reported for the *gluco*-analogue on β -glucosidase. As demonstrated for this last compound, glyconohydroximolactones are transition-state analogues⁴. In addition, like other inhibitors bearing an heteroatom on the "anomeric" position, they are thought to act by hydrogen bonding between this heteroatom and the catalytic carboxylic acid of the active site⁸. The efficiency of the inhibitor varies with the strength of this hydrogen bond, and, consequently, with the basicity of the heteroatom (for example, gluconohydroxymolactam is a better inhibitor of β -glucosidase than gluconohydroximolactone: $K_i = 2.9 \mu\text{M}$, versus $100 \mu\text{M}$ ⁹). **II**, although a good inhibitor, is less performant than **I**: One can expect the nitrogen atom in **II** to be less basic than in **I**; The aromatic ring in **II** is closer to the nitrogen than in **I**, and in that part of the molecule, there is less flexibility in **II** than in **I**. This could prevent the formation of a strong hydrogen bond, since this has been demonstrated to be in the plane of the sugar ring⁸.

References:

- Jacob, G.S. *Current Opinion in Structural Biology* **1995**, *5*, 605.
- Walter, J.K.; Aach, H.G. *Physiol. Plant.* **1987**, *70*, 485. Bewley, J.D.; Leung, D.W.; Ouellette, F.B. *Recent. Adv. Phytochem.* **1983**, *17*, 137. Mc Cleary, B.V.; Matheson, N.K. *Phytochemistry* **1975**, *14*, 1187. Goldberg, R. *C.R. Acad. Sci. Paris, Ser. D* **1967**, 264, 1036.
- Sutherland, I.W. *Carbohydr. Res.* **1984**, *131*, 93; Rouland, C.; Chararas, C.; Renoux, J. *C.R. Acad. Sci. Paris* **1989**, *308 III*, 281; Kohring, S.; Wiegel, J.; Mayer, F. *Appl. Environ. Microbiol.* **1990**, *56*, 3798.
- Beer, D.; Vasella, A. *Helvetica Chim. Acta* **1985** *68*, 2254; Beer, D.; Vasella, A. *Helvetica Chim. Acta* **1986**, *69*, 267.
- Therisod, M.; Therisod, H.; Lubineau, A. *Biomed. Chem. Letters* **1995**, *5*, 2055.
- Beer, D.; Maloisel, J.L.; Rast, D.M.; Vasella, A. *Helvetica Chim. Acta* **1990**, *73*, 1918.
- Ghosez, A.; Göbel, T.; Giese, B. *Chem. Ber.* **1988**, *121*, 1807.
- Heightman, T.; Locatelli, M.; Vasella, A. *Helvetica Chim. Acta* **1996**, *79*, 2190; Granier, T.; Panday, N.; Vasella, A. *Helvetica Chim. Acta* **1997**, *80*, 979.
- Hoos, R.; Vasella, A.; Rupitz, K.; Withers, S. *Carbohydr. Res.* **1997**, 298, 291.