

THROMBIN RECEPTOR (PAR-1) ANTAGONISTS. HETEROCYCLE-BASED PEPTIDOMIMETICS OF THE SFLLR AGONIST MOTIF

William J. Hoekstra,* Becky L. Hulshizer, David F. McComsey,
Patricia Andrade-Gordon, Jack A. Kauffman, Michael F. Addo,
Donna Oksenberg,[†] Robert M. Scarborough,[†] and Bruce E. Maryanoff

*The R. W. Johnson Pharmaceutical Research Institute, Spring House, PA 19477, U.S.A.
and [†]COR Therapeutics, Inc., South San Francisco, CA 94080, U.S.A.*

Received 2 April 1998; accepted 21 May 1998

Abstract: The thrombin receptor (PAR-1) is activated by α -thrombin to stimulate various cell types, including platelets, through the tethered-ligand sequence SFLLRN. A series of azole-based carboxamides, designed after SFLLR, were synthesized and evaluated in vitro. The compounds inhibited platelet aggregation induced by SFLLRN-NH₂ or α -thrombin, and blocked the binding of [³H]-S-(*p*-F-Phe)-Har-L-Har-KY-NH₂ to a CHRF membrane preparation of PAR-1. Oxazole **30** bound to PAR-1 with an IC₅₀ of 1.6 μ M, and gave IC₅₀ values of 25 μ M and 6.6 μ M against α -thrombin- and SFLLRN-NH₂-induced platelet aggregation, respectively.



© 1998 Elsevier Science Ltd. All rights reserved.

Thrombin is a key trypsin-like serine protease in hemostasis that has roles in both coagulation and thrombosis. Its various cellular actions, such as platelet aggregation, lymphocyte mitosis, monocyte chemotaxis, and endothelial cell proliferation,^{1–5} are mediated by specific receptors on the cell surface. The first example of such a receptor was the human thrombin receptor (protease-activated receptor-1; PAR-1), which Coughlin and coworkers cloned, expressed, and identified as a member of the vast G-protein coupled receptor (GPCR) superfamily.⁶ Thrombin activates PAR-1 by proteolytic cleavage of the extracellular domain at the Arg-41/Ser-42 peptide bond to reveal a truncated N-terminus containing the activation sequence SFLLRN, which serves as a “tethered ligand.” Recently, other protease-activated receptors with close homology to the thrombin receptor, PAR-2,^{7,8} PAR-3,^{9a} and PAR-4,^{9b,c} have been cloned as well.



Structure-activity relationships of SFLLR-based agonists have received considerable attention in platelet activation studies.^{4,5,10–12} The minimum structural requirements for agonist peptides are, generally, a small uncapped N-terminal residue at position 1, an aromatic residue at position 2 (with agonist potency enhanced ca. 4-fold by *p*-F-Phe substitution¹³), and a basic or aromatic residue at position 5. A large hydrophobic amino acid at position 4 is important as well, but widely varied substitution is tolerated at position 3.

Some SFLLR analogues have proven to be antagonists of PAR-1 activation in terms of blocking platelet aggregation.^{12–14} In fact, peptide analogues with a cinnamoyl-(*p*-F-Phe) at positions 1/2 and a *p*-guanidino-Phe at position 3 are potent inhibitors of platelet aggregation induced by SFLLRN-NH₂, although no results were reported relative to inhibition of aggregation induced by α -thrombin.^{14,15} For example, *E*-cinnamoyl-(*p*-F-Phe)-(p-guanidino-Phe)-LRR-NH₂ exhibited an IC₅₀ value of 0.021 μ M in platelet aggregation induced by SFLLRN-NH₂ and an IC₅₀ value of 0.0075 μ M in binding to PAR-1 (vs. [³H]-SFLLRN-NH₂).^{14,15} Thus, there is a need for new thrombin receptor antagonists that work against both agonist peptides and α -thrombin, the endogenous activator.

hydrolysis, and coupling again with a basic residue. The stereochemical integrity at the α -position was maintained during saponification of the Cha-type residue with <1% isomerization. Typically, the basic residue was capped as a benzylamide for superior activity (e.g., Arg-NHBn). The arginine side chain was protected with the Pmc (2,2,5,7,8-pentamethylchroman-6-sulfonyl) group, which is conveniently labile to TFA.




The oxazole targets were synthesized by using similar solution-phase methodology. A serine-derived oxazole dipeptide mimetic was prepared by assembling Boc-(*p*-F-Phe)-Ser-OMe (EDC·HCl in CH₂Cl₂; **6**) and cyclizing it with Burgess' reagent to an oxazoline (Eq 3),¹⁸ which was oxidized to an oxazole with *tert*-butylperoxybenzoate/copper (II);¹⁹ intermediate **7** was then used as shown in Eq 2 to prepare target compounds. 5-Methyloxazole **9** was prepared from **8** by reversing the two-step sequence (Eq 4). First, **8** was converted to



a methyl ketone by Dess-Martin oxidation, then cyclized with I_2/Ph_3P to afford **9**.²⁰ Thiazole intermediate **2** turned out to be the most synthetically accessible vis-à-vis oxazoles **7** and **9**; a high-yielding cyclization and a straightforward purification contributed to a useful process for the synthesis of the final thiazole targets.

Fmoc-protected thiazole **10** was introduced at an intermediate stage of our solid-phase synthesis of C-terminal phenylalanine amides (Eq 5). The C-terminal Phe was anchored to a Rink resin for parallel synthesis of arrays of targets. Iterative Fmoc removal and diisopropylcarbodiimide (DIC)/1-hydroxybenzotriazole (HOBT) coupling, then TFA deprotection/resin cleavage, afforded reasonably pure products (>90%) in ca. 50 mg quantities. Given the calculated initial resin loading of 1.0 mmol/g, isolated yields were ca. 40–60%.

Table 1. Platelet aggregation and PAR-1 binding IC_{50} values (μM) for thiazoles




No. ^a	R_1^b	R_2	R_3^c	R_4	R_5	IC_{50} , Human GFP ^d		IC_{50} , PAR-1 Binding ^e
						Thrombin	TRAP-6	
5	Sar	F	Cha	Arg	NHBn	27 ± 4.4	11 ± 1.7	10 ± 3.5
11	Ile	F	Cha	Arg	Phe-NH ₂	32 ± 6.7	4.0 ± 1.3	3.9 ± 0.4
12	Sar	F	Cha	Arg	Phe-NH ₂	41 ± 8.3	14 ± 2.4	5.7 ± 1.8
13	Val	F	Cha	Arg	Phe-NH ₂	22 ± 7.8	4.6 ± 3.5	2.7 ± 0.5
14	Val	OMe	Cha	Arg	Phe-NH ₂	49 ± 13	9.5 ± 1.4	8.2 ± 0.8
15	H	F	Cha	Arg	Phe-NH ₂	28 ± 8.0	16 ± 4.2	8.0
16	Sar	F	Nip	Arg	Phe-NH ₂	>100	>100	>100
17	Sar	F	Arg	Arg	Phe-NH ₂	>100	>100	>100
18	Sar	F	D-Cha	Arg	Phe-NH ₂	96 ± 1.2	42 ± 3.6	94 ± 4.8
19	Ile	F	N-Bn-Gly	Arg	Phe-NH ₂	>100	80 ± 17	>100
20	Sar	F	Cha	Har	Phe-NH ₂	59 ± 7.3	7.7 ± 1.8	5.0
21	Sar	F	Cha	Arg	Cha-NH ₂	46 ± 6.1	6.8 ± 1.7	2.7 ± 0.5
22	Sar	F	Cha	Arg	hPhe-NH ₂	45	5.5	4.1 ± 0.9
23	Sar	F	Cha	Arg	NH(CH ₂) ₂ Ph	27 ± 6.8	10 ± 3.6	5.6 ± 1.8
24	Sar	F	Lys	Arg	NH(CH ₂) ₂ Ph	>100	>100	>100
25	Sar	F	Phe	Arg	NHBn	46 ± 8.8	13 ± 2.8	24 ± 4.8
26	Sar	F	Cha	Arg	NHCH ₂ CH(Me)Ph	40 ± 3.5	16 ± 4.1	24 ± 6

a. See ref 21. b. Sar = sarcosine. c. Cha = cyclohexylalanine; Nip = nipecotic acid. d. α -Thrombin- or SFLLRN-NH₂-induced gel-filtered platelet aggregation (at least $n = 2$; $n = 1$ for values without error limits).²² A level of 80–100% aggregation was achieved at a single concentration per platelet preparation. For SFLLRN-NH₂ ($EC_{50} = 0.30 \pm 0.15 \mu M$), the concentration was 0.5–1.0 μM ; for thrombin ($EC_{50} = 73 \pm 12 pM$), the concentration was 0.05–0.5 nM. e. Inhibition of [³H]-S-(p-F-Phe)-Har-L-Har-KY-NH₂ binding to a thrombin receptor membrane preparation ($n = 2$; $n = 1$ for values without error limits).²³

Biological Results. The thiazole (Table 1) and oxazole (Table 2) derivatives²¹ exemplify 120 target compounds that were synthesized and tested for inhibition of thrombin- and SFLLRN-NH₂-induced platelet aggregation,²² as well as competitive binding of [³H]-S-(*p*-F-Phe)-Har-L-Har-KY-NH₂ (Har = homoarginine) to a membrane preparation of PAR-1.²³ Since little difference in potency was observed between the two heterocycles, thiazoles in excess of oxazoles were prepared because of their relative ease of synthesis. In the oxazole series, H at R₆ was preferred over Me for both binding and inhibition of SFLLR-induced aggregation (**30** vs. **31**). At the N-terminal R₁ position of either series, a small uncapped residue (i.e., Sar or Gly) was preferred for inhibition of binding (**30**, IC₅₀ = 1.6 μM). Fluoro substitution at R₂ was important for activity as H (**34**) or OMe (**14**) substitution afforded only weak antagonists (Table 1). At the R₃ site, a large hydrophobic residue of L-configuration (e.g., L-Cha) appeared necessary, and a guanidine-bearing residue of L-configuration at the R₄ residue was crucial for activity (Table 1). Interestingly, each of these aforementioned amino acid criteria are reflective of the SAR for SFLLR-based agonists. Since an aromatic residue at R₅ had provided good agonist activity,¹² a Phc or aralkyl group was incorporated in these series with success (Tables 1 and 2); an aliphatic residue was also acceptable (**21**). The role of these agents in PAR-2, PAR-3, and PAR-4 recognition is yet to be determined.

Table 2. Platelet aggregation and PAR-1 binding IC₅₀ values (μM) for oxazoles



No. ^a	R ₁ ^b	R ₆	R ₅	IC ₅₀ , Human GFP ^c		IC ₅₀ , PAR-1 Binding ^c
				Thrombin	TRAP-6	
27	Gly	H	H	19 ± 0.6	13 ± 4.2	6.2 ± 0.5
28	Gly	H	CH ₂ Ph	17 ± 4.2	10 ± 1.7	5.4 ± 0.2
29	β-Ala	H	CH ₂ Ph	>100	36 ± 15	6.7 ± 1.3
30	Sar	H	CH ₂ Ph	25 ± 2.5	6.6 ± 1.5	1.6 ± 0.5
31	Sar	Me	CH ₂ Ph	23 ± 3.0	18 ± 2.5	5.0 ± 2.0
32	Dpr	Me	CH ₂ Ph	24 ± 3.1	25 ± 6.9	53 ± 18
33	Ac	Me	CH ₂ Ph	18 ± 2.7	19 ± 4.9	30 ± 3.0
34^d	Sar	Me	CH ₂ Ph	45 ± 4.6	36 ± 7.3	60

a. See ref 21. b. Dpr = 2,3-diaminopropionic acid; Sar = sarcosine. c. See Table 1. d. Des-fluoro-Phe(oxazole).

In conclusion, Sar-oxazole **30** exhibited superior thrombin receptor affinity (IC₅₀ = 1.6 μM), while Ile-thiazole **11** exhibited the best inhibition of platelet aggregation induced by TRAP-6 (IC₅₀ = 4.0 μM). Since thrombin is the endogenous agonist for PAR-1, a prerequisite of pharmacological efficacy for a PAR-1 antagonist would be inhibition of thrombin-induced platelet aggregation. To this end, several analogues (**13**,

27, 28, 30–33) had thrombin-mediated IC_{50} values in the vicinity of 20 μ M, while exhibiting similar inhibitory potency against TRAP-6. Representative compounds from our azole series were inactive in a chromogenic assay for thrombin inhibition at 100 μ M, thereby excluding a direct enzyme-based mechanism of inhibition. Furthermore, antagonist **30** displayed 10-fold weaker platelet aggregation inhibition with arachidonic acid as the agonist (IC_{50} = 63 μ M) than with TRAP-6. Thus, the PAR-1 affinity and inhibition of SFLLRN-NH₂-induced platelet aggregation are consistent with a thrombin receptor mechanism for the biological activity.

References and Notes

- Brass, L. F. *Thromb. Haemostasis* **1995**, *74*, 499.
- Ogletree, M. L.; Natajaran, S.; Seiler, S. M. *Persp. Drug Discovery Des.* **1994**, *1*, 527.
- Coughlin, S. R. *Thromb. Haemostasis* **1993**, *66*, 184.
- Scarborough, R. M.; Naughton, M. A.; Teng, W.; Hung, D. T.; Rose, J. W.; Vu, T.-K.; Wheaton, V. I.; Turck, C. W.; Coughlin, S. R. *J. Biol. Chem.* **1992**, *267*, 13146.
- Darrow, A. L.; Fung-Leung, W.-P.; Ye, R. D.; Santulli, R. J.; Cheung, W.-M.; Derian, C. K.; Burns, C. L.; Damiano, B. P.; Zhou, L.; Keenan, C. M.; Peterson, P. A.; Andrade-Gordon, P. *Thromb. Haemostasis* **1996**, *76*, 860.
- Vu, T.-K. H.; Hung, D. T.; Wheaton, V. I.; Coughlin, S. R. *Cell* **1991**, *64*, 1057.
- Coughlin, S. R. *Proc. Natl. Acad. Sci. U.S.A.* **1994**, *91*, 9200.
- Nystedt, S.; Emilsson, K.; Wahlestedt, C.; Sundelin, J. *Proc. Natl. Acad. Sci. U.S.A.* **1994**, *91*, 9208.
- (a) Ishihara, H.; Connolly, A. J.; Zeng, D.; Kahn, M. L.; Zheng, Y. W.; Timmons, C.; Tram, T.; Coughlin, S. R. *Nature* **1997**, *386*, 502. (b) Coughlin, S.; 1998 ASPET Meeting, San Francisco, CA, Apr. 19, 1998. (c) Davie, E.; Edward P. Kirby Memorial Lecture, Temple University, Philadelphia, PA, May 9, 1998.
- Natarajan, S.; Riexinger, D.; Peluso, M.; Seiler, S. M. *Int. J. Peptide Protein Res.* **1995**, *45*, 145.
- Derian, C. K.; Santulli, R. J.; Tomko, K.; Haertlein, B.; Andrade-Gordon, P. *Thromb. Res.* **1995**, *78*, 505.
- (a) Scarborough, R. M. *PCT Int. Appl.*, 35 pp, WO 9403479. (b) Scarborough, R. M.; Teng, W.; Rose, J. W.; Alves, V.; Arfsten, A.; Naughton, M. A. *Peptides: Chem., Struct. Biol., Proc. 13th Am. Peptide Symp.* **1994**, 695. (Hodges, R. S.; Smith, J. A., Eds.; ESCOM, Leiden, Netherlands).
- Feng, D.-M.; Veber, D. F.; Connolly, T. M.; Nutt, R. F. *Peptides: Biol. Chem., Proc. Third Chin. Peptide Symp.*, **1995**, 84. (Lu, G.-S.; Tam, J. P.; Du, Y.-C., Eds.; ESCOM, Leiden, Netherlands).
- Bernatowicz, M. S.; Klimas, C. E.; Hartl, K. S.; Peluso, M.; Allegretto, N. J.; Seiler, S. M. *J. Med. Chem.* **1996**, *39*, 4879.
- Notably absent from the paper by Bernatowicz et al.¹⁴ were results for inhibition of platelet aggregation induced by the native agonist, α -thrombin.
- A similar approach was used for endothelin receptor antagonists: von Geldern, T. W.; Hutchins, C.; Kester, J. A.; Wu-Wong, J. R.; Chiou, W.; Dixon, D. B.; Oppenorth, T. J. *J. Med. Chem.* **1996**, *39*, 957.
- Boden, C. D. J.; Pattenden, G.; Ye, T. *Synlett* **1995**, 417.
- Wipf, P.; Miller, C. P.; Venkatraman, S.; Fritch, P. C. *Tetrahedron Lett.* **1995**, *36*, 6395.
- Meyers, A. I.; Tavares, F. X. *J. Org. Chem.* **1996**, *61*, 8207.
- Wipf, P.; Miller, C. P. *J. Org. Chem.* **1993**, *58*, 3604.
- All target compounds were homogenous by HPLC with satisfactory ¹H NMR and FABMS data. C-Terminal secondary amides exhibited satisfactory combustion analysis as well. For example, oxazole **30** was isolated as a white powder: mp 127–130°C; [α]_D²⁵ -21.8° (c 0.28, MeOH). FABMS *m/z* 720 (MH⁺); ¹H NMR (DMSO-*d*₆, 400 MHz) δ 0.9 (m, 2 H), 1.1 (m, 4 H), 1.2 (m, 2 H), 1.4–1.8 (m, 12 H), 2.38 (s, 3 H), 3.1 (m, 2 H), 3.3 (m, 2 H), 3.6 (m, 2 H), 4.3 (m, 4 H), 4.6 (m, 1 H), 5.3 (m, 1 H), 7.1 (m, 2 H), 7.2 (m, 7 H), 7.5 (m, 1 H), 7.92 (d, *J* = 6 Hz, 1 H), 8.3 (m, 4 H), 8.60 (s, 1 H), 9.1 (m, 1 H). Analysis calculated for C₃₇H₅₀FN₉O₅•2 TFA: C, 51.95; H, 5.53; N, 13.30. Found: C, 51.55; H, 5.78; N, 13.05.
- Hoekstra, W. J.; Beavers, M. P.; Andrade-Gordon, P.; Evangelisto, M. E.; Keane, P. M.; Press, J. B.; Tomko, K. A.; Fan, F.; Kloczewiak, M.; Mayo, K. H.; Durkin, K. A.; Liotta, D. C. *J. Med. Chem.* **1995**, *38*, 1582.
- A detailed experimental description will be published in due course. Competition assays were performed in CHRF membranes²⁴ by using 10 nM of [³H]-S-(*p*-F-Phe)-Har-L-Har-KY-NH₂. The radiolabeled ligand has a *K_d* value of 15 nM in CHRF membranes. The IC_{50} values reflect the concentration of competing ligand required for 50% inhibition of [³H]-S-(*p*-F-Phe)-Har-L-Har-KY-NH₂ binding.
- Jones, C. L. A.; Witte, D. P.; Feller, M. J.; Fugman, D. A.; Dorn, G. W.; Lieberman, M. A. *Biochim. Biophys. Acta* **1992**, *1136*, 272.