

## SYNTHESIS AND *IN VITRO* CYTOTOXICITY OF C(20)(*RS*)-CAMPTOTHECIN ANALOGUES MODIFIED AT BOTH B (OR A) AND E RING

Sang-sup Jew<sup>1\*</sup>, Myoung Goo Kim<sup>1</sup>, Hee-Jin Kim<sup>1</sup>, Eun-Young Rho<sup>1</sup>, Hyeung-geun Park<sup>1\*</sup>,  
Joon-Kyum Kim<sup>2</sup>, Hyun-Jung Han<sup>2</sup>, and Heesoon Lee<sup>3</sup>

<sup>1</sup>College of Pharmacy, Seoul National University, Seoul 151-742, Korea


<sup>2</sup>Chong Kun Dang Pharmaceutical Co. Seoul 152-070, Korea

<sup>3</sup>College of Pharmacy, Chungbuk National University, Cheongju, Chungbuk 361-763, Korea

Received 3 March 1998; accepted 2 June 1998

**Abstract** : A series of C(7) and C(20)-substituted camptothecin derivatives (**12** - **14**, **16** - **18**) are prepared. Their syntheses and *in vitro* cytotoxicities are reported. © 1998 Elsevier Science Ltd. All rights reserved.


Camptothecin (C(20)(*S*)-**1**) is an alkaloid which was first isolated from chinese tree, *Camptotheca acuminata*<sup>1</sup>. C(20)(*S*)-**1** has been found to have a broad spectrum of antitumor activity<sup>2</sup>, especially against human solid tumor. Mechanistic investigations have shown that the antitumor activity is associated with the inhibition of Topoisomerase I<sup>3</sup> (Topo I) which is essential for the transcription of supercoiled DNA. The above unusual mechanism led to develop semisynthetic analogues, Irinotecan<sup>4</sup> and Topotecan<sup>5</sup> as new commercial antitumor drugs with less toxicity than camptothecin itself. Also the development of total synthetic methods made it possible to investigate the intensive structure-activity relationship studies. Based on the accumulated SAR studies, the introduction of polar groups at C(7), C(9) and C(10) on C(20)(*S*)-**1** generally enhanced cytotoxicity. **2** is one of these compounds modified at A and B ring on C(20)(*S*)-**1** with polar groups<sup>6</sup>. Recently, we have developed a series of C(7)-substituted C(20)(*RS*)-camptothecin analogues<sup>7</sup> and C(20)(*RS*)-desethyl-20-substituted camptothecin analogues<sup>8</sup>. Among these derivatives, **3** and **4** showed significant antitumor activity compared with C(20)(*S*)-**1** in each series of derivatives.



Based on the our previous SAR studies, we expected the antitumor activity could be maximized by combinational modification of **2** and **3**, or **3** and **4**. In this paper, the syntheses and *in vitro* cytotoxicities of C(20)(*RS*)-camptothecin analogues modified at both B (or A) and E ring are reported.

The syntheses of the analogues modified at both B and E ring (**12** - **14**) were accomplished in 5 steps starting from diol **5** which could be easily prepared by previous procedure<sup>7,8</sup> (Scheme 1).


### Scheme 1



Reagents : i) Ac<sub>2</sub>O / CH<sub>2</sub>Cl<sub>2</sub> / pyridine / cat. DMAP., rt, 15 h (94%), ii) 80% TFA, rt, 4 h (89%), iii) **8**/cat. *p*-TsOH/toluene, reflux, 5 h (65%), iv) LiOH/MeOH/H<sub>2</sub>O, rt, 6 h ; then 1 N HCl (pH = 3), rt, 1 h (21%), v) MOMCl (or chloromethyl ethyl ether)/CH<sub>2</sub>Cl<sub>2</sub>/*i*-pr<sub>2</sub>NEt, 0°C to rt, 40 - 44 h (**10** (35%); **11** (52%)), vi) H<sub>2</sub>, 10% Pd on C / AcOH, rt, 5 - 20 h (**12** (31%); **13** (28%); **14** (34%)).

The diacetylation of **5** with Ac<sub>2</sub>O in pyridine gave **6**, which was converted to **7** by deketalization with 80% trifluoroacetic acid. The Friedländer condensation of **7** with **8**<sup>7</sup> followed by hydrolysis and relactonization provided **9**. The treatment of **9** with MOMCl or chloromethyl ethyl ether in basic condition gave **10** and **11** respectively. Finally, the catalytic hydrogenation of **9** - **11** in acetic acid solvent gave **12** - **14** respectively. The other series of analogues (**16** - **18**) were obtained from **7** (Scheme 2). The Friedländer condensation of **7** with amino ketone **15**<sup>6</sup> followed by hydrolysis and relactonization provided **16**. The ether formation was performed by the same method as Scheme 1 to get **17** and **18**.

### Scheme 2



Reagents: i) **15**/cat. *p*-TsOH/toluene, reflux, 15 h; then LiOH/MeOH/H<sub>2</sub>O, rt, 2 h; then 1 N HCl (pH = 3), rt, 3 h (38%), ii) MOMCl (or chloromethyl ethyl ether)/CH<sub>2</sub>Cl<sub>2</sub>/*i*-pr<sub>2</sub>NEt, 0°C to rt, 20 - 26 h (**17** (10%); **18** (9%)),

Table 1. *In vitro* Cytotoxicity<sup>9</sup> of Camptothecin Analogues against Human Tumor Cell Lines<sup>10</sup> (IC<sub>50</sub>,  $\mu$ M).

Compd.	A172	DLD-1	CAOV-3	KATO-III	L1210
C(20)(S)-1	0.029	0.102	0.032	0.448	0.035
<b>2</b>	0.054	2.783	0.087	6.113	14.004
<b>3</b>	2.620	1.860	0.560	0.320	0.440
<b>4</b>	0.167	0.276	0.004	0.623	0.460
<b>12</b>	3.560	0.619	0.483	4.400	6.730
<b>13</b>	12.170	0.549	0.091	3.270	— <sup>a</sup>
<b>14</b>	5.460	2.100	0.024	4.210	14.110
<b>16</b>	0.391	0.055	0.010	0.135	0.898
<b>17</b>	0.029	0.108	0.003	1.460	0.307
<b>18</b>	— <sup>a</sup>	3.876	3.762	5.917	— <sup>a</sup>

<sup>a</sup>The test was not performed.

*In vitro* cytotoxic activities against five human tumor cell lines for the above camptothecin analogues<sup>11</sup> along with C(20)(S)-1 are listed in Table 1. Generally, the both 7 and 20-substituted analogues (**12** - **14**) reduced the potency compared with **3** and **4**. This result was not consistent with our strategy to improve the cytotoxic activity by modification of both B and E ring in camptothecin. We suspect the both modification is not favorable in the binding process because of steric hindrance. On the other hand, A and E ring modified derivatives (**16** - **18**) which were fixed with relatively small chloromethyl group at C(7) showed comparable cytotoxicity with **2**, **3** and C(20)(S)-1. As shown in Table 1, **16** was 2 and 3 times more potent than C(20)(S)-1 in DLD-1 and KATO-III cell lines, respectively. Especially **16** and **17** showed 3- and 10-fold cytotoxicity in CAOV-3 cell line compared with C(20)(S)-1, respectively. In conclusion, the combinational modification of **2** and **3** enhanced the cytotoxicities compared with each **2**, **3** except **18**. Especially the 10 times higher cytotoxicity of **17** compared with C(20)(S)-1 in CAOV-3 cell line give us possible chance to develop specifically effective antitumor agent against ovarian cancer. Considering that these derivatives were racemate, optically active form of **16** and **17** would have even higher cytotoxicity. The study of chiral **16** and **17** is currently being investigated.

**Acknowledgement :** This research was supported by grants of the '96 Good Health R & D project from the Ministry of Health & Welfare, R. O. K. and the Research Center of New Drug Development of Seoul National University.

#### References and Notes :

1. Wall, M. E.; Wani, M. C.; Cook, C. E.; Palmer, K. H.; McPhail, A. T.; Sim, G. A. *J. Am. Chem. Soc.* **1966**, *88*, 3888.
2. Gottlieb, J. A.; Luce, J. K. *Cancer Chemother. Rep.* **1972**, *56*, 103.

3. Hsiang, Y.-H.; Hertzberg, R. P.; Hecht, S.; Liu, L. F. *J. Biol. Chem.* **1985**, *260*, 14873. Hsiang, Y.-H.; Liu, L. F.; Wall, M. E.; Wani, M. C.; Nicholas, A. W.; Manikumar, G.; Kirschenbaum, S.; Silber, R.; Potmesil, M. *Cancer Res.* **1989**, *49*, 4385.
4. Sawada, S.; Okajima, S.; Aiyama, R.; Nokada, K.; Furuta, T.; Yokokura, T.; Sugino, E.; Yamanouchi, K.; Miyasaka, T. *Chem. Pharm. Bull.* **1991**, *39*, 1446.
5. Kingsbury, W. D.; Boehm, T. C.; Jakas, D. R.; Holden, K. G.; Hecht, S. M.; Gallagher, G.; Caranfa, M. J.; McCabe, F. L.; Faucette, L. F.; Johnson, R. K.; Hertzberg, R. P. *J. Med. Chem.* **1991**, *34*, 98.
6. a) Luzzio, M. J.; Besterman, J. M.; Emerson, D. L.; Evans, M. G.; Lackey, K.; Leitner, P. L.; McIntyre, G.; Morton, B.; Myers, P. L.; Peel, M.; Sisco, J. M.; Sternbach, D. D.; Tong, W.-Q.; Truesdale, A.; Uehling, D. E.; Vuong, A.; Yates, J. *J. Med. Chem.* **1995**, *38*, 395. b) Lackey, K.; Sternbach, D. D.; Croom, D. K.; Ermerson, D. L.; Evans, M. G.; Leitner, P. L.; Luzzio, M. J.; McIntyre, G.; Vuong, A.; Yates, J.; Besterman, J. M. *J. Med. Chem.* **1996**, *39*, 713.
7. Jew, S. S.; Kim, H. J.; Kim, M. G.; Roh, E. Y.; Cho, Y. S.; Kim, J. K.; Cha, K. H.; Lee, K. K.; Han, H. J.; Choi, J. Y.; Lee, H. *Bioorg. & Med. Chem. Lett.* **1996**, *6*, 845.
8. Jew, S. S.; Kim, M. G.; Kim, H. J.; Roh, E. Y.; Cho, Y. S.; Kim, J. K.; Cha, K. H.; Lee, K. K.; Han, H. J.; Lee, H. *Bioorg. & Med. Chem. Lett.* **1996**, *6*, 849.
9. Skehan, P.; Storeng, R.; Scudiero, D.; Monks, A.; McMahon, J.; Vistica, D.; Warren, J. T.; Bokesch, H.; Kenny, S.; Boyd, M. R. *J. Natl. Canc. Inst.* **1990**, *82*, 1107.
10. *In vitro* antiproliferative activities of the analogues against five tumor cell lines (A172 human CNS cancer; DLD-1, human colon cancer; CAOV-3, human ovarian cancer; KATO-III, human gastric cancer; L1210, mouse leukemia) were measured by SRB assay<sup>9</sup> after 3 days of incubation and expressed as the doses required to inhibit the growth of 50 % of the cells cultivated (IC<sub>50</sub>,  $\mu$ M).
11. All new compounds gave satisfactory spectroscopic data consistent with the proposed structures.