

## SYNTHESES OF D- AND L-MYO-INOSITOL 1,2,4,5-TETRAKISPHOSPHATE AND STEREOSELECTIVITY OF THE I(1,4,5)P<sub>3</sub> RECEPTOR BINDING

Sung-Kee Chung<sup>\*a</sup>, Boo-Gyo Shin<sup>a</sup>, Young-Tae Chang<sup>a</sup>, Byung-Chang Suh<sup>b</sup> and Kyong-Tai Kim<sup>b</sup>




<sup>a</sup>Department of Chemistry and <sup>b</sup>Department of Life Sciences, Pohang University of Science & Technology,  
Pohang 790-784, Korea.

Received 15 December 1997; accepted 6 February 1998

**Abstract:** D- and L-*myo*-Inositol 1,2,4,5-tetrakisphosphate [D- & L-I(1,2,4,5)P<sub>4</sub>], which are analogues of D-*myo*-Inositol 1,4,5-trisphosphate [D-I(1,4,5)P<sub>3</sub>], a calcium mobilizing second messenger, were synthesized via resolution of the camphanate ester of a *myo*-inositol derivative, and the binding affinities to I(1,4,5)P<sub>3</sub> receptor were measured. © 1998 Elsevier Science Ltd. All rights reserved.


Since the discovery that D-*myo*-inositol-1,4,5-trisphosphate [I(1,4,5)P<sub>3</sub>, **1**] plays a pivotal role as a second messenger in the transmembrane signaling, thus mobilizing calcium ions from the intracellular storage, its interactions with the I(1,4,5)P<sub>3</sub> receptor and metabolic enzymes have been widely studied.<sup>1</sup> One of the major metabolic pathways involves a specific phosphorylation of I(1,4,5)P<sub>3</sub> to I(1,3,4,5)P<sub>4</sub> by I(1,4,5)P<sub>3</sub>-3-kinase [IP3K].<sup>2</sup> It has been suggested that I(1,3,4,5)P<sub>4</sub> also acts as a second messenger mediating the entry of extracellular Ca<sup>2+</sup> through a plasma membrane ion channel,<sup>3</sup> and to mobilize Ca<sup>2+</sup> also from the intracellular calcium stores, albeit less potently than I(1,4,5)P<sub>3</sub>.<sup>4</sup> A study with all possible regioisomers of IP<sub>4</sub>s<sup>5</sup> for their ability to bind to the IP<sub>3</sub> receptor in bovine adrenal cortical membranes, and also for their ability to mobilize Ca<sup>2+</sup> from IP<sub>3</sub>-sensitive Ca<sup>2+</sup> stores in permeabilized CHO cell, indicated that DL-I(1,2,4,5)P<sub>4</sub> had a binding affinity comparable to that of D-I(1,4,5)P<sub>3</sub>.<sup>6</sup>

The syntheses of unnatural I(1,2,4,5)P<sub>4</sub> were reported both in the racemic form<sup>7</sup> and in chiral D-form.<sup>8,9</sup> Racemic I(1,2,4,5)P<sub>4</sub> was found to be 2-3 times less potent than the natural ligand, I(1,4,5)P<sub>3</sub> in terms of the binding affinity and calcium release effect from intracellular calcium store,<sup>10</sup> whereas chiral D-I(1,2,4,5)P<sub>4</sub> (**2D**) was shown to possess the agonistic property only 1.5-2 times less potent than I(1,4,5)P<sub>3</sub>.<sup>9</sup>




As L-I(1,4,5)P<sub>3</sub> is known to be essentially inactive in its binding to the IP<sub>3</sub> receptor or calcium releasing ability,<sup>11</sup> L-I(1,2,4,5)P<sub>4</sub> has also been assumed to be an inactive component in the binding study with the racemic IP<sub>4</sub> sample: an assumption never experimentally confirmed. Here we wish to report the first synthesis of L-I(1,2,4,5)P<sub>4</sub> (**2L**) and its binding property to IP<sub>3</sub> receptor.

Racemic diol **3**<sup>12</sup> was resolved via the diastereomers of its (-)-camphanate ester (Scheme 1), **4D** and **4L**.<sup>13</sup> After silica gel column chromatography, each diastereomer **4D** and **4L** was treated with NaOMe in MeOH to give the enantiomeric pair, **3D** (mp 169–170 °C, [ $\alpha$ ]<sub>D</sub> - 41.7, c 1.58, CH<sub>2</sub>Cl<sub>2</sub>) and **3L** (mp 169–170 °C, [ $\alpha$ ]<sub>D</sub> + 40.2, c 1.21, CH<sub>2</sub>Cl<sub>2</sub>; + 25.7, c 0.69, CH<sub>3</sub>CN; lit.<sup>14</sup> mp 159–161 °C, [ $\alpha$ ]<sub>D</sub> + 22.0, c 1.08, CH<sub>3</sub>CN) (Scheme 1).



Scheme 1. a. (i) (1*S*)-(–)-camphanic chloride (Camp-Cl), pyridine. (ii) separation by column chromatography, **4D** (39%) is less polar than **4L** (35%). b. NaOMe, MeOH,  $\Delta$ , 84%.

Diol **3D** was benzylated under the conventional conditions employing BnBr and NaH in DMF to give **5D**<sup>15</sup>. Acid catalyzed hydrolysis of **5D** in aq. AcOH gave the tetraol, **6D**<sup>16</sup>. Compound **6D** was phosphorylated by successive treatments with dibenzyl *N,N*-diisopropylphosphoramidite and 1*H*-tetrazole, and then H<sub>2</sub>O<sub>2</sub> to give the protected D-I(1,2,4,5)P<sub>4</sub>, **7D**.<sup>17</sup> Hydrogenolysis of **7D** using Pd catalyst on activated charcoal followed by an addition of NaOH to adjust pH 10 gave the sodium salt of D-I(1,2,4,5)P<sub>4</sub>, **2D** (Scheme 2).<sup>18</sup> L-I(1,2,4,5)P<sub>4</sub>, **2L** was synthesized according to the same procedure.



Scheme 2. a. BnBr, NaH, DMF, 87%. b. acetic acid - water (80 : 20), reflux, 80%. c. (i) dibenzyl *N,N*-diisopropylphosphoramidite, 1*H*-tetrazole, (ii) 30% H<sub>2</sub>O<sub>2</sub>, 79%. d. (i) H<sub>2</sub>, Pd-C (10%). (ii) pH 10 (NaOH), quant.

The binding affinities of synthetic **2D** and **2L** were examined by the standard competition binding assay using 1.25 nM [<sup>3</sup>H]-D-I(1,4,5)P<sub>3</sub> and I(1,4,5)P<sub>3</sub> binding protein, which was prepared from bovine adrenal cortex.<sup>19</sup> With D-I(1,4,5)P<sub>3</sub> (IC<sub>50</sub> 15.3 nM) as the reference standard, **2D** showed a comparable binding affinity (IC<sub>50</sub> 13.4 nM) to the natural ligand, while **2L** revealed a much lower affinity (IC<sub>50</sub> 598 nM). It appears quite possible that even the low binding activity of **2L** (about 2% of **2D**) might be due to the contamination of **2D**, since the intermediate **4L** contained about 1.5% **4D**. Thus it is clear that the IP<sub>3</sub> receptor is quite stereospecific in its binding recognition.

In conclusion, we have prepared each enantiomer of I(1,2,4,5)P<sub>4</sub> and demonstrated that the D-form is indeed the active IP<sub>3</sub> receptor agonist as was previously assumed.

**Acknowledgement.** This work was supported by the Korea Science & Engineering Foundation/Center for Biofunctional Molecules and the Ministry of Education/Basic Science Research Institute Fund (97-3437).

#### References and Notes

- a) Majerus, P. W. *Annu. Rev. Biochem.* **1992**, *61*, 225-250. b) Berridge, M. J. *Nature* **1993**, *361*, 315-325.

2. Irvine, R. F.; Letcher, A. J.; Heslop, J. P.; Berridge, M. J. *Nature* **1986**, *320*, 631–634.
3. Irvine, R. F.; Cullen, P. J. *Current Biology* **1993**, *3*, 540–543.
4. Wilcocks, R. A.; Challiss, R. A. J.; Liu, C.; Potter, B. V. L.; Nahorski, S. R. *Mol. Pharmacol.* **1993**, *44*, 810–817.
5. Chung, S. K.; Chang, Y. T. *J. Chem. Soc., Chem. Commun.* **1995**, 11–12.
6. Burford, N. T.; Nahorski, S. R.; Chung, S. K.; Chang, Y. T.; Wilcox, R. A. *Cell Calcium* **1997**, *21*, 301–304.
7. a) Meek, J. L.; Davidson, F.; Hobbs, F. W. Jr. *J. Am. Chem. Soc.* **1988**, *110*, 2317–2318. b) Carless, H. A. J.; Busia, K. *Tetrahedron Lett.* **1990**, *31*, 3449–3452.
8. Chen, J.; Dorman, G.; Prestwich, G. D. *J. Org. Chem.* **1996**, *61*, 393–397.
9. Kozikowski, A. P.; Fauq, A. H.; Wilcox, R. A.; Nahorski, S. R. *Bioorg. Med. Chem. Lett.* **1995**, *5*, 1295–1300.
10. a) Mills, S. J.; Safrany, S. T.; Wilcox, R. A.; Nahorski, S. R.; Potter, B. V. L. *Bioorg. Med. Chem. Lett.* **1993**, *3*, 1505–1510. b) Hirata, M.; Narumoto, N.; Watanabe, Y.; Kanematsu, T.; Koga, T.; Ozaki, S. *Mol. Pharmacol.* **1994**, *45*, 271–276. c) Wilcox, R. A.; Safrany, S. T.; Lampe, D.; Mills, S. J.; Nahorski, S. R.; Potter, B. V. L. *Eur. J. Biochem.* **1994**, *223*, 115–124.
11. a) Polokoff, M. A.; Bencen, G. H.; Vacca, J. P.; DeSolms, S. J.; Young, S. D.; Huff, J. R. *J. Biol. Chem.* **1988**, *263*, 11922–11927. b) Taylor, C. W.; Berridge, M. J.; Brown, K. D.; Cooke, A. M.; Potter, B. V. L. *Biochem. Biophys. Res. Commun.* **1988**, *150*, 626–632. c) Strupish, J.; Cooke, A. M.; Potter, B. V. L.; Gigg, R.; Nahorski, S. R. *Biochem. J.* **1988**, *253*, 901–905.
12. Chung, S. K.; Ryu, Y. *Carbohydr. Res.* **1994**, *258*, 145–167.
13. **4D**: >99% de based on  $^1\text{H}$ -NMR,  $[\alpha]_{\text{D}} + 3.4$  (c 1.04,  $\text{CH}_2\text{Cl}_2$ ); **4L**: ca. 97% de.  $[\alpha]_{\text{D}} - 18.6$  (c 1.50,  $\text{CH}_2\text{Cl}_2$ ).  $R_{\text{f}}$  values on silica gel TLC (ethyl acetate :  $\text{CH}_2\text{Cl}_2 = 1 : 7$ ); **4D**: 0.52; **4L**: 0.44.
14. The assignments of **4D** and **4L** were based on the literature data for **3L**: Jones, M.; Rana, K. K.; Ward, J. G.; Young, R. C. *Tetrahedron Lett.* **1989**, *30*, 5353–5356.
15. **5D**: mp 155–156 °C,  $[\alpha]_{\text{D}} - 45.2$  (c 1.12,  $\text{CH}_2\text{Cl}_2$ ); **5L**: mp 155–156 °C,  $[\alpha]_{\text{D}} + 43.6$  (c 1.43,  $\text{CH}_2\text{Cl}_2$ ).
16. **6D**: mp 169–170 °C,  $[\alpha]_{\text{D}} + 14.7$  (c 1.03,  $\text{CH}_3\text{OH}$ ); **6L**: mp 169–170 °C,  $[\alpha]_{\text{D}} - 12.5$  (c 0.95,  $\text{CH}_3\text{OH}$ ).
17. **7D**: Oil,  $[\alpha]_{\text{D}} - 3.3$  (c 1.02,  $\text{CHCl}_3$ ); **7L**: Oil,  $[\alpha]_{\text{D}} + 2.8$  (c 1.35,  $\text{CHCl}_3$ );  $^{31}\text{P}$ -NMR ( $\text{CDCl}_3$ )  $\delta$  0.70, 1.05, 1.10, 1.61.
18. **2D**:  $[\alpha]_{\text{D}} - 13.3$  (c 1.0,  $\text{H}_2\text{O}$ , pH 10); **2L**:  $[\alpha]_{\text{D}} + 12.1$  (c 1.0,  $\text{H}_2\text{O}$ , pH 10).  $^{31}\text{P}$ -NMR ( $\text{D}_2\text{O}$ , pH 10)  $\delta$  4.52, 5.06, 5.28, 5.37.
19. Suh, B.-C.; Lee, C.-O.; Kim, K.-T. *J. Neurochem.* **1995**, *64*, 1071–1079.