

Statistical learning theory

Joris LIMONIER

September 24, 2022

Contents

1	Proposed exercise lecture 02	1
1.1	Home Exercise	1

1 Proposed exercise lecture 02

1.1 Home Exercise

Demonstrate that the MAP (Maximum A Posteriori) classification rule is quadratic with respect to x^* . We have the following:

$$\hat{y}^* = \arg \max_{k=1,\dots,k} \mathbb{P}(Y = k|X = x^*) \quad (1)$$

We compute the probability:

$$\begin{aligned} \arg \max_{k=1,\dots,k} \mathbb{P}(Y = k|X = x^*) &\stackrel{\text{Bayes}}{=} \frac{\mathbb{P}(Y = k)\mathbb{P}(X = x^*|Y = k)}{\mathbb{P}(X = x^*)} \\ &\propto \mathbb{P}(Y = k)\mathbb{P}(X = x^*|Y = k) \\ &\propto \hat{\Pi}_k \mathcal{N}(x^*, \hat{\mu}_k, \hat{\Sigma}_k) \end{aligned}$$

$$\begin{aligned}
\hat{y}^* &= \arg \max_{k=1,\dots,k} \hat{\Pi}_k \mathcal{N}(x^*, \hat{\mu}_k, \hat{\Sigma}_k) \\
&= \arg \max_{k=1,\dots,k} \log [\hat{\Pi}_k] + \log [\mathcal{N}(x^*, \hat{\mu}_k, \hat{\Sigma}_k)] \\
&= \arg \max_{k=1,\dots,k} \log [\hat{\Pi}_k] + \log \left[\frac{1}{(2\pi)^{N/2} |\Sigma|^{1/2}} e^{-\frac{1}{2} (x^* - \mu)^\top \Sigma^{-1} (x^* - \mu)} \right] \\
&= \arg \max_{k=1,\dots,k} \underbrace{\log [\hat{\Pi}_k]}_{\text{Constant w.r.t. } x^*} + \underbrace{\left[-\frac{1}{2} (x^* - \mu)^\top \Sigma^{-1} (x^* - \mu) \right]}_{\text{Quadratic in } x^*} \\
&\quad - \underbrace{\log [(2\pi)^{N/2} |\Sigma|^{1/2}]}_{\text{Constant w.r.t. } x^*}
\end{aligned}$$

The second term is Quadratic in x^* , the others are constant w.r.t. x^* , which completes the proof.