

Exercise 1

Chapter 3

1. **Monotonicity of Sample Complexity:** Let \mathcal{H} be a hypothesis class for a binary classification task. Suppose that \mathcal{H} is PAC learnable and its sample complexity is given by $m_{\mathcal{H}}(\cdot, \cdot)$. Show that $m_{\mathcal{H}}$ is monotonically nonincreasing in each of its parameters. That is, show that given $\delta \in (0, 1)$, and given $0 < \epsilon_1 \leq \epsilon_2 < 1$, we have that $m_{\mathcal{H}}(\epsilon_1, \delta) \geq m_{\mathcal{H}}(\epsilon_2, \delta)$. Similarly, show that given $\epsilon \in (0, 1)$, and given $0 < \delta_1 \leq \delta_2 < 1$, we have that $m_{\mathcal{H}}(\epsilon, \delta_1) \geq m_{\mathcal{H}}(\epsilon, \delta_2)$.
3. Let $\mathcal{X} = \mathbb{R}^2$, $\mathcal{Y} = \{0, 1\}$, and let \mathcal{H} be the class of concentric circles in the plane, that is, $\mathcal{H} = \{h_r : r \in \mathbb{R}_+\}$, where $h_r(x) = \mathbb{1}_{[\|x\| \leq r]}$. Prove that \mathcal{H} is PAC learnable (assume realizability), and its sample complexity is bounded by

$$m_{\mathcal{H}}(\epsilon, \delta) \leq \left\lceil \frac{\log(1/\delta)}{\epsilon} \right\rceil.$$

7. (*) **The Bayes optimal predictor:** Show that for every probability distribution \mathcal{D} , the Bayes optimal predictor $f_{\mathcal{D}}$ is optimal, in the sense that for every classifier g from \mathcal{X} to $\{0, 1\}$, $L_{\mathcal{D}}(f_{\mathcal{D}}) \leq L_{\mathcal{D}}(g)$.

Chapter 4

1. In this exercise, we show that the (ϵ, δ) requirement on the convergence of errors in our definitions of PAC learning, is, in fact, quite close to a simpler looking requirement about averages (or expectations). Prove that the following two statements are equivalent (for any learning algorithm A , any probability distribution \mathcal{D} , and any loss function whose range is $[0, 1]$):
 1. For every $\epsilon, \delta > 0$, there exists $m(\epsilon, \delta)$ such that $\forall m \geq m(\epsilon, \delta)$

$$\mathbb{P}_{S \sim \mathcal{D}^m} [L_{\mathcal{D}}(A(S)) > \epsilon] < \delta$$

2.

$$\lim_{m \rightarrow \infty} \mathbb{E}_{S \sim \mathcal{D}^m} [L_{\mathcal{D}}(A(S))] = 0$$

2. **Bounded loss functions:** In Corollary 4.6 we assumed that the range of the loss function is $[0, 1]$. Prove that if the range of the loss function is $[a, b]$ then the sample complexity satisfies

$$m_{\mathcal{H}}(\epsilon, \delta) \leq m_{\mathcal{H}}^{\text{uc}}(\epsilon/2, \delta) \leq \left\lceil \frac{2 \log(2|\mathcal{H}|/\delta)(b-a)^2}{\epsilon^2} \right\rceil.$$