


Lecture 2. Diffusion models

Introduction to Bayesian statistical Learning II

21.05.2025 Instructors: Alina Bazarova, Jose Robledo

Denoising Diffusion Probabilistic Models (DDPM)


Another type of generative models. What do they have to do with Bayes?



1. Forward diffusion process: gradually **adding noise** to samples

Denoising Diffusion Probabilistic Models (DDPM)


Another type of generative models. What do they have to do with Bayes?



1. Forward diffusion process: gradually **adding noise** to samples
2. Reverse diffusion process: **recreating** the sample **from noise**

Denoising Diffusion Probabilistic Models (DDPM)

Another type of generative models. What do they have to do with Bayes?



1. Forward diffusion process: gradually **adding noise** to samples
2. Reverse diffusion process: **recreating** the sample **from noise**
3. Heavily relying on conditional probability and **Bayes theorem** in particular

Forward diffusion process

Let $x_0 \sim q(x)$ be a real data distribution.

We produce a sequence of **noisy samples** x_1, \dots, x_T

Forward diffusion process

Let $x_0 \sim q(x)$ be a real data distribution.

We produce a sequence of **noisy samples** x_1, \dots, x_T

Noise steps controlled by a **variance schedule** $\{\beta_t \in (0,1)\}_{t=1}^T$

Forward diffusion process

Let $x_0 \sim q(x)$ be a real data distribution.

We produce a sequence of **noisy samples** x_1, \dots, x_T

Noise steps controlled by a **variance schedule** $\{\beta_t \in (0,1)\}_{t=1}^T$

$$q(x_t | x_{t-1}) \sim \mathcal{N}(x_t; \sqrt{1 - \beta_t} x_{t-1}, \beta_t \mathbf{I}), \quad q(x_{1:T} | x_0) = \prod_{t=1}^T q(x_t | x_{t-1})$$

Forward diffusion process

Let $x_0 \sim q(x)$ be a real data distribution.

We produce a sequence of **noisy samples** x_1, \dots, x_T

Noise steps controlled by a **variance schedule** $\{\beta_t \in (0,1)\}_{t=1}^T$

$$q(x_t | x_{t-1}) \sim \mathcal{N}(x_t; \sqrt{1 - \beta_t} x_{t-1}, \beta_t \mathbf{I}), \quad q(x_{1:T} | x_0) = \prod_{t=1}^T q(x_t | x_{t-1})$$

As $T \rightarrow \infty$, x_T is equivalent to a **Gaussian distribution**

Forward diffusion process

Can we sample x_t at any arbitrary time step t ? Yes!!!

The key: good old **reparametrisation trick!**

Forward diffusion process

Can we sample x_t at any arbitrary time step t ? Yes!!!

The key: good old **reparametrisation trick!**

Let $\alpha_t = 1 - \beta_t$ and $\bar{\alpha}_t = \prod_{i=1}^t \alpha_i$

Forward diffusion process

Can we sample x_t at any arbitrary time step t ? Yes!!!

The key: good old **reparametrisation trick!**

$$\text{Let } \alpha_t = 1 - \beta_t \text{ and } \bar{\alpha}_t = \prod_{i=1}^t \alpha_i$$

$$x_t = \sqrt{\alpha_t} x_{t-1} + \sqrt{1 - \alpha_t} \varepsilon_{t-1} = \sqrt{\alpha_t \alpha_{t-1}} x_{t-2} + \sqrt{1 - \alpha_t \alpha_{t-1}} \bar{\varepsilon}_{t-2}$$

Forward diffusion process

Can we sample x_t at any arbitrary time step t ? Yes!!!

The key: good old **reparametrisation trick!**

$$\text{Let } \alpha_t = 1 - \beta_t \text{ and } \bar{\alpha}_t = \prod_{i=1}^t \alpha_i$$

$$x_t = \sqrt{\bar{\alpha}_t} x_{t-1} + \sqrt{1 - \alpha_t} \varepsilon_{t-1} = \sqrt{\alpha_t \bar{\alpha}_{t-1}} x_{t-2} + \sqrt{1 - \alpha_t \bar{\alpha}_{t-1}} \bar{\varepsilon}_{t-2}$$

$$\text{NB: } \sqrt{\alpha_t} \sqrt{1 - \alpha_t} \varepsilon_{t-1} + \sqrt{1 - \alpha_{t-1}} \varepsilon_{t-2}, \quad \varepsilon_i \sim \mathcal{N}(0, \mathbf{I}), \text{ hence}$$

Forward diffusion process

Can we sample x_t at any arbitrary time step t ? Yes!!!

The key: good old **reparametrisation trick!**

$$\text{Let } \alpha_t = 1 - \beta_t \text{ and } \bar{\alpha}_t = \prod_{i=1}^t \alpha_i$$

$$x_t = \sqrt{\alpha_t}x_{t-1} + \sqrt{1 - \alpha_t}\varepsilon_{t-1} = \sqrt{\alpha_t\alpha_{t-1}}x_{t-2} + \sqrt{1 - \alpha_t\alpha_{t-1}}\bar{\varepsilon}_{t-2}$$

$$\text{NB: } \sqrt{\alpha_t}\sqrt{1 - \alpha_t}\varepsilon_{t-1} + \sqrt{1 - \alpha_{t-1}}\varepsilon_{t-2}, \quad \varepsilon_i \sim \mathcal{N}(0, \mathbf{I}), \text{ hence}$$

$$\sqrt{\alpha_{t-1}}\sqrt{1 - \alpha_t}\varepsilon_{t-1} + \sqrt{1 - \alpha_{t-1}}\varepsilon_{t-2} \sim \mathcal{N}(0, \alpha_{t-1} - \alpha_t\alpha_{t-1} + 1 - \alpha_{t-1})$$

Forward diffusion process

Can we sample x_t at any arbitrary time step t ? Yes!!!

The key: good old **reparametrisation trick!**

$$\text{Let } \alpha_t = 1 - \beta_t \text{ and } \bar{\alpha}_t = \prod_{i=1}^t \alpha_i$$

$$x_t = \sqrt{\alpha_t} x_{t-1} + \sqrt{1 - \alpha_t} \varepsilon_{t-1} = \sqrt{\alpha_t \alpha_{t-1}} x_{t-2} + \sqrt{1 - \alpha_t \alpha_{t-1}} \bar{\varepsilon}_{t-2}$$

$$\text{NB: } \sqrt{\alpha_t} \sqrt{1 - \alpha_t} \varepsilon_{t-1} + \sqrt{1 - \alpha_{t-1}} \varepsilon_{t-2}, \quad \varepsilon_i \sim \mathcal{N}(0, \mathbf{I}), \text{ hence}$$

$$\sqrt{\alpha_{t-1}} \sqrt{1 - \alpha_t} \varepsilon_{t-1} + \sqrt{1 - \alpha_{t-1}} \varepsilon_{t-2} \sim \mathcal{N}(0, \cancel{\alpha_{t-1}} - \alpha_t \alpha_{t-1} + 1 - \cancel{\alpha_{t-1}})$$

Forward diffusion process

Can we sample x_t at any arbitrary time step t ? Yes!!!

The key: good old **reparametrisation trick!**

$$\text{Let } \alpha_t = 1 - \beta_t \text{ and } \bar{\alpha}_t = \prod_{i=1}^t \alpha_i$$

$$x_t = \sqrt{\alpha_t}x_{t-1} + \sqrt{1 - \alpha_t}\varepsilon_{t-1} = \sqrt{\alpha_t\alpha_{t-1}}x_{t-2} + \sqrt{1 - \alpha_t\alpha_{t-1}}\bar{\varepsilon}_{t-2}$$

$$\text{NB: } \sqrt{\alpha_t}\sqrt{1 - \alpha_t}\varepsilon_{t-1} + \sqrt{1 - \alpha_{t-1}}\varepsilon_{t-2}, \quad \varepsilon_i \sim \mathcal{N}(0, \mathbf{I}), \text{ hence}$$

$$\sqrt{\alpha_{t-1}}\sqrt{1 - \alpha_t}\varepsilon_{t-1} + \sqrt{1 - \alpha_{t-1}}\varepsilon_{t-2} \sim \mathcal{N}(0, \alpha_{t-1} - \alpha_t\alpha_{t-1} + 1 - \alpha_{t-1})$$

$$\sqrt{\alpha_{t-1}}\sqrt{1 - \alpha_t}\varepsilon_{t-1} + \sqrt{1 - \alpha_{t-1}}\varepsilon_{t-2} \sim \sqrt{1 - \alpha_t\alpha_{t-1}}\mathcal{N}(0, \mathbf{I})$$

Forward diffusion process

Following the same argument

$$x_t = \sqrt{\alpha_t}x_{t-1} + \sqrt{1 - \alpha_t}\varepsilon_{t-1} = \dots = \sqrt{\bar{\alpha}_t}x_0 + \sqrt{1 - \bar{\alpha}_t}\varepsilon, \quad \varepsilon \sim \mathcal{N}(0, \mathbf{I}), \quad \bar{\alpha}_t = \prod_{i=1}^t \alpha_i$$

Forward diffusion process

Following the same argument

$$x_t\sqrt{\alpha_t}x_{t-1} + \sqrt{1 - \alpha_t}\varepsilon_{t-1} = \dots = \sqrt{\bar{\alpha}_t}x_0 + \sqrt{1 - \bar{\alpha}_t}\varepsilon, \quad \varepsilon \sim \mathcal{N}(0, \mathbf{I}), \quad \bar{\alpha}_t = \prod_{i=1}^t \alpha_i$$

Hence $q(x_t | x_0) \sim \mathcal{N}(x_t; \sqrt{\bar{\alpha}_t}x_0, (1 - \bar{\alpha}_t)\mathbf{I})$

Forward diffusion process

Following the same argument

$$x_t\sqrt{\alpha_t}x_{t-1} + \sqrt{1 - \alpha_t}\varepsilon_{t-1} = \dots = \sqrt{\bar{\alpha}_t}x_0 + \sqrt{1 - \bar{\alpha}_t}\varepsilon, \quad \varepsilon \sim \mathcal{N}(0, \mathbf{I}), \quad \bar{\alpha}_t = \prod_{i=1}^t \alpha_i$$

Hence $q(x_t | x_0) \sim \mathcal{N}(x_t; \sqrt{\bar{\alpha}_t}x_0, (1 - \bar{\alpha}_t)\mathbf{I})$

Larger update step when the sample gets noisier: $\beta_1 < \beta_2 < \dots < \beta_T$,

and therefore $\bar{\alpha}_1 > \dots > \bar{\alpha}_T$

Reverse diffusion process

If we reverse the above process and sample from $q(x_{t-1} | x_t)$,

we can recreate a sample from $x_T \sim \mathcal{N}(0, \mathbf{I})$

We need to estimate $q(x_{t-1} | x_t)$. We do it with another probability density function p_θ

Reverse diffusion process

If we reverse the above process and sample from $q(x_{t-1} | x_t)$,

we can recreate a sample from $x_T \sim \mathcal{N}(0, \mathbf{I})$

We need to estimate $q(x_{t-1} | x_t)$. We do it with another probability density function p_θ

$$p_\theta(x_{0:T}) = p(x_T) \prod_{t=1}^T p_\theta(x_{t-1} | x_t),$$

Reverse diffusion process

If we reverse the above process and sample from $q(x_{t-1} | x_t)$,

we can recreate a sample from $x_T \sim \mathcal{N}(0, \mathbf{I})$

We need to estimate $q(x_{t-1} | x_t)$. We do it with another probability density function p_θ

$$p_\theta(x_{0:T}) = p(x_T) \prod_{t=1}^T p_\theta(x_{t-1} | x_t), \quad p_\theta(x_{t-1} | x_t) = \mathcal{N}(x_{t-1}; \mu_\theta(x_t, t), \Sigma_\theta(x_t, t))$$

Which is learned via a neural network (θ are the parameters of NN)!

Reverse diffusion process

If we reverse the above process and sample from $q(x_{t-1} | x_t)$,

we can recreate a sample from $x_T \sim \mathcal{N}(0, \mathbf{I})$

We need to estimate $q(x_{t-1} | x_t)$. We do it with another probability density function p_θ

$$p_\theta(x_{0:T}) = p(x_T) \prod_{t=1}^T p_\theta(x_{t-1} | x_t), \quad p_\theta(x_{t-1} | x_t) = \mathcal{N}(x_{t-1}; \mu_\theta(x_t, t), \Sigma_\theta(x_t, t))$$

Which is learned via a neural network (θ are the parameters of NN)!

Important: $q(x_{t-1} | x_t, x_0) \sim \mathcal{N}(x_{t-1}; \tilde{\mu}(x_t, x_0), \tilde{\beta}\mathbf{I})$ **tractable!**

Conditioning on x_0 . Bayes rule to the rescue! $P(A | B) = \frac{P(B)P(B | A)}{P(A)}$

Conditioning on x_0 . Bayes rule to the rescue! $P(A | B) = \frac{P(B)P(B | A)}{P(A)}$

$$P(A | B, C) = \frac{P(AB | C)}{P(B | C)} = \frac{P(B | A, C)P(A | C)}{P(B | C)}$$

Conditioning on x_0 . Bayes rule to the rescue! $P(A | B) = \frac{P(B)P(B | A)}{P(A)}$

$$P(A | B, C) = \frac{P(AB | C)}{P(B | C)} = \frac{P(B | A, C)P(A | C)}{P(B | C)}$$

$$q(x_{t-1} | x_t, x_0) = q(x_t | x_{t-1}, x_0) \frac{q(x_{t-1} | x_0)}{q(x_t | x_0)} \propto \exp\left(-\frac{1}{2}\left(\frac{(x_t - \sqrt{\alpha_t}x_{t-1})^2}{1 - \alpha_t} + \frac{(x_{t-1} - \sqrt{\bar{\alpha}_{t-1}}x_0)^2}{1 - \bar{\alpha}_{t-1}} - \frac{(x_t - \sqrt{\bar{\alpha}_t}x_0)^2}{1 - \bar{\alpha}_t}\right)\right)$$

Conditioning on x_0 . Bayes rule to the rescue! $P(A | B) = \frac{P(B)P(B | A)}{P(A)}$

$$P(A | B, C) = \frac{P(AB | C)}{P(B | C)} = \frac{P(B | A, C)P(A | C)}{P(B | C)}$$

$$\begin{aligned} q(x_{t-1} | x_t, x_0) &= q(x_t | x_{t-1}, x_0) \frac{q(x_{t-1} | x_0)}{q(x_t | x_0)} \propto \exp\left(-\frac{1}{2}\left(\frac{(x_t - \sqrt{\alpha_t}x_{t-1})^2}{1 - \alpha_t} + \frac{(x_{t-1} - \sqrt{\bar{\alpha}_{t-1}}x_0)^2}{1 - \bar{\alpha}_{t-1}} - \frac{(x_t - \sqrt{\bar{\alpha}_t}x_0)^2}{1 - \bar{\alpha}_t}\right)\right) \\ &= \exp\left(-\frac{1}{2}\left(\left(\frac{\alpha_t}{\beta_t} + \frac{1}{1 - \bar{\alpha}_{t-1}}\right)x_{t-1}^2 - 2\left(\frac{\sqrt{\alpha_t}}{\beta_t}x_t + \frac{\sqrt{\bar{\alpha}_{t-1}}}{1 - \bar{\alpha}_{t-1}}x_0\right)x_{t-1} + C(x_0, x_1)\right)\right) \end{aligned}$$

Conditioning on x_0 . Bayes rule to the rescue! $P(A | B) = \frac{P(B)P(B | A)}{P(A)}$

$$P(A | B, C) = \frac{P(AB | C)}{P(B | C)} = \frac{P(B | A, C)P(A | C)}{P(B | C)}$$

$$\begin{aligned} q(x_{t-1} | x_t, x_0) &= q(x_t | x_{t-1}, x_0) \frac{q(x_{t-1} | x_0)}{q(x_t | x_0)} \propto \exp\left(-\frac{1}{2}\left(\frac{(x_t - \sqrt{\alpha_t}x_{t-1})^2}{1 - \alpha_t} + \frac{(x_{t-1} - \sqrt{\bar{\alpha}_{t-1}}x_0)^2}{1 - \bar{\alpha}_{t-1}} - \frac{(x_t - \sqrt{\bar{\alpha}_t}x_0)^2}{1 - \bar{\alpha}_t}\right)\right) \\ &= \exp\left(-\frac{1}{2}\left(\left(\frac{\alpha_t}{\beta_t} + \frac{1}{1 - \bar{\alpha}_{t-1}}\right)x_{t-1}^2 - 2\left(\frac{\sqrt{\alpha_t}}{\beta_t}x_t + \frac{\sqrt{\bar{\alpha}_{t-1}}}{1 - \bar{\alpha}_{t-1}}x_0\right)x_{t-1} + C(x_0, x_1)\right)\right) \end{aligned}$$

$$\text{Hence } \tilde{\beta}_t = 1/\left(\frac{\alpha_t}{\beta_t} + \frac{1}{1 - \bar{\alpha}_{t-1}}\right) = \frac{1 - \bar{\alpha}_{t-1}}{1 - \bar{\alpha}_t}\beta_t,$$

Conditioning on x_0 . Bayes rule to the rescue! $P(A | B) = \frac{P(B)P(B | A)}{P(A)}$

$$P(A | B, C) = \frac{P(AB | C)}{P(B | C)} = \frac{P(B | A, C)P(A | C)}{P(B | C)}$$

$$\begin{aligned} q(x_{t-1} | x_t, x_0) &= q(x_t | x_{t-1}, x_0) \frac{q(x_{t-1} | x_0)}{q(x_t | x_0)} \propto \exp\left(-\frac{1}{2}\left(\frac{(x_t - \sqrt{\alpha_t}x_{t-1})^2}{1 - \alpha_t} + \frac{(x_{t-1} - \sqrt{\bar{\alpha}_{t-1}}x_0)^2}{1 - \bar{\alpha}_{t-1}} - \frac{(x_t - \sqrt{\bar{\alpha}_t}x_0)^2}{1 - \bar{\alpha}_t}\right)\right) \\ &= \exp\left(-\frac{1}{2}\left(\left(\frac{\alpha_t}{\beta_t} + \frac{1}{1 - \bar{\alpha}_{t-1}}\right)x_{t-1}^2 - 2\left(\frac{\sqrt{\alpha_t}}{\beta_t}x_t + \frac{\sqrt{\bar{\alpha}_{t-1}}}{1 - \bar{\alpha}_{t-1}}x_0\right)x_{t-1} + C(x_0, x_1)\right)\right) \end{aligned}$$

$$\text{Hence } \tilde{\beta}_t = 1/\left(\frac{\alpha_t}{\beta_t} + \frac{1}{1 - \bar{\alpha}_{t-1}}\right) = \frac{1 - \bar{\alpha}_{t-1}}{1 - \bar{\alpha}_t} \beta_t,$$

$$\tilde{\mu}_t = \left(\frac{\sqrt{\alpha_t}}{\beta_t}x_t + \frac{\sqrt{\bar{\alpha}_{t-1}}}{1 - \bar{\alpha}_{t-1}}x_0\right)/\tilde{\beta}_t = \left(\frac{\sqrt{\alpha_t}}{\beta_t}x_t + \frac{\sqrt{\bar{\alpha}_{t-1}}}{1 - \bar{\alpha}_{t-1}}x_0\right)\frac{1 - \bar{\alpha}_{t-1}}{1 - \bar{\alpha}_t}\beta_t$$

Conditioning on x_0 . Bayes rule to the rescue! $P(A | B) = \frac{P(B)P(B | A)}{P(A)}$

$$P(A | B, C) = \frac{P(AB | C)}{P(B | C)} = \frac{P(B | A, C)P(A | C)}{P(B | C)}$$

$$q(x_{t-1} | x_t, x_0) = q(x_t | x_{t-1}, x_0) \frac{q(x_{t-1} | x_0)}{q(x_t | x_0)} \propto \exp\left(-\frac{1}{2}\left(\frac{(x_t - \sqrt{\alpha_t}x_{t-1})^2}{1 - \alpha_t} + \frac{(x_{t-1} - \sqrt{\bar{\alpha}_{t-1}}x_0)^2}{1 - \bar{\alpha}_{t-1}} - \frac{(x_t - \sqrt{\bar{\alpha}_t}x_0)^2}{1 - \bar{\alpha}_t}\right)\right)$$

$$= \exp\left(-\frac{1}{2}\left(\left(\frac{\alpha_t}{\beta_t} + \frac{1}{1 - \bar{\alpha}_{t-1}}\right)x_{t-1}^2 - 2\left(\frac{\sqrt{\alpha_t}}{\beta_t}x_t + \frac{\sqrt{\bar{\alpha}_{t-1}}}{1 - \bar{\alpha}_{t-1}}x_0\right)x_{t-1} + C(x_0, x_1)\right)\right)$$

$$\text{Hence } \tilde{\beta}_t = 1/\left(\frac{\alpha_t}{\beta_t} + \frac{1}{1 - \bar{\alpha}_{t-1}}\right) = \frac{1 - \bar{\alpha}_{t-1}}{1 - \bar{\alpha}_t}\beta_t,$$

$$\tilde{\mu}_t = \left(\frac{\sqrt{\alpha_t}}{\beta_t}x_t + \frac{\sqrt{\bar{\alpha}_{t-1}}}{1 - \bar{\alpha}_{t-1}}x_0\right)/\tilde{\beta}_t = \left(\frac{\sqrt{\alpha_t}}{\beta_t}x_t + \frac{\sqrt{\bar{\alpha}_{t-1}}}{1 - \bar{\alpha}_{t-1}}x_0\right)\frac{1 - \bar{\alpha}_{t-1}}{1 - \bar{\alpha}_t}\beta_t$$

We can reparametrise it further! \rightarrow

$$= \frac{\sqrt{\alpha_t}(1 - \bar{\alpha}_{t-1})}{1 - \bar{\alpha}_t}x_t + \frac{\sqrt{\bar{\alpha}_{t-1}}\beta_t}{1 - \bar{\alpha}_t}x_0$$

Remember! $x_t = \sqrt{\bar{\alpha}_t}x_0 + \sqrt{1 - \bar{\alpha}_t}\varepsilon,$ $x_0 = \frac{1}{\sqrt{\bar{\alpha}_t}}(x_t - \sqrt{1 - \bar{\alpha}_t}\varepsilon_t)$

Remember! $x_t = \sqrt{\bar{\alpha}_t}x_0 + \sqrt{1 - \bar{\alpha}_t}\varepsilon,$ $x_0 = \frac{1}{\sqrt{\bar{\alpha}_t}}(x_t - \sqrt{1 - \bar{\alpha}_t}\varepsilon_t)$

Substitute this into expression of $\tilde{\mu}_t$ and get $\tilde{\mu}_t = \frac{1}{\sqrt{\alpha_t}}(x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}}\varepsilon_t)$

Remember! $x_t = \sqrt{\bar{\alpha}_t}x_0 + \sqrt{1 - \bar{\alpha}_t}\varepsilon,$ $x_0 = \frac{1}{\sqrt{\bar{\alpha}_t}}(x_t - \sqrt{1 - \bar{\alpha}_t}\varepsilon_t)$

Substitute this into expression of $\tilde{\mu}_t$ and get $\tilde{\mu}_t = \frac{1}{\sqrt{\alpha_t}}(x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}}\varepsilon_t)$

Recap! We need to learn the distributions $p_{\theta}(x_{t-1} | x_t) \sim \mathcal{N}(x_{t-1}; \mu_{\theta}(x_t, t), \Sigma_{\theta}(x_t, t))$

So we will train μ_{θ} to predict $\tilde{\mu}_t$

Remember! $x_t = \sqrt{\bar{\alpha}_t}x_0 + \sqrt{1 - \bar{\alpha}_t}\varepsilon,$ $x_0 = \frac{1}{\sqrt{\bar{\alpha}_t}}(x_t - \sqrt{1 - \bar{\alpha}_t}\varepsilon_t)$

Substitute this into expression of $\tilde{\mu}_t$ and get $\tilde{\mu}_t = \frac{1}{\sqrt{\alpha_t}}(x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}}\varepsilon_t)$

Recap! We need to learn the distributions $p_{\theta}(x_{t-1} | x_t) \sim \mathcal{N}(x_{t-1}; \mu_{\theta}(x_t, t), \Sigma_{\theta}(x_t, t))$

So we will train μ_{θ} to predict $\tilde{\mu}_t$

How are we going to do that?

Remember! $x_t = \sqrt{\bar{\alpha}_t}x_0 + \sqrt{1 - \bar{\alpha}_t}\varepsilon$, $x_0 = \frac{1}{\sqrt{\bar{\alpha}_t}}(x_t - \sqrt{1 - \bar{\alpha}_t}\varepsilon_t)$

Substitute this into expression of $\tilde{\mu}_t$ and get $\tilde{\mu}_t = \frac{1}{\sqrt{\alpha_t}}(x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}}\varepsilon_t)$

Recap! We need to learn the distributions $p_{\theta}(x_{t-1} | x_t) \sim \mathcal{N}(x_{t-1}; \mu_{\theta}(x_t, t), \Sigma_{\theta}(x_t, t))$

So we will train μ_{θ} to predict $\tilde{\mu}_t$

How are we going to do that? Minimise negative log-likelihood $-\log p_{\theta}(x_0)$

Remember! $x_t = \sqrt{\bar{\alpha}_t}x_0 + \sqrt{1 - \bar{\alpha}_t}\varepsilon,$ $x_0 = \frac{1}{\sqrt{\bar{\alpha}_t}}(x_t - \sqrt{1 - \bar{\alpha}_t}\varepsilon_t)$

Substitute this into expression of $\tilde{\mu}_t$ and get $\tilde{\mu}_t = \frac{1}{\sqrt{\alpha_t}}(x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}}\varepsilon_t)$

Recap! We need to learn the distributions $p_{\theta}(x_{t-1} | x_t) \sim \mathcal{N}(x_{t-1}; \mu_{\theta}(x_t, t), \Sigma_{\theta}(x_t, t))$

So we will train μ_{θ} to predict $\tilde{\mu}_t$

How are we going to do that? Minimise negative log-likelihood $-\log p_{\theta}(x_0)$

$$-\log p_{\theta}(x_0) \leq -\log p_{\theta}(x_0) + D_{KL}(q(x_{1:T} | x_0) || p_{\theta}(x_{1:T} | x_0))$$

Remember! $x_t = \sqrt{\bar{\alpha}_t}x_0 + \sqrt{1 - \bar{\alpha}_t}\varepsilon,$ $x_0 = \frac{1}{\sqrt{\bar{\alpha}_t}}(x_t - \sqrt{1 - \bar{\alpha}_t}\varepsilon_t)$

Substitute this into expression of $\tilde{\mu}_t$ and get $\tilde{\mu}_t = \frac{1}{\sqrt{\alpha_t}}(x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}}\varepsilon_t)$

Recap! We need to learn the distributions $p_\theta(x_{t-1} | x_t) \sim \mathcal{N}(x_{t-1}; \mu_\theta(x_t, t), \Sigma_\theta(x_t, t))$

So we will train μ_θ to predict $\tilde{\mu}_t$

How are we going to do that? Minimise negative log-likelihood $-\log p_\theta(x_0)$

Kullback-Leibler divergence ≥ 0

$-\log p_\theta(x_0) \leq -\log p_\theta(x_0) + D_{KL}(q(x_{1:T} | x_0) || p_\theta(x_{1:T} | x_0))$

Remember! $x_t = \sqrt{\bar{\alpha}_t}x_0 + \sqrt{1 - \bar{\alpha}_t}\varepsilon, \quad x_0 = \frac{1}{\sqrt{\bar{\alpha}_t}}(x_t - \sqrt{1 - \bar{\alpha}_t}\varepsilon_t)$

Substitute this into expression of $\tilde{\mu}_t$ and get $\tilde{\mu}_t = \frac{1}{\sqrt{\alpha_t}}(x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}}\varepsilon_t)$

Recap! We need to learn the distributions $p_\theta(x_{t-1} | x_t) \sim \mathcal{N}(x_{t-1}; \mu_\theta(x_t, t), \Sigma_\theta(x_t, t))$

So we will train μ_θ to predict $\tilde{\mu}_t$

How are we going to do that? Minimise negative log-likelihood $-\log p_\theta(x_0)$

Kullback-Leibler divergence ≥ 0

$-\log p_\theta(x_0) \leq -\log p_\theta(x_0) + D_{KL}(q(x_{1:T} | x_0) || p_\theta(x_{1:T} | x_0)) =$

$-\log p_\theta(x_0) + E_{x_{1:T} \sim q(x_{1:T} | x_0)}[\log \frac{q(x_{1:T} | x_0)}{p_\theta(x_{0:T})/p_\theta(x_0)}] = -\log p_\theta(x_0) + E_q[\log \frac{q(x_{1:T} | x_0)}{p_\theta(x_{0:T})}] + \log p_\theta(x_0)$

Remember! $x_t = \sqrt{\bar{\alpha}_t}x_0 + \sqrt{1 - \bar{\alpha}_t}\varepsilon, \quad x_0 = \frac{1}{\sqrt{\bar{\alpha}_t}}(x_t - \sqrt{1 - \bar{\alpha}_t}\varepsilon_t)$

Substitute this into expression of $\tilde{\mu}_t$ and get $\tilde{\mu}_t = \frac{1}{\sqrt{\alpha_t}}(x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}}\varepsilon_t)$

Recap! We need to learn the distributions $p_\theta(x_{t-1} | x_t) \sim \mathcal{N}(x_{t-1}; \mu_\theta(x_t, t), \Sigma_\theta(x_t, t))$

So we will train μ_θ to predict $\tilde{\mu}_t$

How are we going to do that? Minimise negative log-likelihood $-\log p_\theta(x_0)$

Kullback-Leibler divergence ≥ 0



$$-\log p_\theta(x_0) \leq -\log p_\theta(x_0) + D_{KL}(q(x_{1:T}|x_0) || p_\theta(x_{1:T}|x_0)) =$$

$$-\log p_\theta(x_0) + E_{x_{1:T} \sim q(x_{1:T}|x_0)}[\log \frac{q(x_{1:T}|x_0)}{p_\theta(x_{0:T})/p_\theta(x_0)}] = -\cancel{\log p_\theta(x_0)} + E_q[\log \frac{q(x_{1:T}|x_0)}{p_\theta(x_{0:T})}] + \cancel{\log p_\theta(x_0)} = E_q[\log \frac{q(x_{1:T}|x_0)}{p_\theta(x_{0:T})}]$$

$$\text{Let } L_{VLB} = E_{q(x_{0:T})}[\log \frac{q(x_{1:T}|x_0)}{p_\theta(x_{0:T})}] \geq -E_{q(x_0)} \log p_\theta(x_0)$$

$$\text{Let } L_{VLB} = E_{q(x_{0:T})}[\log \frac{q(x_{1:T}|x_0)}{p_\theta(x_{0:T})}] \geq -E_{q(x_0)} \log p_\theta(x_0)$$

We used that $E_{q(x_0)}E_{q(x_{1:T}|x_0)}f(x_{0:T}) = E_{q(x_{0:T})}f(x_{0:T})$ for any f of our interest

$$\text{Let } L_{VLB} = E_{q(x_{0:T})}[\log \frac{q(x_{1:T}|x_0)}{p_\theta(x_{0:T})}] \geq -E_{q(x_0)}p_\theta(x_0)$$

We used that $E_{q(x_0)}E_{q(x_{1:T}|x_0)}f(x_{0:T}) = E_{q(x_{0:T})}f(x_{0:T})$ for any f of our interest

$$L_{VLB} = E_{q(x_{0:T})}[\log \frac{q(x_{1:T}|x_0)}{p_\theta(x_{0:T})}]$$

$$\text{Let } L_{VLB} = E_{q(x_{0:T})}[\log \frac{q(x_{1:T}|x_0)}{p_\theta(x_{0:T})}] \geq - E_{q(x_0)} p_\theta(x_0)$$

We used that $E_{q(x_0)} E_{q(x_{1:T}|x_0)} f(x_{0:T}) = E_{q(x_{0:T})} f(x_{0:T})$ for any f of our interest

$$L_{VLB} = E_{q(x_{0:T})}[\log \frac{q(x_{1:T}|x_0)}{p_\theta(x_{0:T})}] = E_q[\log \frac{\prod_{t=1}^T q(x_t|x_{t-1})}{p_\theta(x_T) \prod_{t=1}^T p_\theta(x_{t-1}|x_t)}]$$

$$\text{Let } L_{VLB} = E_{q(x_{0:T})}[\log \frac{q(x_{1:T}|x_0)}{p_\theta(x_{0:T})}] \geq -E_{q(x_0)}p_\theta(x_0)$$

We used that $E_{q(x_0)}E_{q(x_{1:T}|x_0)}f(x_{0:T}) = E_{q(x_{0:T})}f(x_{0:T})$ for any f of our interest

$$L_{VLB} = E_{q(x_{0:T})}[\log \frac{q(x_{1:T}|x_0)}{p_\theta(x_{0:T})}] = E_q[\log \frac{\prod_{t=1}^T q(x_t|x_{t-1})}{p_\theta(x_T)\prod_{t=1}^T p_\theta(x_{t-1}|x_t)}] = E_q[-\log p_\theta(x_T) + \sum_{t=1}^T \log \frac{q(x_t|x_{t-1})}{p_\theta(x_{t-1}|x_t)}]$$

Let $L_{VLB} = E_{q(x_{0:T})}[\log \frac{q(x_{1:T} | x_0)}{p_\theta(x_{0:T})}] \geq -E_{q(x_0)} p_\theta(x_0)$

We used that $E_{q(x_0)} E_{q(x_{1:T} | x_0)} f(x_{0:T}) = E_{q(x_{0:T})} f(x_{0:T})$ for any f of our interest

$$\begin{aligned} L_{VLB} &= E_{q(x_{0:T})}[\log \frac{q(x_{1:T} | x_0)}{p_\theta(x_{0:T})}] = E_q[\log \frac{\prod_{t=1}^T q(x_t | x_{t-1})}{p_\theta(x_T) \prod_{t=1}^T p_\theta(x_{t-1} | x_t)}] = E_q[-\log p_\theta(x_T) + \sum_{t=1}^T \log \frac{q(x_t | x_{t-1})}{p_\theta(x_{t-1} | x_t)}] \\ &= E_q[-\log p_\theta(x_T) + \sum_{t=2}^T \log \frac{q(x_t | x_{t-1})}{p_\theta(x_{t-1} | x_t)} + \log \frac{q(x_1 | x_0)}{p_\theta(x_0 | x_1)}] \end{aligned}$$

$$L_{VLB} = E_{q(x_{0:T})}[\log \frac{q(x_{1:T}|x_0)}{p_\theta(x_{0:T})}] = E_q[\log \frac{\prod_{t=1}^T q(x_t|x_{t-1})}{p_\theta(x_T) \prod_{t=1}^T p_\theta(x_{t-1}|x_t)}] = E_q[-\log p_\theta(x_T) + \sum_{t=1}^T \log \frac{q(x_t|x_{t-1})}{p_\theta(x_{t-1}|x_t)}]$$

$$\begin{aligned}
L_{VLB} &= E_{q(x_{0:T})}[\log \frac{q(x_{1:T}|x_0)}{p_\theta(x_{0:T})}] = E_q[\log \frac{\prod_{t=1}^T q(x_t|x_{t-1})}{p_\theta(x_T) \prod_{t=1}^T p_\theta(x_{t-1}|x_t)}] = E_q[-\log p_\theta(x_T) + \sum_{t=1}^T \log \frac{q(x_t|x_{t-1})}{p_\theta(x_{t-1}|x_t)}] \\
&= E_q[-\log p_\theta(x_T) + \sum_{t=2}^T \log \frac{q(x_t|x_{t-1})}{p_\theta(x_{t-1}|x_t)} + \log \frac{q(x_1|x_0)}{p_\theta(x_0|x_1)}]
\end{aligned}$$

$$\begin{aligned}
L_{VLB} &= E_{q(x_{0:T})}[\log \frac{q(x_{1:T}|x_0)}{p_\theta(x_{0:T})}] = E_q[\log \frac{\prod_{t=1}^T q(x_t|x_{t-1})}{p_\theta(x_T) \prod_{t=1}^T p_\theta(x_{t-1}|x_t)}] = E_q[-\log p_\theta(x_T) + \sum_{t=1}^T \log \frac{q(x_t|x_{t-1})}{p_\theta(x_{t-1}|x_t)}] \\
&= E_q[-\log p_\theta(x_T) + \sum_{t=2}^T \log \frac{q(x_t|x_{t-1})}{p_\theta(x_{t-1}|x_t)} + \log \frac{q(x_1|x_0)}{p_\theta(x_0|x_1)}] = E_q[-\log p_\theta(x_T) + \sum_{t=2}^T \log \frac{q(x_{t-1}|x_t, x_0)}{p_\theta(x_{t-1}|x_t)} \frac{q(x_t|x_0)}{q(x_{t-1}|x_0)} + \dots]
\end{aligned}$$

$$\begin{aligned}
L_{VLB} &= E_{q(x_{0:T})}[\log \frac{q(x_{1:T}|x_0)}{p_\theta(x_{0:T})}] = E_q[\log \frac{\prod_{t=1}^T q(x_t|x_{t-1})}{p_\theta(x_T) \prod_{t=1}^T p_\theta(x_{t-1}|x_t)}] = E_q[-\log p_\theta(x_T) + \sum_{t=1}^T \log \frac{q(x_t|x_{t-1})}{p_\theta(x_{t-1}|x_t)}] \\
&= E_q[-\log p_\theta(x_T) + \sum_{t=2}^T \log \frac{q(x_t|x_{t-1})}{p_\theta(x_{t-1}|x_t)} + \log \frac{q(x_1|x_0)}{p_\theta(x_0|x_1)}] = E_q[-\log p_\theta(x_T) + \sum_{t=2}^T \log \frac{q(x_{t-1}|x_t, x_0)}{p_\theta(x_{t-1}|x_t)} \frac{q(x_t|x_0)}{q(x_{t-1}|x_0)} + \dots] \\
&= E_q[-\log p_\theta(x_T) + \sum_{t=2}^T \log \frac{q(x_{t-1}|x_t, x_0)}{p_\theta(x_{t-1}|x_t)} + \sum_{t=2}^T \frac{q(x_t|x_0)}{q(x_{t-1}|x_0)} + \log \frac{q(x_1|x_0)}{p_\theta(x_0|x_1)}]
\end{aligned}$$

$$\begin{aligned}
L_{VLB} &= E_{q(x_{0:T})}[\log \frac{q(x_{1:T}|x_0)}{p_\theta(x_{0:T})}] = E_q[\log \frac{\prod_{t=1}^T q(x_t|x_{t-1})}{p_\theta(x_T) \prod_{t=1}^T p_\theta(x_{t-1}|x_t)}] = E_q[-\log p_\theta(x_T) + \sum_{t=1}^T \log \frac{q(x_t|x_{t-1})}{p_\theta(x_{t-1}|x_t)}] \\
&= E_q[-\log p_\theta(x_T) + \sum_{t=2}^T \log \frac{q(x_t|x_{t-1})}{p_\theta(x_{t-1}|x_t)} + \log \frac{q(x_1|x_0)}{p_\theta(x_0|x_1)}] = E_q[-\log p_\theta(x_T) + \sum_{t=2}^T \log \frac{q(x_{t-1}|x_t, x_0)}{p_\theta(x_{t-1}|x_t)} \frac{q(x_t|x_0)}{q(x_{t-1}|x_0)} + \dots] \\
&= E_q[-\log p_\theta(x_T) + \sum_{t=2}^T \log \frac{q(x_{t-1}|x_t, x_0)}{p_\theta(x_{t-1}|x_t)} + \sum_{t=2}^T \log \frac{q(x_t|x_0)}{q(x_{t-1}|x_0)} + \log \frac{q(x_1|x_0)}{p_\theta(x_0|x_1)}] \\
&= E_q[-\log p_\theta(x_T) + \sum_{t=2}^T \log \frac{q(x_{t-1}|x_t, x_0)}{p_\theta(x_{t-1}|x_t)} + \log \frac{q(x_T|x_0)}{q(x_1|x_0)} + \log \frac{q(x_1|x_0)}{p_\theta(x_0|x_1)}]
\end{aligned}$$

$$\begin{aligned}
L_{VLB} &= E_{q(x_{0:T})}[\log \frac{q(x_{1:T}|x_0)}{p_\theta(x_{0:T})}] = E_q[\log \frac{\prod_{t=1}^T q(x_t|x_{t-1})}{p_\theta(x_T) \prod_{t=1}^T p_\theta(x_{t-1}|x_t)}] = E_q[-\log p_\theta(x_T) + \sum_{t=1}^T \log \frac{q(x_t|x_{t-1})}{p_\theta(x_{t-1}|x_t)}] \\
&= E_q[-\log p_\theta(x_T) + \sum_{t=2}^T \log \frac{q(x_t|x_{t-1})}{p_\theta(x_{t-1}|x_t)} + \log \frac{q(x_1|x_0)}{p_\theta(x_0|x_1)}] = E_q[-\log p_\theta(x_T) + \sum_{t=2}^T \log \frac{q(x_{t-1}|x_t, x_0)}{p_\theta(x_{t-1}|x_t)} \frac{q(x_t|x_0)}{q(x_{t-1}|x_0)} + \dots] \\
&= E_q[-\log p_\theta(x_T) + \sum_{t=2}^T \log \frac{q(x_{t-1}|x_t, x_0)}{p_\theta(x_{t-1}|x_t)} + \sum_{t=2}^T \log \frac{q(x_t|x_0)}{q(x_{t-1}|x_0)} + \log \frac{q(x_1|x_0)}{p_\theta(x_0|x_1)}] \\
&= E_q[-\log p_\theta(x_T) + \sum_{t=2}^T \log \frac{q(x_{t-1}|x_t, x_0)}{p_\theta(x_{t-1}|x_t)} + \log \frac{q(x_T|x_0)}{q(\cancel{x_1}|x_0)} + \log \frac{q(\cancel{x_1}|x_0)}{p_\theta(x_0|x_1)}]
\end{aligned}$$

$$\begin{aligned}
L_{VLB} &= E_{q(x_{0:T})}[\log \frac{q(x_{1:T}|x_0)}{p_\theta(x_{0:T})}] = E_q[\log \frac{\prod_{t=1}^T q(x_t|x_{t-1})}{p_\theta(x_T) \prod_{t=1}^T p_\theta(x_{t-1}|x_t)}] = E_q[-\log p_\theta(x_T) + \sum_{t=1}^T \log \frac{q(x_t|x_{t-1})}{p_\theta(x_{t-1}|x_t)}] \\
&= E_q[-\log p_\theta(x_T) + \sum_{t=2}^T \log \frac{q(x_t|x_{t-1})}{p_\theta(x_{t-1}|x_t)} + \log \frac{q(x_1|x_0)}{p_\theta(x_0|x_1)}] = E_q[-\log p_\theta(x_T) + \sum_{t=2}^T \log \frac{q(x_{t-1}|x_t, x_0)}{p_\theta(x_{t-1}|x_t)} \frac{q(x_t|x_0)}{q(x_{t-1}|x_0)} + \dots] \\
&= E_q[-\log p_\theta(x_T) + \sum_{t=2}^T \log \frac{q(x_{t-1}|x_t, x_0)}{p_\theta(x_{t-1}|x_t)} + \sum_{t=2}^T \log \frac{q(x_t|x_0)}{q(x_{t-1}|x_0)} + \log \frac{q(x_1|x_0)}{p_\theta(x_0|x_1)}] \\
&= E_q[-\log p_\theta(x_T) + \sum_{t=2}^T \log \frac{q(x_{t-1}|x_t, x_0)}{p_\theta(x_{t-1}|x_t)} + \log \frac{q(x_T|x_0)}{q(\cancel{x_1}|x_0)} + \log \frac{q(\cancel{x_1}|x_0)}{p_\theta(x_0|x_1)}] \\
&= E_q[\log \frac{q(x_T|x_0)}{p_\theta(x_T)} + \sum_{t=2}^T \log \frac{q(x_{t-1}|x_t, x_0)}{p_\theta(x_{t-1}|x_t)} - \log p_\theta(x_0|x_1)]
\end{aligned}$$

$$\begin{aligned}
L_{VLB} &= E_{q(x_{0:T})}[\log \frac{q(x_{1:T}|x_0)}{p_\theta(x_{0:T})}] = E_q[\log \frac{\prod_{t=1}^T q(x_t|x_{t-1})}{p_\theta(x_T) \prod_{t=1}^T p_\theta(x_{t-1}|x_t)}] = E_q[-\log p_\theta(x_T) + \sum_{t=1}^T \log \frac{q(x_t|x_{t-1})}{p_\theta(x_{t-1}|x_t)}] \\
&= E_q[-\log p_\theta(x_T) + \sum_{t=2}^T \log \frac{q(x_t|x_{t-1})}{p_\theta(x_{t-1}|x_t)} + \log \frac{q(x_1|x_0)}{p_\theta(x_0|x_1)}] = E_q[-\log p_\theta(x_T) + \sum_{t=2}^T \log \frac{q(x_{t-1}|x_t, x_0)}{p_\theta(x_{t-1}|x_t)} \frac{q(x_t|x_0)}{q(x_{t-1}|x_0)} + \dots] \\
&= E_q[-\log p_\theta(x_T) + \sum_{t=2}^T \log \frac{q(x_{t-1}|x_t, x_0)}{p_\theta(x_{t-1}|x_t)} + \sum_{t=2}^T \frac{q(x_t|x_0)}{q(x_{t-1}|x_0)} + \log \frac{q(x_1|x_0)}{p_\theta(x_0|x_1)}] \\
&= E_q[-\log p_\theta(x_T) + \sum_{t=2}^T \log \frac{q(x_{t-1}|x_t, x_0)}{p_\theta(x_{t-1}|x_t)} + \log \frac{q(x_T|x_0)}{q(\cancel{x_1}|x_0)} + \log \frac{q(\cancel{x_1}|x_0)}{p_\theta(x_0|x_1)}] \\
&= E_q[\log \frac{q(x_T|x_0)}{p_\theta(x_T)} + \sum_{t=2}^T \log \frac{q(x_{t-1}|x_t, x_0)}{p_\theta(x_{t-1}|x_t)} - \log p_\theta(x_0|x_1)] \\
&= E_q[\underbrace{D_{KL}(q(x_T|x_0) || p_\theta(x_T))}_{L_T} + \sum_{t=2}^T \underbrace{D_{KL}(q(x_{t-1}|x_t, x_0) || p_\theta(x_{t-1}|x_t))}_{L_{t-1}} - \underbrace{\log p_\theta(x_0|x_1)}_{L_0}]
\end{aligned}$$

Simplifying the loss further

$$L_{VLB} = L_T + L_{T-1} + \dots + L_0$$

Simplifying the loss further

$$L_{VLB} = L_T + L_{T-1} + \dots + L_0$$

$$L_T = D_{KL}(q(x_T|x_0) || p_{\theta}(x_T)) \quad \text{Constant, since no trainable parameters and } x_T \text{ is Gaussian noise}$$

Simplifying the loss further

$$L_{VLB} = L_T + L_{T-1} + \dots + L_0$$

$$L_T = D_{KL}(q(x_T | x_0) || p_{\theta}(x_T)) \quad \text{Constant, since no trainable parameters and } x_T \text{ is Gaussian noise}$$

$$L_t = D_{KL}(q(x_{t-1} | x_t, x_0) || p_{\theta}(x_{t-1} | x_t)) \quad \text{The main thing!}$$

Simplifying the loss further

$$L_{VLB} = L_T + L_{T-1} + \dots + L_0$$

$$L_T = D_{KL}(q(x_T|x_0) || p_{\theta}(x_T)) \quad \text{Constant, since no trainable parameters and } x_T \text{ is Gaussian noise}$$

$$L_t = D_{KL}(q(x_{t-1}|x_t, x_0) || p_{\theta}(x_{t-1}|x_t)) \quad \text{The main thing!}$$

$$L_0 = -\log p_{\theta}(x_0|x_1) \quad \text{Can be omitted or modelled via a separate decoder, derived from}$$

$$\mathcal{N}(x_0; \mu_{\theta}(x_1, 1), \Sigma_{\theta}(x_1, 1))$$

Simplifying the loss further

$$L_{VLB} = L_T + L_{T-1} + \dots + L_0$$

$$L_T = D_{KL}(q(x_T | x_0) || p_{\theta}(x_T)) \quad \text{Constant, since no trainable parameters and } x_T \text{ is Gaussian noise}$$

$$L_t = D_{KL}(q(x_{t-1} | x_t, x_0) || p_{\theta}(x_{t-1} | x_t)) \quad \text{The main thing!}$$

$$L_0 = -\log p_{\theta}(x_0 | x_1) \quad \text{Can be omitted or modelled via a separate decoder, derived from}$$

$$\mathcal{N}(x_0; \mu_{\theta}(x_1, 1), \Sigma_{\theta}(x_1, 1))$$

Remember again! We need to learn the distributions $p_{\theta}(x_{t-1} | x_t) \sim \mathcal{N}(x_{t-1}; \mu_{\theta}(x_t, t), \Sigma_{\theta}(x_t, t))$

That is for L_t !

Parametrisation of L_t for Training Loss

$$p_{\theta}(x_{t-1} | x_t) \sim \mathcal{N}(x_{t-1}; \mu_{\theta}(x_t, t), \Sigma_{\theta}(x_t, t))$$

We would like to train μ_{θ} to predict $\tilde{\mu}_t = \frac{1}{\sqrt{\alpha_t}}(x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}}\epsilon_t)$

Parametrisation of L_t for Training Loss

$$p_{\theta}(x_{t-1} | x_t) \sim \mathcal{N}(x_{t-1}; \mu_{\theta}(x_t, t), \Sigma_{\theta}(x_t, t))$$

We would like to train μ_{θ} to predict $\tilde{\mu}_t = \frac{1}{\sqrt{\alpha_t}}(x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}}\epsilon_t)$

But x_t is available during training time!

Parametrisation of L_t for Training Loss

$$p_{\theta}(x_{t-1} | x_t) \sim \mathcal{N}(x_{t-1}; \mu_{\theta}(x_t, t), \Sigma_{\theta}(x_t, t))$$

We would like to train μ_{θ} to predict $\tilde{\mu}_t = \frac{1}{\sqrt{\alpha_t}}(x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}}\epsilon_t)$

But x_t is available during training time!

Hence, let us **predict the noise term instead!**

Parametrisation of L_t for Training Loss

$$p_{\theta}(x_{t-1} | x_t) \sim \mathcal{N}(x_{t-1}; \mu_{\theta}(x_t, t), \Sigma_{\theta}(x_t, t))$$

We would like to train μ_{θ} to predict $\tilde{\mu}_t = \frac{1}{\sqrt{\alpha_t}}(x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}}\varepsilon_t)$

But x_t is available during training time!

Hence, let us **predict the noise term instead!**

$$\mu_{\theta}(x_t, t) = \frac{1}{\sqrt{\alpha_t}}(x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}}\varepsilon_{\theta}(x_t, t))$$

Parametrisation of L_t for Training Loss

$$p_{\theta}(x_{t-1} | x_t) \sim \mathcal{N}(x_{t-1}; \mu_{\theta}(x_t, t), \Sigma_{\theta}(x_t, t))$$

We would like to train μ_{θ} to predict $\tilde{\mu}_t = \frac{1}{\sqrt{\alpha_t}}(x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}}\varepsilon_t)$

But x_t is available during training time!

Hence, let us **predict the noise term instead!**

$$\mu_{\theta}(x_t, t) = \frac{1}{\sqrt{\alpha_t}}(x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}}\varepsilon_{\theta}(x_t, t)) \quad x_{t-1} \sim \mathcal{N}(x_{t-1}; \frac{1}{\sqrt{\alpha_t}}(x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}}\varepsilon_{\theta}(x_t, t)), \Sigma_{\theta}(x_t, t))$$

Parametrisation of L_t for Training Loss

$$L_t = E_{x_0, \varepsilon} \left[\frac{1}{2 ||\Sigma_{\theta}(x_t, t)||_2^2} ||\tilde{\mu}_t(x_t, x_0) - \mu_{\theta}(x_t, t)||^2 \right]$$

Parametrisation of L_t for Training Loss

$$\begin{aligned} L_t &= E_{x_0, \varepsilon} \left[\frac{1}{2 ||\Sigma_{\theta}(x_t, t)||_2^2} ||\tilde{\mu}_t(x_t, x_0) - \mu_{\theta}(x_t, t)||^2 \right] \\ &= E_{x_0, \varepsilon} \left[\frac{1}{2 ||\Sigma_{\theta}||_2^2} || \frac{1}{\sqrt{\alpha_t}} (x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}} \varepsilon_t) - \frac{1}{\sqrt{\alpha_t}} (x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}} \varepsilon_{\theta}(x_t, t)) ||^2 \right] \end{aligned}$$

Parametrisation of L_t for Training Loss

$$\begin{aligned} L_t &= E_{x_0, \varepsilon} \left[\frac{1}{2 ||\Sigma_{\theta}(x_t, t)||_2^2} ||\tilde{\mu}_t(x_t, x_0) - \mu_{\theta}(x_t, t)||^2 \right] \\ &= E_{x_0, \varepsilon} \left[\frac{1}{2 ||\Sigma_{\theta}||_2^2} || \cancel{\frac{1}{\sqrt{\alpha_t}} x_t} - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}} \varepsilon_t - \cancel{\frac{1}{\sqrt{\alpha_t}} x_t} + \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}} \varepsilon_{\theta}(x_t, t) ||^2 \right] \\ &= E_{x_0, \varepsilon} \left[\frac{(1 - \alpha_t)^2}{2 \alpha_t (1 - \bar{\alpha}_t) ||\Sigma_{\theta}||_2^2} ||\varepsilon_t - \varepsilon_{\theta}(x_t, t)||^2 \right] \end{aligned}$$

Parametrisation of L_t for Training Loss

$$\begin{aligned} L_t &= E_{x_0, \varepsilon} \left[\frac{1}{2 ||\Sigma_\theta(x_t, t)||_2^2} ||\tilde{\mu}_t(x_t, x_0) - \mu_\theta(x_t, t)||^2 \right] \\ &= E_{x_0, \varepsilon} \left[\frac{1}{2 ||\Sigma_\theta||_2^2} || \frac{1}{\sqrt{\alpha_t}} (x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}} \varepsilon_t) - \frac{1}{\sqrt{\alpha_t}} (x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}} \varepsilon_\theta(x_t, t)) ||^2 \right] \\ &= E_{x_0, \varepsilon} \left[\frac{(1 - \alpha_t)^2}{2\alpha_t(1 - \bar{\alpha}_t) ||\Sigma_\theta||_2^2} ||\varepsilon_t - \varepsilon_\theta(x_t, t)||^2 \right] = E_{x_0, \varepsilon} \left[\frac{(1 - \alpha_t)^2}{2\alpha_t(1 - \bar{\alpha}_t) ||\Sigma_\theta||_2^2} ||\varepsilon_t - \varepsilon_\theta(\sqrt{\bar{\alpha}_t}x_0 + \sqrt{1 - \bar{\alpha}_t}\varepsilon_t, t)||^2 \right] \end{aligned}$$

Parametrisation of L_t for Training Loss

$$\begin{aligned} L_t &= E_{x_0, \varepsilon} \left[\frac{1}{2 ||\Sigma_{\theta}(x_t, t)||_2^2} ||\tilde{\mu}_t(x_t, x_0) - \mu_{\theta}(x_t, t)||^2 \right] \\ &= E_{x_0, \varepsilon} \left[\frac{1}{2 ||\Sigma_{\theta}||_2^2} || \frac{1}{\sqrt{\alpha_t}} (x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}} \varepsilon_t) - \frac{1}{\sqrt{\alpha_t}} (x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}} \varepsilon_{\theta}(x_t, t)) ||^2 \right] \\ &= E_{x_0, \varepsilon} \left[\frac{(1 - \alpha_t)^2}{2\alpha_t(1 - \bar{\alpha}_t) ||\Sigma_{\theta}||_2^2} ||\varepsilon_t - \varepsilon_{\theta}(x_t, t)||^2 \right] = E_{x_0, \varepsilon} \left[\frac{(1 - \alpha_t)^2}{2\alpha_t(1 - \bar{\alpha}_t) ||\Sigma_{\theta}||_2^2} ||\varepsilon_t - \varepsilon_{\theta}(\sqrt{\bar{\alpha}_t}x_0 + \sqrt{1 - \bar{\alpha}_t}\varepsilon_t, t)||^2 \right] \end{aligned}$$

In practice can simplify even further!

Parametrisation of L_t for Training Loss

$$\begin{aligned} L_t &= E_{x_0, \varepsilon} \left[\frac{1}{2 ||\Sigma_\theta(x_t, t)||_2^2} ||\tilde{\mu}_t(x_t, x_0) - \mu_\theta(x_t, t)||^2 \right] \\ &= E_{x_0, \varepsilon} \left[\frac{1}{2 ||\Sigma_\theta||_2^2} || \frac{1}{\sqrt{\alpha_t}} (x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}} \varepsilon_t) - \frac{1}{\sqrt{\alpha_t}} (x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}} \varepsilon_\theta(x_t, t)) ||^2 \right] \\ &= E_{x_0, \varepsilon} \left[\frac{(1 - \alpha_t)^2}{2\alpha_t(1 - \bar{\alpha}_t) ||\Sigma_\theta||_2^2} ||\varepsilon_t - \varepsilon_\theta(x_t, t)||^2 \right] = E_{x_0, \varepsilon} \left[\frac{(1 - \alpha_t)^2}{2\alpha_t(1 - \bar{\alpha}_t) ||\Sigma_\theta||_2^2} ||\varepsilon_t - \varepsilon_\theta(\sqrt{\bar{\alpha}_t}x_0 + \sqrt{1 - \bar{\alpha}_t}\varepsilon_t, t)||^2 \right] \end{aligned}$$

In practice can simplify even further!

$$L_t^{simple} = E_{t, x_0, \varepsilon} [||\varepsilon_t - \varepsilon_\theta(\sqrt{\bar{\alpha}_t}x_0 + \sqrt{1 - \bar{\alpha}_t}\varepsilon_t, t)||^2]$$

Parametrisation of L_t for Training Loss

$$\begin{aligned} L_t &= E_{x_0, \varepsilon} \left[\frac{1}{2 || \Sigma_{\theta}(x_t, t) ||_2^2} || \tilde{\mu}_t(x_t, x_0) - \mu_{\theta}(x_t, t) ||^2 \right] \\ &= E_{x_0, \varepsilon} \left[\frac{1}{2 || \Sigma_{\theta} ||_2^2} || \frac{1}{\sqrt{\alpha_t}} (x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}} \varepsilon_t) - \frac{1}{\sqrt{\alpha_t}} (x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}} \varepsilon_{\theta}(x_t, t)) ||^2 \right] \\ &= E_{x_0, \varepsilon} \left[\frac{(1 - \alpha_t)^2}{2 \alpha_t (1 - \bar{\alpha}_t) || \Sigma_{\theta} ||_2^2} || \varepsilon_t - \varepsilon_{\theta}(x_t, t) ||^2 \right] = E_{x_0, \varepsilon} \left[\frac{(1 - \alpha_t)^2}{2 \alpha_t (1 - \bar{\alpha}_t) || \Sigma_{\theta} ||_2^2} || \varepsilon_t - \varepsilon_{\theta}(\sqrt{\bar{\alpha}_t} x_0 + \sqrt{1 - \bar{\alpha}_t} \varepsilon_t, t) ||^2 \right] \end{aligned}$$

In practice can simplify even further!

$$L_t^{simple} = E_{t, x_0, \varepsilon} [|| \varepsilon_t - \varepsilon_{\theta}(\sqrt{\bar{\alpha}_t} x_0 + \sqrt{1 - \bar{\alpha}_t} \varepsilon_t, t) ||^2]$$


Bottom line: what we are doing is predicting the noise!

Parametrisation of L_t for Training Loss

$$\begin{aligned}
 L_t &= E_{x_0, \epsilon} \left[\frac{1}{2 ||\Sigma_{\theta}(x_t, t)||_2^2} ||\tilde{\mu}_t(x_t, x_0) - \mu_{\theta}(x_t, t)||^2 \right] \\
 &= E_{x_0, \epsilon} \left[\frac{1}{2 ||\Sigma_{\theta}||_2^2} || \frac{1}{\sqrt{\alpha_t}} (x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}} \epsilon_t) - \frac{1}{\sqrt{\alpha_t}} (x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}} \epsilon_{\theta}(x_t, t)) ||^2 \right] \\
 &= E_{x_0, \epsilon} \left[\frac{(1 - \alpha_t)^2}{2\alpha_t(1 - \bar{\alpha}_t) ||\Sigma_{\theta}||_2^2} ||\epsilon_t - \epsilon_{\theta}(x_t, t)||^2 \right] = E_{x_0, \epsilon} \left[\frac{(1 - \alpha_t)^2}{2\alpha_t(1 - \bar{\alpha}_t) ||\Sigma_{\theta}||_2^2} ||\epsilon_t - \epsilon_{\theta}(\sqrt{\bar{\alpha}_t}x_0 + \sqrt{1 - \bar{\alpha}_t}\epsilon_t, t)||^2 \right]
 \end{aligned}$$

In practice can simplify even further!

$$L_t^{simple} = E_{t, x_0, \epsilon} [||\epsilon_t - \epsilon_{\theta}(\sqrt{\bar{\alpha}_t}x_0 + \sqrt{1 - \bar{\alpha}_t}\epsilon_t, t)||^2]$$



Bottom line: what we are doing is predicting the noise! Unet architecture is used for that

Parametrization of β_t

Typically a sequence of linearly increasing constants: e.g. from $\beta_1 = 10^{-4}$ to $\beta_T = 0.02$

Parametrization of β_t

Typically a sequence of linearly increasing constants: e.g. from $\beta_1 = 10^{-4}$ to $\beta_T = 0.02$

Can be fixed (not learnable)

Parametrization of β_t

Typically a sequence of linearly increasing constants: e.g. from $\beta_1 = 10^{-4}$ to $\beta_T = 0.02$

Can be fixed (not learnable)

Parametrization of reverse process variance Σ_θ

Parametrization of β_t

Typically a sequence of linearly increasing constants: e.g. from $\beta_1 = 10^{-4}$ to $\beta_T = 0.02$

Can be fixed (not learnable)

Parametrization of reverse process variance Σ_θ

Can also be fixed $\Sigma_\theta(x_t, t) = \sigma_t^2 \mathbf{I}$, where σ_t is set to be a function of β_t

Parametrization of β_t

Typically a sequence of linearly increasing constants: e.g. from $\beta_1 = 10^{-4}$ to $\beta_T = 0.02$

Can be fixed (not learnable)

Parametrization of reverse process variance Σ_θ

Can also be fixed $\Sigma_\theta(x_t, t) = \sigma_t^2 \mathbf{I}$, where σ_t is set to be a function of β_t

Alternative $\Sigma_\theta(x_t, t) = \exp(v \log \beta_t + (1 - v) \log \tilde{\beta}_t)$, $\tilde{\beta}_t = \frac{1 - \bar{\alpha}_{t-1}}{1 - \bar{\alpha}_t}$, v is learnable

Parametrization of β_t

Typically a sequence of linearly increasing constants: e.g. from $\beta_1 = 10^{-4}$ to $\beta_T = 0.02$

Can be fixed (not learnable)

Parametrization of reverse process variance Σ_θ

Can also be fixed $\Sigma_\theta(x_t, t) = \sigma_t^2 \mathbf{I}$, where σ_t is set to be a function of β_t

Alternative $\Sigma_\theta(x_t, t) = \exp(v \log \beta_t + (1 - v) \log \tilde{\beta}_t)$, $\tilde{\beta}_t = \frac{1 - \bar{\alpha}_{t-1}}{1 - \bar{\alpha}_t}$, v is learnable

Hence, the loss $L = L_{simple} + \lambda L_{VLB}$, λ is small ~ 0.001 and L_{VLB} only guides the training of Σ_θ

(In L_{VLB} stop gradient with respect to μ_θ)

Bonus! Score networks and guided diffusion

The score of each sample's x probability density function is defined as $\nabla_x \log q(x)$

Bonus! Score networks and guided diffusion

The score of each sample's x probability density function is defined as $\nabla_x \log q(x)$

Langevin dynamics can sample data points from a probability density distribution using only the score $\nabla_x \log q(x)$ in an iterative process.

Bonus! Score networks and guided diffusion

The score of each sample's x probability density function is defined as $\nabla_x \log q(x)$

Langevin dynamics can sample data points from a probability density distribution using only the score $\nabla_x \log q(x)$ in an iterative process.

Score network $s_\theta(x_t, t) \approx \nabla_x \log q(x)$

Bonus! Score networks and guided diffusion

The score of each sample's x probability density function is defined as $\nabla_x \log q(x)$

Langevin dynamics can sample data points from a probability density distribution using only the score $\nabla_x \log q(x)$ in an iterative process.

Score network $s_\theta(x_t, t) \approx \nabla_x \log q(x)$

$$\text{If } x \sim \mathcal{N}(\mu, \sigma^2 \mathbf{I}), \quad \nabla_x \log p(x) = \nabla_x \left(-\frac{1}{2\sigma^2} (x - \mu)^2 \right) = -\frac{x - \mu}{\sigma^2} = -\frac{\varepsilon}{\sigma}, \quad \varepsilon \sim \mathcal{N}(0, \mathbf{I})$$

Bonus! Score networks and guided diffusion

The score of each sample's x probability density function is defined as $\nabla_x \log q(x)$

Langevin dynamics can sample data points from a probability density distribution using only the score $\nabla_x \log q(x)$ in an iterative process.

Score network $s_\theta(x_t, t) \approx \nabla_x \log q(x)$

$$\text{If } x \sim \mathcal{N}(\mu, \sigma^2 \mathbf{I}), \quad \nabla_x \log p(x) = \nabla_x \left(-\frac{1}{2\sigma^2} (x - \mu)^2 \right) = -\frac{x - \mu}{\sigma^2} = -\frac{\varepsilon}{\sigma}, \quad \varepsilon \sim \mathcal{N}(0, \mathbf{I})$$

$$q(x_t | x_0) \sim \mathcal{N}(\sqrt{\bar{\alpha}_t} x_0, (1 - \bar{\alpha}_t) \mathbf{I})$$

Bonus! Score networks and guided diffusion

The score of each sample's x probability density function is defined as $\nabla_x \log q(x)$

Langevin dynamics can sample data points from a probability density distribution using only the score $\nabla_x \log q(x)$ in an iterative process.

Score network $s_\theta(x_t, t) \approx \nabla_x \log q(x)$

$$\text{If } x \sim \mathcal{N}(\mu, \sigma^2 \mathbf{I}), \quad \nabla_x \log p(x) = \nabla_x \left(-\frac{1}{2\sigma^2} (x - \mu)^2 \right) = -\frac{x - \mu}{\sigma^2} = -\frac{\varepsilon}{\sigma}, \quad \varepsilon \sim \mathcal{N}(0, \mathbf{I})$$

$$q(x_t | x_0) \sim \mathcal{N}(\sqrt{\bar{\alpha}_t} x_0, (1 - \bar{\alpha}_t) \mathbf{I})$$

$$\text{And therefore } s_\theta(x_t, t) \approx \nabla_{x_t} \log q(x_t) = E_{q(x_0)} \nabla_{x_t} q(x_t | x_0) = E_{q(x_0)} \left[-\frac{\varepsilon_\theta(x_t, t)}{\sqrt{1 - \bar{\alpha}_t}} \right] = -\frac{\varepsilon_\theta(x_t, t)}{\sqrt{1 - \bar{\alpha}_t}}$$

Guided diffusion

- We have additional input y (a class label in classifier guided diffusion)
- We want to model a conditional distribution $p(x | y)$ instead



Conditioned on dogs

Classifier guided diffusion

We separately train a classifier $f_\phi(y \mid x_t, t)$ on a noisy image x_t , and use gradients $\nabla_x \log f_\phi(y \mid x_t)$ to guide the diffusion. Let us have a joint distribution $q(x_t, y)$, y is e.g. the image label

Classifier guided diffusion

We separately train a classifier $f_\phi(y \mid x_t, t)$ on a noisy image x_t , and use gradients $\nabla_x \log f_\phi(y \mid x_t)$

to guide the diffusion. Let us have a joint distribution $q(x_t, y)$, y is e.g. the image label

$$\nabla_{x_t} \log q(x_t \mid y) = \nabla_{x_t} \log q(x_t) + \nabla_{x_t} \log q(y \mid x_t) - \nabla_{x_t} \log q(y)$$

Classifier guided diffusion

We separately train a classifier $f_\phi(y | x_t, t)$ on a noisy image x_t , and use gradients $\nabla_x \log f_\phi(y | x_t)$

to guide the diffusion. Let us have a joint distribution $q(x_t, y)$, y is e.g. the image label

$$\nabla_{x_t} \log q(x_t | y) = \nabla_{x_t} \log q(x_t) + \nabla_{x_t} \log q(y | x_t) - \nabla_{x_t} \log q(y) \approx -\frac{1}{\sqrt{1 - \bar{\alpha}_t}} \varepsilon_\theta(x_t, t) + \nabla_{x_t} \log f_\phi(y | x_t)$$

Classifier guided diffusion

We separately train a classifier $f_\phi(y \mid x_t, t)$ on a noisy image x_t , and use gradients $\nabla_x \log f_\phi(y \mid x_t)$ to guide the diffusion. Let us have a joint distribution $q(x_t, y)$, y is e.g. the image label

$$\begin{aligned}\nabla_{x_t} \log q(x_t \mid y) &= \nabla_{x_t} \log q(x_t) + \nabla_{x_t} \log q(y \mid x_t) - \nabla_{x_t} \log q(y) \approx -\frac{1}{\sqrt{1 - \bar{\alpha}_t}} \varepsilon_\theta(x_t, t) + \nabla_{x_t} \log f_\phi(y \mid x_t) \\ &= -\frac{1}{\sqrt{1 - \bar{\alpha}_t}} (\varepsilon_\theta(x_t, t) - \sqrt{1 - \bar{\alpha}_t} \nabla_{x_t} \log f_\phi(y \mid x_t))\end{aligned}$$

Classifier guided diffusion

We separately train a classifier $f_\phi(y | x_t, t)$ on a noisy image x_t , and use gradients $\nabla_x \log f_\phi(y | x_t)$ to guide the diffusion. Let us have a joint distribution $q(x_t, y)$, y is e.g. the image label

$$\begin{aligned}\nabla_{x_t} \log q(x_t | y) &= \nabla_{x_t} \log q(x_t) + \nabla_{x_t} \log q(y | x_t) - \nabla_{x_t} \log q(y) \approx -\frac{1}{\sqrt{1 - \bar{\alpha}_t}} \varepsilon_\theta(x_t, t) + \nabla_{x_t} \log f_\phi(y | x_t) \\ &= -\frac{1}{\sqrt{1 - \bar{\alpha}_t}} (\varepsilon_\theta(x_t, t) - \sqrt{1 - \bar{\alpha}_t} \nabla_{x_t} \log f_\phi(y | x_t))\end{aligned}$$

Classifier guided predictor $\bar{\varepsilon}_\theta(x_t, t) = \varepsilon_\theta(x_t, t) - \sqrt{1 - \bar{\alpha}_t} w \nabla_{x_t} \log f_\phi(y | x_t)$

Classifier guided diffusion

We separately train a classifier $f_\phi(y | x_t, t)$ on a noisy image x_t , and use gradients $\nabla_x \log f_\phi(y | x_t)$ to guide the diffusion. Let us have a joint distribution $q(x_t, y)$, y is e.g. the image label

$$\begin{aligned}\nabla_{x_t} \log q(x_t | y) &= \nabla_{x_t} \log q(x_t) + \nabla_{x_t} \log q(y | x_t) - \nabla_{x_t} \log q(y) \approx -\frac{1}{\sqrt{1 - \bar{\alpha}_t}} \varepsilon_\theta(x_t, t) + \nabla_{x_t} \log f_\phi(y | x_t) \\ &= -\frac{1}{\sqrt{1 - \bar{\alpha}_t}} (\varepsilon_\theta(x_t, t) - \sqrt{1 - \bar{\alpha}_t} \nabla_{x_t} \log f_\phi(y | x_t))\end{aligned}$$

Classifier guided predictor $\bar{\varepsilon}_\theta(x_t, t) = \varepsilon_\theta(x_t, t) - \sqrt{1 - \bar{\alpha}_t} w \nabla_{x_t} \log f_\phi(y | x_t)$

Where w is the strength of the guidance

Classifier guided diffusion

Algorithm 1 Classifier guided diffusion sampling, given a diffusion model $(\mu_\theta(x_t), \Sigma_\theta(x_t))$, classifier $f_\phi(y|x_t)$, and gradient scale s .

Input: class label y , gradient scale s

$x_T \leftarrow$ sample from $\mathcal{N}(0, \mathbf{I})$

for all t from T to 1 **do**

$\mu, \Sigma \leftarrow \mu_\theta(x_t), \Sigma_\theta(x_t)$

$x_{t-1} \leftarrow$ sample from $\mathcal{N}(\mu + s\Sigma \nabla_{x_t} \log f_\phi(y|x_t), \Sigma)$

end for

return x_0

Downsides:

Classifier guided diffusion

Algorithm 1 Classifier guided diffusion sampling, given a diffusion model $(\mu_\theta(x_t), \Sigma_\theta(x_t))$, classifier $f_\phi(y|x_t)$, and gradient scale s .

Input: class label y , gradient scale s

$x_T \leftarrow$ sample from $\mathcal{N}(0, \mathbf{I})$

for all t from T to 1 **do**

$\mu, \Sigma \leftarrow \mu_\theta(x_t), \Sigma_\theta(x_t)$

$x_{t-1} \leftarrow$ sample from $\mathcal{N}(\mu + s\Sigma \nabla_{x_t} \log f_\phi(y|x_t), \Sigma)$

end for

return x_0

Downsides:

- The classifier has to cope with noise, might need to be trained separately

Classifier guided diffusion

Algorithm 1 Classifier guided diffusion sampling, given a diffusion model $(\mu_\theta(x_t), \Sigma_\theta(x_t))$, classifier $f_\phi(y|x_t)$, and gradient scale s .

Input: class label y , gradient scale s

$x_T \leftarrow$ sample from $\mathcal{N}(0, \mathbf{I})$

for all t from T to 1 **do**

$\mu, \Sigma \leftarrow \mu_\theta(x_t), \Sigma_\theta(x_t)$

$x_{t-1} \leftarrow$ sample from $\mathcal{N}(\mu + s\Sigma \nabla_{x_t} \log f_\phi(y|x_t), \Sigma)$

end for

return x_0

Downsides:

- The classifier has to cope with noise, might need to be trained separately
- If noise-robust might be inefficient: most of the information on x is irrelevant to y

Classifier free guidance

- Train a diffusion model $p(x | y)$ with conditioning dropout: conditional information is removed some percentage of the time

Classifier free guidance

- Train a diffusion model $p(x | y)$ with conditioning dropout: conditional information is removed some percentage of the time
- Resulting model can be conditional $p(x | y)$ and unconditional $p(x)$

Classifier free guidance

- Train a diffusion model $p(x | y)$ with conditioning dropout: conditional information is removed some percentage of the time
- Resulting model can be conditional $p(x | y)$ and unconditional $p(x)$
- No need for a separate classifier!

Classifier free guidance

Guide the diffusion without an independent classifier f_ϕ

Classifier free guidance

Guide the diffusion without an independent classifier f_ϕ

- Unconditional denoising diffusion model $p_\theta(x)$, parametrised through a score estimator $\varepsilon_\theta(x_t, t)$

Classifier free guidance

Guide the diffusion without an independent classifier f_ϕ

- Unconditional denoising diffusion model $p_\theta(x)$, parametrised through a score estimator $\varepsilon_\theta(x_t, t)$
- Conditional model $p_\theta(x | y)$ parametrised through $\varepsilon_\theta(x_t, t, y)$

Classifier free guidance

Guide the diffusion without an independent classifier f_ϕ

- Unconditional denoising diffusion model $p_\theta(x)$, parametrised through a score estimator $\varepsilon_\theta(x_t, t)$
- Conditional model $p_\theta(x | y)$ parametrised through $\varepsilon_\theta(x_t, t, y)$
- Conditional model is trained on paired data (x, y)

Classifier free guidance

Guide the diffusion without an independent classifier f_ϕ

- Unconditional denoising diffusion model $p_\theta(x)$, parametrised through a score estimator $\varepsilon_\theta(x_t, t)$
- Conditional model $p_\theta(x | y)$ parametrised through $\varepsilon_\theta(x_t, t, y)$
- Conditional model is trained on paired data (x, y)
- Conditional information y gets discarded periodically: $\varepsilon_\theta(x_t, t) = \varepsilon_\theta(x_t, t, y = \emptyset)$

Classifier free guidance

Guide the diffusion without an independent classifier f_ϕ

- Unconditional denoising diffusion model $p_\theta(x)$, parametrised through a score estimator $\varepsilon_\theta(x_t, t)$
- Conditional model $p_\theta(x | y)$ parametrised through $\varepsilon_\theta(x_t, t, y)$
- Conditional model is trained on paired data (x, y)
- Conditional information y gets discarded periodically: $\varepsilon_\theta(x_t, t) = \varepsilon_\theta(x_t, t, y = \emptyset)$

$$\nabla_{x_t} \log p(y | x_t) = \nabla_{x_t} \log p(x_t | y) - \nabla_{x_t} \log p(x_t)$$

Classifier free guidance

Guide the diffusion without an independent classifier f_ϕ

- Unconditional denoising diffusion model $p_\theta(x)$, parametrised through a score estimator $\varepsilon_\theta(x_t, t)$
- Conditional model $p_\theta(x | y)$ parametrised through $\varepsilon_\theta(x_t, t, y)$
- Conditional model is trained on paired data (x, y)
- Conditional information y gets discarded periodically: $\varepsilon_\theta(x_t, t) = \varepsilon_\theta(x_t, t, y = \emptyset)$

$$\nabla_{x_t} \log p(y | x_t) = \nabla_{x_t} \log p(x_t | y) - \nabla_{x_t} \log p(x_t) = -\frac{1}{\sqrt{1 - \bar{\alpha}_t}} (\varepsilon_\theta(x_t, t, y) - \varepsilon_\theta(x_t, t))$$

Classifier free guidance

Guide the diffusion without an independent classifier f_ϕ

- Unconditional denoising diffusion model $p_\theta(x)$, parametrised through a score estimator $\varepsilon_\theta(x_t, t)$
- Conditional model $p_\theta(x | y)$ parametrised through $\varepsilon_\theta(x_t, t, y)$
- Conditional model is trained on paired data (x, y)
- Conditional information y gets discarded periodically: $\varepsilon_\theta(x_t, t) = \varepsilon_\theta(x_t, t, y = \emptyset)$

$$\nabla_{x_t} \log p(y | x_t) = \nabla_{x_t} \log p(x_t | y) - \nabla_{x_t} \log p(x_t) = -\frac{1}{\sqrt{1 - \bar{\alpha}_t}} (\varepsilon_\theta(x_t, t, y) - \varepsilon_\theta(x_t, t))$$

$$\text{Now } \nabla_{x_t} \log p(x_t, y) = \nabla_{x_t} \log p(x_t) + \nabla_{x_t} \log p(y | x_t) = \nabla_{x_t} \log p(x_t | y) + \nabla_{x_t} \log y + \nabla_{x_t} p(y | x_t)$$

Classifier free guidance

Guide the diffusion without an independent classifier f_ϕ

- Unconditional denoising diffusion model $p_\theta(x)$, parametrised through a score estimator $\varepsilon_\theta(x_t, t)$
- Conditional model $p_\theta(x_t | y)$ parametrised through $\varepsilon_\theta(x_t, t, y)$
- Conditional model is trained on paired data (x, y)
- Conditional information y gets discarded periodically: $\varepsilon_\theta(x_t, t) = \varepsilon_\theta(x_t, t, y = \emptyset)$

$$\nabla_{x_t} \log p(y | x_t) = \nabla_{x_t} \log p(x_t | y) - \nabla_{x_t} \log p(x_t) = -\frac{1}{\sqrt{1 - \bar{\alpha}_t}} (\varepsilon_\theta(x_t, t, y) - \varepsilon_\theta(x_t, t))$$

$$\text{Now } \nabla_{x_t} \log p(x_t, y) = \nabla_{x_t} \log p(x_t) + \nabla_{x_t} \log p(y | x_t) = \nabla_{x_t} \log p(x_t | y) + \cancel{\nabla_{x_t} \log y} + \nabla_{x_t} p(y | x_t)$$

Classifier free guidance

$$\nabla_{x_t} \log p(x_t, y) = \nabla_{x_t} \log p(x_t | y) + \nabla_{x_t} \log p(y | x_t)$$

$$\nabla_{x_t} \log p(y | x_t) = -\frac{1}{\sqrt{1 - \bar{\alpha}_t}} (\varepsilon_{\theta}(x_t, t, y) - \varepsilon_{\theta}(x_t, t))$$

Hence, analogously to the case of classifier guided diffusion

Classifier free guidance

$$\nabla_{x_t} \log p(x_t, y) = \nabla_{x_t} \log p(x_t | y) + \nabla_{x_t} \log p(y | x_t)$$

$$\nabla_{x_t} \log p(y | x_t) = -\frac{1}{\sqrt{1 - \bar{\alpha}_t}} (\varepsilon_{\theta}(x_t, t, y) - \varepsilon_{\theta}(x_t, t))$$

Hence, analogously to the case of classifier guided diffusion

$$\bar{\varepsilon}_{\theta}(x_t, t, y) = \varepsilon_{\theta}(x_t, t, y) - \sqrt{1 - \bar{\alpha}_t} w \nabla_{x_t} \log p(y | x_t), \text{ where } \varepsilon_{\theta}(x_t, t, y) \text{ parametrises } p(x_t | y)$$

Classifier free guidance

$$\nabla_{x_t} \log p(x_t, y) = \nabla_{x_t} \log p(x_t | y) + \nabla_{x_t} \log p(y | x_t)$$

$$\nabla_{x_t} \log p(y | x_t) = -\frac{1}{\sqrt{1 - \bar{\alpha}_t}} (\varepsilon_{\theta}(x_t, t, y) - \varepsilon_{\theta}(x_t, t))$$

Hence, analogously to the case of classifier guided diffusion

$$\begin{aligned} \bar{\varepsilon}_{\theta}(x_t, t, y) &= \varepsilon_{\theta}(x_t, t, y) - \sqrt{1 - \bar{\alpha}_t} w \nabla_{x_t} \log p(y | x_t), \text{ where } \varepsilon_{\theta}(x_t, t, y) \text{ parametrises } p(x_t | y) \\ &= \varepsilon_{\theta}(x_t, t, y) + w(\varepsilon_{\theta}(x_t, t, y) - \varepsilon_{\theta}(x_t, t)) \end{aligned}$$