

Lecture 2. Computational methods

Markov Chain Monte Carlo, Laplace approximation

24.03.2025-28.03.2025 Instructors: Alina Bazarova, Jose Robledo

Why computational methods?

Recall that in our target formula for posterior $p(\theta | x) = \frac{p(\theta)p(x | \theta)}{\int_{\mathbb{R}} p(\theta)p(x | \theta)d\theta}$

where θ are our parameters. The **integral** below can get **really nasty**!

BUT: this integral is just a constant! Rewrite $p(\theta | x) = \frac{1}{Z}p(x, \theta)$, where Z is just a normalising constant, although possibly varying over a large range.

What to do?

Markov Chain Monte Carlo (MCMC) algorithm

Monte Carlo integration.

Assume we want to compute $E f(\theta | X) = \frac{\int f(\theta) p(\theta) p(X | \theta) d\theta}{\int p(\theta) p(X | \theta) d\theta}$

where f is some function of parameters θ given the data X .

Markov Chain Monte Carlo (MCMC) algorithm

Monte Carlo integration.

Assume we want to compute $E f(\theta | X) = \frac{\int f(\theta) p(\theta) p(X | \theta) d\theta}{\int p(\theta) p(X | \theta) d\theta}$

where f is some function of parameters θ given the data X .

Monte Carlo integration evaluates this integral by drawing **independent samples** $\{\theta_t, t = 1, \dots, n\}$ from posterior distribution $p(\theta | X)$

Markov Chain Monte Carlo (MCMC) algorithm

Monte Carlo integration.

Assume we want to compute $E f(\theta | X) = \frac{\int f(\theta) p(\theta) p(X | \theta) d\theta}{\int p(\theta) p(X | \theta) d\theta}$

where f is some function of parameters θ given the data X .

Monte Carlo integration evaluates this integral by drawing **independent samples** $\{\theta_t, t = 1, \dots, n\}$ from posterior distribution $p(\theta | X)$ and then approximating $E f(\theta | X) \approx \frac{1}{n} \sum_{t=1}^n f(\theta_t)$

Markov Chain Monte Carlo (MCMC) algorithm

Monte Carlo integration.

Assume we want to compute $E f(\theta | X) = \frac{\int f(\theta) p(\theta) p(X | \theta) d\theta}{\int p(\theta) p(X | \theta) d\theta}$

where f is some function of parameters θ given the data X .

Monte Carlo integration evaluates this integral by drawing **independent samples** $\{\theta_t, t = 1, \dots, n\}$ from posterior distribution $p(\theta | X)$ and then approximating $E f(\theta | X) \approx \frac{1}{n} \sum_{t=1}^n f(\theta_t)$

(law of large numbers)

Markov Chain Monte Carlo (MCMC) algorithm

However:

1. $p(\theta | X)$ can be non-standard, and hence sampling independently from it would not be feasible.

Markov Chain Monte Carlo (MCMC) algorithm

However:

1. $p(\theta | X)$ can be non-standard, and hence sampling independently from it would not be feasible.
2. Good news: $\{\theta_t\}$ does not necessarily need to be independent. One of the ways of tackling the above problem is to

Markov Chain Monte Carlo (MCMC) algorithm

However:

1. $p(\theta | X)$ can be non-standard, and hence sampling independently from it would not be feasible.
2. Good news: $\{\theta_t\}$ does not necessarily need to be independent. One of the ways of tackling the above problem is to do it through a Markov chain having $p(\theta | X)$ as its stationary distribution.

This is called Markov chain Monte Carlo.

MCMC algorithm II

Markov chain. Suppose we generate a sequence of random variables $\{\theta_0, \theta_1, \dots\}$.

MCMC algorithm II

Markov chain. Suppose we generate a sequence of random variables $\{\theta_0, \theta_1, \dots\}$.

Each time $t \geq 0$ the next state θ_{t+1} is sampled from a distribution $P(\theta_{t+1} \mid \theta_t)$, which depends **only on the current state of the chain** θ_t and does not depend on its history $\{\theta_0, \dots, \theta_{t-1}\}$.

MCMC algorithm II

Markov chain. Suppose we generate a sequence of random variables $\{\theta_0, \theta_1, \dots\}$.

Each time $t \geq 0$ the next state θ_{t+1} is sampled from a distribution $P(\theta_{t+1} | \theta_t)$, which depends **only on the current state of the chain** θ_t and does not depend on its history $\{\theta_0, \dots, \theta_{t-1}\}$.

Subject to certain conditions the chain will gradually “**forget**” **its initial state** θ_0 and the distribution $P(\theta_t | \theta_0)$ will not depend on t or θ_0 and converge to a unique stationary distribution

MCMC algorithm II

Markov chain. Suppose we generate a sequence of random variables $\{\theta_0, \theta_1, \dots\}$.

Each time $t \geq 0$ the next state θ_{t+1} is sampled from a distribution $P(\theta_{t+1} | \theta_t)$, which depends **only on the current state of the chain** θ_t and does not depend on its history $\{\theta_0, \dots, \theta_{t-1}\}$.

Subject to certain conditions the chain will gradually “**forget**” **its initial state** θ_0 and the distribution $P(\theta_t | \theta_0)$ will not depend on t or θ_0 and converge to a unique stationary distribution

Hence, after **sufficiently long burn-in** of m iterations points of $\{\theta_t, t = m + 1, \dots, n\}$ will be samples from the stationary distribution and the desired integral can be re-written as

MCMC algorithm II

Markov chain. Suppose we generate a sequence of random variables $\{\theta_0, \theta_1, \dots\}$.

Each time $t \geq 0$ the next state θ_{t+1} is sampled from a distribution $P(\theta_{t+1} | \theta_t)$, which depends **only on the current state of the chain** θ_t and does not depend on its history $\{\theta_0, \dots, \theta_{t-1}\}$.

Subject to certain conditions the chain will gradually “**forget**” its **initial state** θ_0 and the distribution $P(\theta_t | \theta_0)$ will not depend on t or θ_0 and converge to a unique stationary distribution

Hence, after **sufficiently long burn-in** of m iterations points of $\{\theta_t, t = m + 1, \dots, n\}$ will be samples from the stationary distribution and the desired integral can be re-written as

$$E f(\theta | X) \approx \frac{1}{n - m} \sum_{t=m+1}^n f(\theta_t)$$

MCMC algorithm II

Markov chain. Suppose we generate a sequence of random variables $\{\theta_0, \theta_1, \dots\}$.

Each time $t \geq 0$ the next state θ_{t+1} is sampled from a distribution $P(\theta_{t+1} | \theta_t)$, which depends **only on the current state of the chain** θ_t and does not depend on its history $\{\theta_0, \dots, \theta_{t-1}\}$.

Subject to certain conditions the chain will gradually “**forget**” its **initial state** θ_0 and the distribution $P(\theta_t | \theta_0)$ will not depend on t or θ_0 and converge to a unique stationary distribution

Hence, after **sufficiently long burn-in** of m iterations points of $\{\theta_t, t = m + 1, \dots, n\}$ will be samples from the stationary distribution and the desired integral can be re-written as

$$E f(\theta | X) \approx \frac{1}{n - m} \sum_{t=m+1}^n f(\theta_t)$$

Important: We can construct an MCMC algorithm which will have $p(\theta | X)$ as the stationary distribution!

Metropolis-Hastings sampler

At each time t the next state θ_{t+1} is chosen by first sampling a candidate Y from a ***proposal*** distribution $q(\cdot | \theta_t)$ which **depends only on the current state θ_t** (or not even that)

Metropolis-Hastings sampler

At each time t the next state θ_{t+1} is chosen by first sampling a candidate Y from a ***proposal*** distribution $q(\cdot | \theta_t)$ which **depends only on the current state** θ_t (or not even that)

Candidate Y is then accepted to be the next state of the chain with probability $\alpha(\theta_t, Y)$,
where $\alpha(\theta, Y) = \min \left(1, \frac{p(Y)p(X|Y)q(\theta|Y)}{p(\theta)p(X|\theta)q(Y|\theta)} \right)$.

Metropolis-Hastings sampler

At each time t the next state θ_{t+1} is chosen by first sampling a candidate Y from a ***proposal*** distribution $q(\cdot | \theta_t)$ which **depends only on the current state** θ_t (or not even that)

Candidate Y is then accepted to be the next state of the chain with probability $\alpha(\theta_t, Y)$,
where $\alpha(\theta, Y) = \min \left(1, \frac{p(Y)p(X|Y)q(\theta|Y)}{p(\theta)p(X|\theta)q(Y|\theta)} \right)$.

Now denote $\pi(\theta) = p(\theta | X)$

Metropolis-Hastings sampler

At each time t the next state θ_{t+1} is chosen by first sampling a candidate Y from a ***proposal*** distribution $q(\cdot | \theta_t)$ which **depends only on the current state** θ_t (or not even that)

Candidate Y is then accepted to be the next state of the chain with probability $\alpha(\theta_t, Y)$,
where $\alpha(\theta, Y) = \min \left(1, \frac{p(Y)p(X|Y)q(\theta|Y)}{p(\theta)p(X|\theta)q(Y|\theta)} \right)$.

Now denote $\pi(\theta) = p(\theta | X) = \frac{p(\theta)p(X|\theta)}{\int p(\theta)p(X|\theta)d\theta}$

$$P(\theta_{t+1} | \theta_t) = q(\theta_{t+1} | \theta_t)\alpha(\theta_t, \theta_{t+1}) + I(\theta_{t+1} = \theta_t)[1 - \int q(Y | \theta_t)\alpha(\theta_t, Y)dY] \quad (1)$$

Metropolis-Hastings sampler

At each time t the next state θ_{t+1} is chosen by first sampling a candidate Y from a **proposal** distribution $q(\cdot | \theta_t)$ which **depends only on the current state** θ_t (or not even that)

Candidate Y is then accepted to be the next state of the chain with probability $\alpha(\theta_t, Y)$,
where $\alpha(\theta, Y) = \min \left(1, \frac{p(Y)p(X|Y)q(\theta|Y)}{p(\theta)p(X|\theta)q(Y|\theta)} \right)$.

Now denote $\pi(\theta) = p(\theta | X) = \frac{p(\theta)p(X|\theta)}{\int p(\theta)p(X|\theta)d\theta}$

$$P(\theta_{t+1} | \theta_t) = q(\theta_{t+1} | \theta_t) \alpha(\theta_t, \theta_{t+1}) + I(\theta_{t+1} = \theta_t) \left[1 - \int q(Y | \theta_t) \alpha(\theta_t, Y) dY \right] \quad (1)$$

acceptance of candidate $Y = \theta_{t+1}$

Metropolis-Hastings sampler

At each time t the next state θ_{t+1} is chosen by first sampling a candidate Y from a **proposal** distribution $q(\cdot | \theta_t)$ which **depends only on the current state** θ_t (or not even that)

Candidate Y is then accepted to be the next state of the chain with probability $\alpha(\theta_t, Y)$,
where $\alpha(\theta, Y) = \min \left(1, \frac{p(Y)p(X|Y)q(\theta|Y)}{p(\theta)p(X|\theta)q(Y|\theta)} \right)$.

Now denote $\pi(\theta) = p(\theta | X) = \frac{p(\theta)p(X|\theta)}{\int p(\theta)p(X|\theta)d\theta}$

$$P(\theta_{t+1} | \theta_t) = q(\theta_{t+1} | \theta_t)\alpha(\theta_t, \theta_{t+1}) + I(\theta_{t+1} = \theta_t)[1 - \int q(Y | \theta_t)\alpha(\theta_t, Y)dY] \quad (1)$$

acceptance of candidate $Y = \theta_{t+1}$

rejection of all possible candidates Y

Metropolis-Hastings sampler II

Recall $\alpha(\theta, Y) = \min \left(1, \frac{\pi(Y)q(\theta | Y)}{\pi(\theta)q(Y | \theta)} \right)$, and hence

$$\pi(\theta_t)q(\theta_{t+1} | \theta_t)\alpha(\theta_t, \theta_{t+1}) = \pi(\theta_{t+1})q(\theta_t | \theta_{t+1})\alpha(\theta_{t+1}, \theta_t) \quad (2)$$

Metropolis-Hastings sampler II

Recall $\alpha(\theta, Y) = \min \left(1, \frac{\pi(Y)q(\theta | Y)}{\pi(\theta)q(Y | \theta)} \right)$, and hence

$$\pi(\theta_t)q(\theta_{t+1} | \theta_t)\alpha(\theta_t, \theta_{t+1}) = \pi(\theta_{t+1})q(\theta_t | \theta_{t+1})\alpha(\theta_{t+1}, \theta_t) \quad (2)$$

Hint: one of the α s in the equality above is equal to 1. Moreover, multiply (1) by $\pi(\theta_t)$

Metropolis-Hastings sampler II

Recall $\alpha(\theta, Y) = \min \left(1, \frac{\pi(Y)q(\theta | Y)}{\pi(\theta)q(Y | \theta)} \right)$, and hence

$$\pi(\theta_t)q(\theta_{t+1} | \theta_t)\alpha(\theta_t, \theta_{t+1}) = \pi(\theta_{t+1})q(\theta_t | \theta_{t+1})\alpha(\theta_{t+1}, \theta_t) \quad (2)$$

Hint: one of the α s in the equality above is equal to 1. Moreover, multiply (1) by $\pi(\theta_t)$

$$\pi(\theta_t)P(\theta_{t+1} | \theta_t) = \pi(\theta_t)q(\theta_{t+1} | \theta_t)\alpha(\theta_t, \theta_{t+1}) + \pi(\theta_t)I(\theta_{t+1} = \theta_t)[1 - \int q(Y | \theta_t)\alpha(\theta_t, Y)dY] \quad (3)$$

Metropolis-Hastings sampler II

Recall $\alpha(\theta, Y) = \min \left(1, \frac{\pi(Y)q(\theta | Y)}{\pi(\theta)q(Y | \theta)} \right)$, and hence

$$\pi(\theta_t)q(\theta_{t+1} | \theta_t)\alpha(\theta_t, \theta_{t+1}) = \pi(\theta_{t+1})q(\theta_t | \theta_{t+1})\alpha(\theta_{t+1}, \theta_t) \quad (2)$$

Hint: one of the α s in the equality above is equal to 1. Moreover, multiply (1) by $\pi(\theta_t)$

$$\pi(\theta_t)P(\theta_{t+1} | \theta_t) = \pi(\theta_t)q(\theta_{t+1} | \theta_t)\alpha(\theta_t, \theta_{t+1}) + \pi(\theta_t)I(\theta_{t+1} = \theta_t)[1 - \int q(Y | \theta_t)\alpha(\theta_t, Y)dY] \quad (3)$$

$$\pi(\theta_{t+1})P(\theta_t | \theta_{t+1}) = \pi(\theta_{t+1})q(\theta_t | \theta_{t+1})\alpha(\theta_{t+1}, \theta_t) + \pi(\theta_{t+1})I(\theta_{t+1} = \theta_t)[1 - \int q(Y | \theta_{t+1})\alpha(\theta_{t+1}, Y)dY] \quad (4)$$

Metropolis-Hastings sampler II

Recall $\alpha(\theta, Y) = \min \left(1, \frac{\pi(Y)q(\theta | Y)}{\pi(\theta)q(Y | \theta)} \right)$, and hence

$$\pi(\theta_t)q(\theta_{t+1} | \theta_t)\alpha(\theta_t, \theta_{t+1}) = \pi(\theta_{t+1})q(\theta_t | \theta_{t+1})\alpha(\theta_{t+1}, \theta_t) \quad (2)$$

Hint: one of the α s in the equality above is equal to 1. Moreover, multiply (1) by $\pi(\theta_t)$

$$\pi(\theta_t)P(\theta_{t+1} | \theta_t) = \pi(\theta_t)q(\theta_{t+1} | \theta_t)\alpha(\theta_t, \theta_{t+1}) + \pi(\theta_t)I(\theta_{t+1} = \theta_t)[1 - \int q(Y | \theta_t)\alpha(\theta_t, Y)dY] \quad (3)$$

$$\pi(\theta_{t+1})P(\theta_t | \theta_{t+1}) = \pi(\theta_{t+1})q(\theta_t | \theta_{t+1})\alpha(\theta_{t+1}, \theta_t) + \pi(\theta_{t+1})I(\theta_{t+1} = \theta_t)[1 - \int q(Y | \theta_{t+1})\alpha(\theta_{t+1}, Y)dY] \quad (4)$$

The first terms on the right-hand side of (3) and (4) are equal by (2), and the second ones by equality $\theta_t = \theta_{t+1}$, therefore

Metropolis-Hastings sampler II

Recall $\alpha(\theta, Y) = \min \left(1, \frac{\pi(Y)q(\theta | Y)}{\pi(\theta)q(Y | \theta)} \right)$, and hence

$$\pi(\theta_t)q(\theta_{t+1} | \theta_t)\alpha(\theta_t, \theta_{t+1}) = \pi(\theta_{t+1})q(\theta_t | \theta_{t+1})\alpha(\theta_{t+1}, \theta_t) \quad (2)$$

Hint: one of the α s in the equality above is equal to 1. Moreover, multiply (1) by $\pi(\theta_t)$

$$\pi(\theta_t)P(\theta_{t+1} | \theta_t) = \pi(\theta_t)q(\theta_{t+1} | \theta_t)\alpha(\theta_t, \theta_{t+1}) + \pi(\theta_t)I(\theta_{t+1} = \theta_t)[1 - \int q(Y | \theta_t)\alpha(\theta_t, Y)dY] \quad (3)$$

$$\pi(\theta_{t+1})P(\theta_t | \theta_{t+1}) = \pi(\theta_{t+1})q(\theta_t | \theta_{t+1})\alpha(\theta_{t+1}, \theta_t) + \pi(\theta_{t+1})I(\theta_{t+1} = \theta_t)[1 - \int q(Y | \theta_{t+1})\alpha(\theta_{t+1}, Y)dY] \quad (4)$$

The first terms on the left-hand side of (3) and (4) are equal by (2), and the second ones by equality $\theta_t = \theta_{t+1}$, therefore

$\pi(\theta_t)P(\theta_{t+1} | \theta_t) = \pi(\theta_{t+1})P(\theta_t | \theta_{t+1})$. Let us integrate both sides with respect to θ_t

Metropolis-Hastings sampler II

Recall $\alpha(\theta, Y) = \min \left(1, \frac{\pi(Y)q(\theta | Y)}{\pi(\theta)q(Y | \theta)} \right)$, and hence

$$\pi(\theta_t)q(\theta_{t+1} | \theta_t)\alpha(\theta_t, \theta_{t+1}) = \pi(\theta_{t+1})q(\theta_t | \theta_{t+1})\alpha(\theta_{t+1}, \theta_t) \quad (2)$$

Hint: one of the α s in the equality above is equal to 1. Moreover, multiply (1) by $\pi(\theta_t)$

$$\pi(\theta_t)P(\theta_{t+1} | \theta_t) = \pi(\theta_t)q(\theta_{t+1} | \theta_t)\alpha(\theta_t, \theta_{t+1}) + \pi(\theta_t)I(\theta_{t+1} = \theta_t)[1 - \int q(Y | \theta_t)\alpha(\theta_t, Y)dY] \quad (3)$$

$$\pi(\theta_{t+1})P(\theta_t | \theta_{t+1}) = \pi(\theta_{t+1})q(\theta_t | \theta_{t+1})\alpha(\theta_{t+1}, \theta_t) + \pi(\theta_{t+1})I(\theta_{t+1} = \theta_t)[1 - \int q(Y | \theta_{t+1})\alpha(\theta_{t+1}, Y)dY] \quad (4)$$

The first terms on the left-hand side of (3) and (4) are equal by (2), and the second ones by equality $\theta_t = \theta_{t+1}$, therefore

$\pi(\theta_t)P(\theta_{t+1} | \theta_t) = \pi(\theta_{t+1})P(\theta_t | \theta_{t+1})$. Let us integrate both sides with respect to θ_t

$$\int \pi(\theta_t)P(\theta_{t+1} | \theta_t)d\theta_t = \pi(\theta_{t+1}) \quad \textbf{Meaning:} \text{ if } \theta_t \text{ is from the distribution } \pi(.), \text{ then } \theta_{t+1} \text{ will be also.}$$

Metropolis-Hastings sampler II

Recall $\alpha(\theta, Y) = \min \left(1, \frac{\pi(Y)q(\theta | Y)}{\pi(\theta)q(Y | \theta)} \right)$, and hence

$$\pi(\theta_t)q(\theta_{t+1} | \theta_t)\alpha(\theta_t, \theta_{t+1}) = \pi(\theta_{t+1})q(\theta_t | \theta_{t+1})\alpha(\theta_{t+1}, \theta_t) \quad (2)$$

Hint: one of the α s in the equality above is equal to 1. Moreover, multiply (1) by $\pi(\theta_t)$

$$\pi(\theta_t)P(\theta_{t+1} | \theta_t) = \pi(\theta_t)q(\theta_{t+1} | \theta_t)\alpha(\theta_t, \theta_{t+1}) + \pi(\theta_t)I(\theta_{t+1} = \theta_t)[1 - \int q(Y | \theta_t)\alpha(\theta_t, Y)dY] \quad (3)$$

$$\pi(\theta_{t+1})P(\theta_t | \theta_{t+1}) = \pi(\theta_{t+1})q(\theta_t | \theta_{t+1})\alpha(\theta_{t+1}, \theta_t) + \pi(\theta_{t+1})I(\theta_{t+1} = \theta_t)[1 - \int q(Y | \theta_{t+1})\alpha(\theta_{t+1}, Y)dY] \quad (4)$$

The first terms on the left-hand side of (3) and (4) are equal by (2), and the second ones by equality $\theta_t = \theta_{t+1}$, therefore

$\pi(\theta_t)P(\theta_{t+1} | \theta_t) = \pi(\theta_{t+1})P(\theta_t | \theta_{t+1})$. Let us integrate both sides with respect to θ_t

$$\int \pi(\theta_t)P(\theta_{t+1} | \theta_t)d\theta_t = \pi(\theta_{t+1}) \quad \textbf{Meaning:} \text{ if } \theta_t \text{ is from the distribution } \pi(.), \text{ then } \theta_{t+1} \text{ will be also.}$$

Hence, once sample from stationary has been obtained, all subsequent samples are going to be from it. This means MCMC has converged. The period before convergence is called burn-in

Metropolis-Hastings: how it works in practice

1. Start at **current position** X .
2. Propose moving to a **new position** Y using proposal $q(Y|X)$
3. Accept/Reject the new position based on the position's adherence to the data and prior distributions using $\alpha(X, Y)$
 - If you accept: Move to the new position Y . Return to Step 1.
 - Else: Do not move to new position, stay at X . Return to Step 1.
4. After a large number of iterations, return **all accepted positions**.

Metropolis-Hastings sampler III

The natural question: **what should be the proposal distribution $q(Y | \theta)$?**

Metropolis-Hastings sampler III

The natural question: **what should be the proposal distribution $q(Y | \theta)$?**

1. The rate of convergence to the stationary distribution depends on it! And hence the **compute time**.

Metropolis-Hastings sampler III

The natural question: **what should be the proposal distribution $q(Y | \theta)$?**

1. The rate of convergence to the stationary distribution depends on it! And hence the **compute time**.
2. Even if the chain converged it may **mix** slowly (move around the states). And hence one needs to **run it for longer** to obtain **reliable estimates**.

Metropolis-Hastings sampler III

The natural question: **what should be the proposal distribution $q(Y | \theta)$?**

1. The rate of convergence to the stationary distribution depends on it! And hence the **compute time**.
2. Even if the chain converged it may **mix** slowly (move around the states). And hence one needs to **run it for longer** to obtain **reliable estimates**.
3. Proposal has to **explore the space efficiently**, sometimes it requires to perform experimentation and craftsmanship to construct a good one.

Jupyter notebook 2

Typical proposal distributions

Most typical one: **random walk**, $q(Y | \theta) = q(|Y - \theta|)$.

Typical proposal distributions

Most typical one: **random walk**, $q(Y | \theta) = q(|Y - \theta|)$.

Example: $Y \sim N(\theta_t, s)$, where N is a normal distribution and s is the custom standard deviation

Typical proposal distributions

Most typical one: **random walk**, $q(Y | \theta) = q(|Y - \theta|)$.

Example: $Y \sim N(\theta_t, s)$, where N is a normal distribution and s is the custom standard deviation

Important property: **acceptance rate** - how frequently the proposal gets accepted. Ideally should be 0.2-0.4

Typical proposal distributions

Most typical one: **random walk**, $q(Y | \theta) = q(|Y - \theta|)$.

Example: $Y \sim N(\theta_t, s)$, where N is a normal distribution and s is the custom standard deviation

Important property: **acceptance rate** - how frequently the proposal gets accepted. Ideally should be 0.2-0.4

This can be tuned during the **burn-in** period. In general:

Typical proposal distributions

Most typical one: **random walk**, $q(Y | \theta) = q(|Y - \theta|)$.

Example: $Y \sim N(\theta_t, s)$, where N is a normal distribution and s is the custom standard deviation

Important property: **acceptance rate - how frequently the proposal gets accepted. Ideally should be 0.2-0.4**

This can be tuned during the **burn-in** period. In general:

1. Acceptance **too high** -> chain mixes slowly. Acceptance **too low** -> chain stops moving.

Typical proposal distributions

Most typical one: **random walk**, $q(Y | \theta) = q(|Y - \theta|)$.

Example: $Y \sim N(\theta_t, s)$, where N is a normal distribution and s is the custom standard deviation

Important property: **acceptance rate - how frequently the proposal gets accepted. Ideally should be 0.2-0.4**

This can be tuned during the **burn-in** period. In general:

1. Acceptance **too high** -> chain mixes slowly. Acceptance **too low** -> chain stops moving.
2. **The larger the variance** of the proposal is the lower the acceptance rate is.

Typical proposal distributions

Most typical one: **random walk**, $q(Y | \theta) = q(|Y - \theta|)$.

Example: $Y \sim N(\theta_t, s)$, where N is a normal distribution and s is the custom standard deviation

Important property: **acceptance rate - how frequently the proposal gets accepted. Ideally should be 0.2-0.4**

This can be tuned during the **burn-in** period. In general:

1. Acceptance **too high** -> chain mixes slowly. Acceptance **too low** -> chain stops moving.
2. **The larger the variance** of the proposal is the lower the acceptance rate is.
3. This can be used during burn-in **to reach the desired acceptance** rate.

Single component MH and Gibbs sampler

Instead of updating θ *en bloc* it is often more convenient and computationally efficient to divide θ into components $\{\theta_1 \dots \theta_h\}$ and update them one by one.

Single component MH and Gibbs sampler

Instead of updating θ *en bloc* it is often more convenient and computationally efficient to divide θ into components $\{\theta_1 \dots \theta_h\}$ and update them one by one.

This means that instead of $q(Y | \theta)$ we will have $q(Y_i | \theta_{-i}, \theta_i)$, where $\theta_{-i} = \{\theta_1 \dots \theta_{i-1}, \theta_{i+1} \dots \theta_h\}$.

Single component MH and Gibbs sampler

Instead of updating θ *en bloc* it is often more convenient and computationally efficient to divide θ into components $\{\theta_1 \dots \theta_h\}$ and update them one by one.

This means that instead of $q(Y | \theta)$ we will have $q(Y_i | \theta_{-i}, \theta_i)$, where $\theta_{-i} = \{\theta_1 \dots \theta_{i-1}, \theta_{i+1} \dots \theta_h\}$.

Acceptance probability will then be $\alpha(\theta_{-i}, \theta_i, Y_i) = \min \left(1, \frac{\pi(Y_i | \theta_{-i}) q(\theta_i | Y_i, \theta_{-i})}{\pi(\theta_i | \theta_{-i}) q(Y_i | \theta_i, \theta_{-i})} \right)$

Single component MH and Gibbs sampler

Instead of updating θ *en bloc* it is often more convenient and computationally efficient to divide θ into components $\{\theta_1 \dots \theta_h\}$ and update them one by one.

This means that instead of $q(Y | \theta)$ we will have $q(Y_i | \theta_{-i}, \theta_i)$, where $\theta_{-i} = \{\theta_1 \dots \theta_{i-1}, \theta_{i+1} \dots \theta_h\}$.

Acceptance probability will then be $\alpha(\theta_{-i}, \theta_i, Y_i) = \min \left(1, \frac{\pi(Y_i | \theta_{-i}) q(\theta_i | Y_i, \theta_{-i})}{\pi(\theta_i | \theta_{-i}) q(Y_i | \theta_i, \theta_{-i})} \right)$

Gibbs sampler: $q(Y_i | \theta_i, \theta_{-i}) = \pi(Y_i | \theta_{-i})$. **Acceptance probability in this case is always equals to 1!**

Single component MH and Gibbs sampler

Instead of updating θ *en bloc* it is often more convenient and computationally efficient to divide θ into components $\{\theta_1 \dots \theta_h\}$ and update them one by one.

This means that instead of $q(Y | \theta)$ we will have $q(Y_i | \theta_{-i}, \theta_i)$, where $\theta_{-i} = \{\theta_1 \dots \theta_{i-1}, \theta_{i+1} \dots \theta_h\}$.

Acceptance probability will then be $\alpha(\theta_{-i}, \theta_i, Y_i) = \min \left(1, \frac{\pi(Y_i | \theta_{-i}) q(\theta_i | Y_i, \theta_{-i})}{\pi(\theta_i | \theta_{-i}) q(Y_i | \theta_i, \theta_{-i})} \right)$

Gibbs sampler: $q(Y_i | \theta_i, \theta_{-i}) = \pi(Y_i | \theta_{-i})$. **Acceptance probability in this case is always equal to 1!**

Gibbs sampling uses the property of tractability of all **conditional** posterior distributions to get samples from the unknown **full** posterior distribution of all model variables.

Gibbs sampling scheme

Assume we have data $X \sim p(X | \theta_1, \theta_2)$

1. Randomly initialize $\theta_1^{(0)}$ and sample $\theta_2^{(0)} \sim p(\theta_2 | X, \theta_1^{(0)})$

2. For step $t = 1, \dots, T$

(a) Sample $\theta_1^{(t)} \sim p(\theta_1 | X, \theta_2^{t-1})$

(b) Sample $\theta_2^{(t)} \sim p(\theta_2 | X, \theta_1^t)$