

**Input:** subjects attitude:  $A = \{a_i\}_{i=1}^n$ , where  $a_i \in \{0, 1\}$  and  $n \in \mathbb{N}$ ;  
distribution of subjects  $D_i$  with  $i \in I$  where  $I = \{1, \dots, 4\}$

**Output:** a set  $P$  with all pairs in the current session;  
an updated distribution of subjects  $D'_i$

**begin**

create an empty set  $P = \{\}$  for all pairs in this session

**while** *subject attitude sequence is not empty* **do**

create a positive and negative attitude set. Let  $i$  be such that

$a_i = j$ :

$S^+ = \{i : a_i = 1\}$

$S^- = \{i : a_i = 0\}$

**if**  $|A| > 1$  **then**

**if**  $|S^+| = 0$  **then**

**if** *the number of subjects in  $D_1 \leq 50$*  **then**

$P = P \cup \{S_1^-\} \times \{S_2^-\}$

$a_{i-}$ , where  $i = S_1^-$  and  $a_{j-}$ , where  $j = S_2^-$

$D'_i = \{(D_1 + 2), D_2, D_3, D_4\}$

**else**

assign  $a_i$  to study 2 and  $a_{i-}$ , where  $i = S_1^-$

assign  $a_j$  to study 2 and  $a_{j-}$ , where  $j = S_2^-$

**else if**  $|S^+| = 1$  **then**

**if** *the number of subjects in  $D_2 \leq 50 \wedge D_3 \leq 50$*  **then**

$P = P \cup \{S_1^-\} \times \{S_1^+\}$

$a_{i-}$ , where  $i = S_1^-$  and  $a_{j-}$ , where  $j = S_1^+$

$D'_i = \{D_1, (D_2 + 1), (D_3 + 1), D_4\}$

**else**

assign  $a_i$  to study 2 and  $a_{i-}$ , where  $i = S_1^-$

assign  $a_j$  to study 2 and  $a_{j-}$ , where  $j = S_1^+$

**else**

**if** *the number of subjects in  $D_4 \leq 50$*  **then**

$P = P \cup \{S_1^+\} \times \{S_2^+\}$

$a_{i-}$ , where  $i = S_1^+$  and  $a_{j-}$ , where  $j = S_2^+$

$D'_i = \{D_1, D_2, D_3, (D_4 + 2)\}$

**else**

assign  $a_i$  to study 2 and  $a_{i-}$ , where  $i = S_1^+$

assign  $a_j$  to study 2 and  $a_{j-}$ , where  $j = S_2^+$

**else**

assign user  $a_1$  to study 3 and empty  $A = \{\}$

**end**

**end**