

We consider sequences formed from the addition of terms of a given sequence. Let $\{a_n\}$, $n = 1, 2, 3, \dots$, be an arbitrary sequence of integer numbers; d a positive integer. We construct another sequence $\{b_m\}$, $m = 1, 2, 3, \dots$, by defining b_m as consisting of $n \times d$ occurrences of the term a_n :

$$b_1 = \underbrace{a_1, \dots, a_1}_{d \text{ occurrences of } a_1}, b_2 = \underbrace{a_2, \dots, a_2}_{2d \text{ occurrences of } a_2}, b_3 = \underbrace{a_3, \dots, a_3}_{3d \text{ occurrences of } a_3}, \dots$$

For example, if $a_n = n$, and $d = 1$, then the resulting sequence $\{b_m\}$ is:

$$\underbrace{1}_{b_1}, \underbrace{2, 2}_{b_2}, \underbrace{3, 3, 3}_{b_3}, \underbrace{4, 4, 4, 4}_{b_4}, \dots$$

Given a_n and d we want to obtain the corresponding k -th integer in the sequence $\{b_m\}$. For example, with $a_n = n$ and $d = 1$ we have 3 for $k = 6$; we have 4 for $k = 7$. With $a_n = n$ and $d = 2$, we have 2 for $k = 6$; we have 3 for $k = 7$.

Input

The first line of input contains C ($0 < C < 100$), the number of test cases that follows.
Each of the C test cases consists of three lines:

1. The first line represents a_n — a polynomial in n of degree i with non-negative integer coefficients in increasing order of the power:

$$a_n = c_0 + c_1n + c_2n^2 + c_3n^3 + \dots + c_in^i$$

where $c_j \in \mathbb{N}_0$, $j = 0, \dots, i$. This polynomial a_n is codified by its degree i followed by the coefficients c_j , $j = 0, \dots, i$. All the numbers are separated by a single space.

2. The second line is the positive integer d .
3. The third line is the positive integer k .

It is assumed that the polynomial a_n is a polynomial of degree less or equal than 20 ($1 \leq i \leq 20$) with non-negative integer coefficients less or equal than 10000 ($0 \leq c_j \leq 10000$, $j = 0, \dots, i$); $1 \leq d \leq 100000$; $1 \leq k \leq 1000000$.

Output

The output is a sequence of lines, one for each test case. Each of these lines contains the k -th integer in the sequence $\{b_m\}$ for the corresponding test case. This value is less or equal than $2^{63} - 1$.

Sample Input

```
2
4 3 0 0 0 23
25
100
1 0 1
1
6
```

Sample Output

```
1866
3
```