

Interval dependency and unequally spaced measurements


Modeling Intensive Longitudinal Data

Noémi K. Schuurman




Equal spacing between measurements

Researcher 1: half-hourly effect




Equal spacing between measurements

Researcher 1: half-hourly effect




Researcher 2: hourly effect




Equal spacing between measurements

Researcher 1: half-hourly effect




Researcher 2: hourly effect




Both sets of measurements are neatly evenly spaced.

Effects depend on interval size

Researcher 1: half-hourly effect



Researcher 2: hourly effect



AR model: effect depends on interval


► Half-hourly effect:

$$\phi_{res1} = .5$$

► Hourly effect: $\phi_{res2} = \phi_{res1}^2 = .25$

Unequal spacing between measurements

Researcher 1: half-hourly effect



Researcher 2: hourly effect



Researcher 3: blended effect



Unequal spacing between measurements

Solutions for unequally spaced measurements:

- ▶ AR models are discrete time models: *Continuous time models* can take the length of the time intervals into account.

Unequal spacing between measurements

Solutions for unequally spaced measurements:

- ▶ AR models are discrete time models: *Continuous time models* can take the length of the time intervals into account.
- ▶ Ad hoc solution: add in missing observations to equally space measurements

Unequal spacing between measurements

Solutions for unequally spaced measurements:

- ▶ AR models are discrete time models: *Continuous time models* can take the length of the time intervals into account.
- ▶ Ad hoc solution: add in missing observations to equally space measurements
 - TINTERVAL function in Mplus
 - r-package ...



**Utrecht
University**

Sharing science,
shaping tomorrow