

# Introduction to Differential Manifolds

---

**José Sá**

*Wolfson College, University of Cambridge, UK*

*E-mail:* [jac295@cam.ac.uk](mailto:jac295@cam.ac.uk)

ABSTRACT: Notes based on the lectures by Peter Gothen at University of Porto with personal inclusions (pun intended) here and there.

---

# Contents

<b>1</b>	<b>Differential Manifolds and Differentiable Maps</b>	<b>2</b>
1.1	Review of General Topology	2
1.2	Differential Manifolds	3
1.3	Differential Maps	6
1.4	Inverse Function Theorem	6
1.5	Submanifolds and the Submersion Theorem	6
1.6	Compactness	6
1.7	Quotient Manifolds	6
<b>2</b>	<b>Curves and Tangent Spaces</b>	<b>7</b>
2.1	Curves as Tangent Vectors	7
2.2	Tangent spaces of Submanifolds	7
2.3	Tangent vectors as Derivatives	7
2.4	The Tangent Bundle	7
2.5	Vector Fields	7
2.6	Integral Curves and 1-parameter Subgroup of Diffeomorphisms	7
2.7	Tangent vectors as Derivations	7
<b>3</b>	<b>Differential Forms</b>	<b>8</b>
3.1	One-forms and the Cotangent Bundle	8
3.2	Wedge Product and Exterior Algebra	8
3.3	Pullback of Forms	8
3.4	Orientation and Volume Forms	8
3.5	Integration in Differential Manifolds	8
3.6	Exterior Derivative	8
3.7	Manifolds with boundary	8
3.8	Stokes Theorem	8
3.9	Bubbling Forms	8
<b>4</b>	<b>de Rham Cohomology</b>	<b>9</b>
4.1	Definition	9
4.2	Aside: Cohomology as a Functor	9
4.3	Cochain Maps and Cochain Homotopies	9
4.4	Poincaré Lemma	9
4.5	Mayer-Vietories sequence	9
4.6	Degree of mappings	9
4.7	Index of a Vector Field and Poincaré-Hopf theorem	9
<b>5</b>	<b>Lie Groups as Manifolds</b>	<b>10</b>
5.1	Lie groups	10

5.2	Lie algebras	10
5.3	Exponential Map	10
5.4	Lie derivative	10

---

# 1 Differential Manifolds and Differentiable Maps

## 1.1 Review of General Topology

Let  $S$  be a set.

**Definition 1.1.** A topology is a collection  $\mathcal{T}$  of subsets of  $S$ , called the open sets, such that:

- (i)  $\emptyset, S \in \mathcal{T}$ , where  $\emptyset$  is the empty set.
- (ii) if  $U_\alpha \in \mathcal{T}$ , for  $\alpha \in A$ , then  $\bigcup_{\alpha \in A} U_\alpha \in \mathcal{T}$
- (iii) if  $U_1, \dots, U_n \in \mathcal{T}$ ,  $n \in \mathbb{N}$ , then  $\bigcap_{i=1}^n U_i \in \mathcal{T}$

**Example 1.1.**

- 1)  $S = \mathbb{R}^n$ ,  $U \in \mathcal{T}$  iff  $U \subseteq S$  is open in the usual sense.
- 2) If  $(S, d)$  is a metric space, then it is a topological space.


**Definition 1.2.** Let  $(S, \mathcal{T})$  be a topological space. A basis for the topology of  $S$  is the collection  $B \subseteq \mathcal{T}$  so that any  $U \in \mathcal{T}$  is the union of sets from  $B$ .

**Example 1.2.**

- 1)  $\{B(x; \epsilon) \mid x \in \mathbb{R}^n, \epsilon \in \mathbb{R}^+\}$  is a basis for the usual topology in  $\mathbb{R}^n$ .
- 2)  $\{B(x; \epsilon) \mid x \in \mathbb{Q}^n, \epsilon \in \mathbb{Q}^+\}$  is a countable basis for  $\mathbb{R}^n$ .

**Definition 1.3.**  $(S, \mathcal{T})$  is second-countable if the topology  $\mathcal{T}$  has a countable basis.

**Definition 1.4.**  $(S, \mathcal{T})$  is Hausdorff if for all  $x, y \in S$ , with  $x \neq y$ , there are open sets  $U, V \subseteq S$  so that  $x \in U$ ,  $y \in V$  and  $U \cap V = \emptyset$ .




The second-countability and “Hausdorffness” seems to be strange requirements but they are natural conditions, since The prototype of a manifold is the Euclidean space, which has these properties. If we do not impose these then many important results do not hold. For instance, we need Hausdorff condition in order to have unique convergence of sequences and uniqueness of ODE solutions. Without second-countability, it is not certain that we can embed the manifold in a finite-dimensional Euclidean space.

Let’s now define continuity in terms using topology language. Let  $X, Y$  be any two topological spaces.

**Definition 1.5.** A map  $f : X \longrightarrow Y$  is continuous at  $x \in X$  if for any open  $V \subseteq Y$  containing  $y$ , there exists a open  $U \subseteq X$  containing  $x$  so that  $f(U) \subseteq V$ .

**Definition 1.6.** A map  $f : X \rightarrow Y$  is continuous if it is continuous at all  $x \in X$ .

**Proposition 1.1.** A map  $f : X \rightarrow Y$  is continuous iff for all open  $V \subseteq Y$ , the preimage  $f^{-1}(V) \equiv \{x \in X \mid f(x) \in V\} \subseteq X$  is open in  $X$ .



**Figure 1:** Add caption to this figure

**Definition 1.7.** A set  $F \subseteq X$  is closed if  $X \setminus F \subseteq X$  is open.

**Proposition 1.2.** A map  $f : X \rightarrow Y$  is continuous iff  $f^{-1}(F) \subseteq X$  is closed in  $X$ , for every closed  $F \subseteq Y$  in  $Y$ .

**Definition 1.8.** A continuous map  $f : X \rightarrow Y$  is called a homeomorphism iff it has a continuous inverse  $g : Y \rightarrow X$ , such that  $f \circ g = \text{id}_Y$  and  $g \circ f = \text{id}_X$ . We say that  $X$  and  $Y$  are homeomorphic. We write  $g = f^{-1}$  and  $X \cong Y$ .

## 1.2 Differential Manifolds

**Definition 1.9.** A topological manifold of dimension  $m$  is a second countable, Hausdorff topological space  $M$ , so that any  $p \in M$  has a open neighborhood  $x \ni U \subseteq M$ , which is homeomorphic with  $\mathbb{R}^m$ . For convenience, we will sometimes write  $M^m \equiv M$  such that  $\dim(M) = m$ .

*Remark.* We might as well have said homeomorphic with an open subset of  $\mathbb{R}^m$ , because  $B(0, \epsilon) = \{x \in \mathbb{R}^m \mid \|x\| < \epsilon\} \cong \mathbb{R}^m$ . To see that we can use the function

$$f : B(0, \epsilon) \rightarrow \mathbb{R}^m$$

$$x \mapsto \frac{x}{\epsilon - \|x\|}$$

*Remark.* The *Theorem of Invariance of Domain* states that if exists  $\phi : \mathbb{R}^m \xrightarrow{\cong} \mathbb{R}^n$ , then  $m = n$ . Hence the definition is consistent.

**Definition 1.10.** A chart on a topological manifold  $M^m$  is a tuple  $(U, \phi)$ , where  $U \subseteq M$  is a open subset in  $M$  and  $\phi$  is a homeomorphism from  $U$  to  $V = \phi(U) \subseteq \mathbb{R}^m$  open in  $\mathbb{R}^m$ . If  $0 \in V$ , we say that the chart is centered on  $p$  if  $\phi(0) = p$ .

- $(U, \phi)$  is sometimes referred to as a coordinate patch.
- $(x_1(p), \dots, x_m(p)) = \phi(p)$  are called local coordinates of the point  $p$  in the chart  $(U, \phi)$ .

- $\phi^{-1}$  is also a homeomorphism and it is called a parametrisation.

We now are going to define the differential structure of the manifold.

**Definition 1.11.** A  $\mathcal{C}^0$ -atlas for  $M$  is a collection  $\mathcal{A} = \{(U_\alpha, \phi_\alpha) \mid \alpha \in A\}$  of charts so that  $\bigcup_{\alpha \in A} U_\alpha = M$ .


**Definition 1.12.** A  $\mathcal{C}^k$ -atlas for  $M$  is a  $\mathcal{C}^0$ -atlas such that

$$\theta_{\beta\alpha} \equiv \phi_\beta \circ (\phi_\alpha|_{U_{\alpha\beta}})^{-1} : \phi_\alpha(U_{\alpha\beta}) \longrightarrow \phi_\beta(U_{\alpha\beta}) ,$$

are differentiable of class  $\mathcal{C}^k$  for all  $\alpha, \beta \in A$ , where  $U_{\alpha\beta} \equiv U_\alpha \cap U_\beta$ .

*Remark.*

- The mappings  $\theta_{\alpha\beta}$  are called transition maps (mathematics) or coordinate transformations (physics).
- Note that  $U_{\alpha\beta}$  is open in  $U_\alpha$  and  $U_\beta$ , so both  $\phi_\alpha(U_{\alpha\beta})$  and  $\phi_\beta(U_{\alpha\beta})$  are open in  $\phi_\alpha(U_\alpha)$  and  $\phi_\beta(U_\beta)$ , respectively. Therefore  $\phi_\alpha(U_{\alpha\beta})$  and  $\phi_\beta(U_{\alpha\beta})$  are homeomorphic open sets in  $\mathbb{R}^m$ .



**Figure 2:** Figure describing homeomorphism between open sets in  $M$  and in  $\mathbb{R}^m$ , with transition maps  $\theta_{\alpha\beta}$ .

**Definition 1.13** (Provisional). A differential manifold of class  $\mathcal{C}^k$  is a topological manifold  $M$  together with a  $\mathcal{C}^k$  atlas.

*Remark* (Smoothness). A smooth atlas is one where the transitions functions are of the class  $\mathcal{C}^\infty$ , i.e. infinitely differentiable.

**Example 1.3.**

- (1)  $M = \mathbb{R}^m$  is a differential manifold, with the trivial smooth  $\mathcal{C}^\infty$ -atlas  $\mathcal{A} = \{(\mathbb{R}^m, \text{id})\}$ , where  $\text{id}(x) = x$ . Note that  $\mathbb{R}^m$  is open in  $\mathbb{R}^m$ .
- (2) Let  $M = S^m \equiv \{x \in \mathbb{R}^{m+1} \mid \|x\| = 1\}$ .

We write  $x = (x_0, x_1, \dots, x_m)$ . The stereographic projections from  $N = (1, 0, \dots, 0)$  and  $S = (-1, 0, \dots, 0)$  are given from the following charts

$$\begin{aligned} \phi_N : U_N \equiv S^m \setminus \{N\} &\longrightarrow \mathbb{R}^m & \phi_S : U_S \equiv S^m \setminus \{S\} &\longrightarrow \mathbb{R}^m \\ (x_0, \dots, x_m) &\longmapsto \frac{1}{1 - x_0}(x_1, \dots, x_m) & (x_0, \dots, x_m) &\longmapsto \frac{1}{1 + x_0}(x_1, \dots, x_m) \end{aligned}$$



**Figure 3:** Insert figure about describing the stereographic projection

*Fact.*  $\|\phi_N(p)\| \|\phi_S(p)\| = 1$ , for all  $p \in U_N \cap U_S$ .

$$\implies \theta_{NS}(y) = \phi_N \circ (\phi_S|_{U_{NS}})^{-1}(y) = \frac{(y_1, \dots, y_m)}{(y_1)^2 + \dots + (y_m)^2} = \frac{1}{\|y\|^2} y$$

Since this transition map is a map<sup>1</sup> from  $\mathbb{R}^m \setminus \{0\}$  to  $\mathbb{R}^m \setminus \{0\}$  of class  $\mathcal{C}^\infty$ , then  $S^m$  is a smooth  $m$ -dimensional manifold.

*Note.* The topology on  $S^m$  is the subspace topology, i.e.  $U \subset S^m$  is defined to be open if it is of the form  $U = S^m \cap V$  for some open  $V \subseteq \mathbb{R}^m$ .

**Definition 1.14.** Two charts  $(\phi_\alpha, U_\alpha)$  and  $(\phi_\beta, U_\beta)$  are said to be  $\mathcal{C}^k$ -compatible if  $\theta_{\beta\alpha} \equiv \phi_\beta \circ (\phi_\alpha|_{U_{\alpha\beta}})^{-1}$  and  $\theta_{\alpha\beta} = (\theta_{\beta\alpha})^{-1}$  are differentiable of class  $\mathcal{C}^k$ . Note if  $U \cap V = \emptyset$ , we still say the transition map is smooth.

**Definition 1.15.** A  $\mathcal{C}^k$ -atlas is maximal if it contains all  $\mathcal{C}^k$ -compatible charts.

<sup>1</sup>This is called the inversion map and is actually one-to-one in the  $m$ -dimensional punctured Euclidean space  $\mathbb{R}^m \setminus \{0\}$ . It maps the region  $0 < \|y\| < 1$  to the region  $\|y\| > 1$  and vice-versa, while leaving the unit sphere unchanged.

**1.3 Differential Maps**

**1.4 Inverse Function Theorem**

**1.5 Submanifolds and the Submersion Theorem**

**1.6 Compactness**

**1.7 Quotient Manifolds**



## 2 Curves and Tangent Spaces

### 2.1 Curves as Tangent Vectors

### 2.2 Tangent spaces of Submanifolds

### 2.3 Tangent vectors as Derivatives

### 2.4 The Tangent Bundle

### 2.5 Vector Fields

### 2.6 Integral Curves and 1-parameter Subgroup of Diffeomorphisms

### 2.7 Tangent vectors as Derivations

## 3 Differential Forms

### 3.1 One-forms and the Cotangent Bundle

### 3.2 Wedge Product and Exterior Algebra

### 3.3 Pullback of Forms

### 3.4 Orientation and Volume Forms

### 3.5 Integration in Differential Manifolds

### 3.6 Exterior Derivative

### 3.7 Manifolds with boundary

### 3.8 Stokes Theorem

### 3.9 Bubbling Forms

## 4 de Rham Cohomology

### 4.1 Definition

### 4.2 Aside: Cohomology as a Functor

### 4.3 Cochain Maps and Cochain Homotopies

### 4.4 Poincaré Lemma

### 4.5 Mayer-Vietories sequence

### 4.6 Degree of mappings

### 4.7 Index of a Vector Field and Poincaré-Hopf theorem

## 5 Lie Groups as Manifolds

### 5.1 Lie groups

### 5.2 Lie algebras

### 5.3 Exponential Map

### 5.4 Lie derivative

## References

- [1] D. Barden and C. B. Thomas, *An Introduction to Differential Manifolds* (Imperial College Press, 2003).
- [2] J. Lafontaine, *An Introduction to Differential Manifolds* (Springer International Publishing, 2015).
- [3] J. M. Lee, *Introduction to Smooth Manifolds*, Graduate Texts in Mathematics, Vol. 218 (Springer-Verlag New York, 2003).