

Testing fundamental physics with astrophysical and cosmological observations

José Sá^a , João G. Rosa^a

*^aDepartamento de Física, Universidade de Aveiro,
Campus de Santiago, 3810-183 Aveiro, Portugal*

E-mail: jose.a.sa@ua.pt, joao.rosa@ua.pt

Contents

1	Introduction	2
2	Superradiant scattering	3
	References	4

1 Introduction

Some text transitioning from MSc thesis to research work ...

2 Superradiant scattering

The general scattering theory of perturbations off rotating black holes (BH) has long been known in General Relativity (GR) [1]. We will take particular focus on neutral BHs since any intrinsic charge would increase electromagnetic (EM) forces on opposite-charged plasma. The Kerr BH is a vacuum solution for the Einstein's field equations, which generalized the well-known Schwarzschild spherical geometry.

In this work we use the Boyer-Linquist coordinates [2], where the metric takes the form

$$ds^2 = \left(1 - \frac{2Mr}{\rho^2}\right) dt^2 - 2a \sin^2 \theta \frac{(r^2 + a^2 - \Delta)}{\rho^2} dt d\varphi - \frac{(r^2 + a^2)^2 - \Delta a^2 \sin^2 \theta}{\rho^2} \sin^2 \theta d\varphi^2 - \frac{\rho^2}{\Delta} dr^2 - \rho^2 d\theta^2 \quad (2.1)$$

where $\Delta = r^2 - 2Mr + a^2$, $\bar{\rho} = r + ia \cos \theta$ and $\rho^2 \equiv |\bar{\rho}|^2$.

It was Newman and Penrose that develop the necessary formalism of spinor calculus for the study of perturbations [3]. The NP formalism focus on choosing a non-local tetrad $(\mathbf{l}, \mathbf{n}, \mathbf{m}, \bar{\mathbf{m}})$ of complex null vectors and projecting the relevant tensors in this basis. For example, for electromagnetic waves we use the Faraday to define the relevant NP scalars

$$\phi_0 = F_{\mu\nu} l^\mu m^\nu, \quad \phi_2 = F_{\mu\nu} \bar{m}^\mu n^\nu, \quad (2.2)$$

while for gravitational perturbations we use the Weyl tensor

$$\psi_0 = -C_{\mu\nu\sigma\rho} l^\mu m^\nu l^\sigma m^\rho, \quad \psi_4 = -C_{\mu\nu\sigma\rho} n^\mu \bar{m}^\nu n^\sigma \bar{m}^\rho. \quad (2.3)$$

Choosing a suitable tetrad [4], Teukolsky showed that is possible to obtain a separable wave equation for all types of massless perturbations (scalar, electromagnetic, gravitational) that are characterized by a *spin-weight* parameter s [5–8]. Due to the underlying symmetries of the geometry we can perform a mode decomposition of the form

$$\Upsilon_s = \int d\omega \sum_{\ell, m} e^{-i\omega t + im\varphi} {}_s S_{\ell m}(\theta) {}_s R_{\ell m}(r), \quad (2.4)$$

where Υ_0 obeys the Klein-Gordon wave equation in curved spacetime, $g^{\mu\nu} \nabla_\mu \partial_\nu \Upsilon_0 = 0$. The other bosonic perturbations are characterized by two polarizations mixed differently in the corresponding complex NP quantities $\Upsilon_{+1} = \phi_0$ and $\Upsilon_{-1} = 2(\bar{\rho}^*)^2 \phi_2$ for electromagnetic waves and $\Upsilon_{+2} = \psi_0$ and $\Upsilon_{-2} = 4(\bar{\rho}^*)^4 \psi_4$ for gravitational waves.

References

- [1] J. A. H. Futterman, F. A. Handler, and R. A. Matzner, *Scattering from Black Holes*, Cambridge Monographs on Mathematical Physics (Cambridge University Press, 1988).
- [2] R. H. Boyer and R. W. Lindquist, *Maximal Analytic Extension of the Kerr Metric*, [J. Math. Phys. **8**, 265 \(1967\)](#).
- [3] E. Newman and R. Penrose, *An Approach to Gravitational Radiation by a Method of Spin Coefficients*, [J. Math. Phys. **3**, 566 \(1962\)](#).
- [4] W. Kinnersley, *Type D Vacuum Metrics*, [J. Math. Phys. **10**, 1195 \(1969\)](#).
- [5] S. A. Teukolsky, *Rotating Black Holes: Separable Wave Equations for Gravitational and Electromagnetic Perturbations*, [Phys. Rev. Lett. **29**, 1114 \(1972\)](#).
- [6] S. A. Teukolsky, *Perturbations of a rotating black hole. I. Fundamental equations for gravitational electromagnetic and neutrino field perturbations*, [Astrophys. J. **185**, 635 \(1973\)](#).
- [7] W. H. Press and S. A. Teukolsky, *Perturbations of a Rotating Black Hole. II. Dynamical Stability of the Kerr Metric*, [Astrophys. J. **185**, 649 \(1973\)](#).
- [8] S. A. Teukolsky and W. H. Press, *Perturbations of a rotating black hole. III. Interaction of the hole with gravitational and electromagnetic radiation*, [Astrophys. J. **193**, 443 \(1974\)](#).