

Superradiant scattering at Kerr black holes

José Sá

Mestrado em Física

Departamento de Física e Astronomia



2017

Orientador

[João Rosa](#), Professor Auxiliar Convidado, [Faculdade de Ciências da Universidade do Porto](#)

Coorientador

[Orfeu Bertolami](#), Professor Catedrático, [Faculdade de Ciências da Universidade do Porto](#)







Todas as correções determinadas
pelo júri, e só essas, foram efetuadas.

O Presidente do Júri,

Porto, ____/____/____



UNIVERSIDADE DO PORTO

MASTER'S THESIS

Superradiant scattering at Kerr black holes

Author:

José SÁ

Supervisor:

João ROSA

Co-supervisor:

Orfeu BERTOLAMI

*A thesis submitted in fulfilment of the requirements
for the degree of Master of Science*

at the

Faculdade de Ciências da Universidade do Porto
Departamento de Física e Astronomia

July 2017

Acknowledgements

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

UNIVERSIDADE DO PORTO

Abstract

Faculdade de Ciências da Universidade do Porto

Departamento de Física e Astronomia

Master of Science

Superradiant scattering at Kerr black holes

by José Sá

The Thesis Abstract is written here (and usually kept to just this page). The page is kept centered vertically so can expand into the blank space above the title too...

UNIVERSIDADE DO PORTO

Resumo

Faculdade de Ciências da Universidade do Porto

Departamento de Física e Astronomia

Mestre de Ciência

Superradiant scattering at Kerr black holes

por José Sá

Tradução em português do “Abstract” escrito em inglês mais a cima. A página é centrada vertical e horizontalmente, podendo expandir para o espaço superior da página em branco ...

Contents

Acknowledgements	iii
Abstract	v
Resumo	vii
Contents	ix
List of Figures	xi
List of Tables	xiii
Notation and Conventions	xv
Abbreviations	xvii
1 Superradiance	1
1.1 Introduction	1
1.2 Black hole superradiance	2
2 Mathematical preliminaries	3
2.1 General Relativity	3
2.2 Kerr black hole	3
2.3 Kinnersley tetrad	3
2.4 Newman-Penrose formalism	3
3 Teukolsky master equation	5
3.1 Angular solutions	5
3.2 Asymptotic radial solution	5
3.3 Amplification factor Z_{slm}	5
4 Numerical method	7
4.1 Eigenvalues	7
4.1.1 Leaver method	7
4.1.2 Spectral	7
4.2 Radial ansatz	7
4.3 Amplification factor as a first test	7

5	Scattering problem	9
5.1	Plane wave decomposition	9
A	Spin-weighted spherical harmonics	11
A.1	Connection with spheroidal harmonics	11
A.2	Spin raising/lowering differential operators	12
A.3	Generalized addition of angular momentum formula	12
A.4	Some useful harmonics	12
	Bibliography	13

List of Figures

List of Tables

Notation and Conventions

Units

Unit conventions

Tensors and GR related

Metric definitions and stuff

Abbreviations

QM	Quantum Mechanics
BH	Black Hole
GR	General Relativity
GW	Gravitational Wave
LIGO	Laser Interferometric Gravitational Wave Observatory
SWSH	Spin-Weighted Spheroidal Harmonic

Chapter 1

Superradiance

1.1 Introduction

The first appearance of the concept of *superradiance* was in 1954, when Dicke [1] showed that a gas could be excited by a pulse into “superradiant states” from thermal equilibrium and then emit coherent radiation.

Zel’dovich [2, 3] showed that a absorbing surface rotating with an angular velocity Ω could scatter incident wave with frequency ω which satisfies

$$\omega - m \Omega < 0 \tag{1.1}$$

where m is the usual azimuthal number of the monochromatic plane wave relative to the rotation axis. Condition (1.1) was to become one of the most important results of (rotational) superradiance as it presents itself in multiple examples in the literature, *i.e.* Vavilov-Cherekov effect and anomalous Doppler effect.

Actually, radiation amplification can be traced to birth of Quantum Mechanics, in the beginnings of the 20th century. First studies of the Dirac equation by Klein [4] revealed the possibility of electrons propagating in a region with a sufficiently large potential barrier without the expected dampening from non-relativistic QM tunnel effect. Due to some confusion, this result was wrongly interpreted by some authors as fermionic superradiance, as if the reflected current by the barrier could be greater than the incident current. The problem was named *Klein paradox* by Sauter [5] and this misleading result was due to a incorrect calculation of the group velocities of the reflected and transmitted waves.

Today, it is known that fermionic currents cannot be amplified for this particular problem [6], result that was correctly obtained by Klein in his original paper. On the contrary, superradiant scattering can indeed occur for bosonic fields.

1.2 Black hole superradiance

Chapter 2

Mathematical preliminaries

2.1 General Relativity

2.2 Kerr black hole

2.3 Kinnersley tetrad

2.4 Newman-Penrose formalism

Chapter 3

Teukolsky master equation

3.1 Angular solutions

3.2 Asymptotic radial solution

3.3 Amplification factor Z_{slm}

Chapter 4

Numerical method

4.1 Eigenvalues

4.1.1 Leaver method

4.1.2 Spectral

4.2 Radial ansatz

4.3 Amplification factor as a first test

Chapter 5

Scattering problem

5.1 Plane wave decomposition

Appendix A

Spin-weighted spherical harmonics

SWSHs play an important role in BH physics and was first introduced by Teukolsky when considering non-scalar wave perturbations on a Kerr background, obtaining a separable master equation in four dimensions. After the usual change of coordinates, the polar differential equation goes as

$$\frac{1}{S} \frac{d}{dx} \left((1-x^2) \frac{dS}{dx} \right) + (cx)^2 - 2csx - \frac{(m+sx)^2}{1-x^2} + s = -\lambda \quad (\text{A.1})$$

with $x = \cos \theta$, where λ is the eigenvalue for a given SWSH solution. Periodic boundary conditions on the azimuthal wave function constrains m to the integers.

A.1 Connection with spheroidal harmonics

By setting $s = 0$ (scalar) and $c = 0$ (spherical), then it's clear that (A.1) appears as a generalization of the spherical harmonics equation. In this last case, the solution are given by the associated Legendre polynomials, $P_\ell^m(x)$, for which the eigenvalue is $\ell(\ell+1)$, restricted to the condition of $|m| \leq \ell$. The closed form for spherical harmonics, after normalization, is

$${}_0Y_\ell^m(x) = (-1)^m \sqrt{\frac{(2\ell+1)}{4\pi} \frac{(\ell-m)!}{(\ell+m)!}} P_\ell^m(x) \quad (\text{A.2})$$

where P_ℓ^m are the associated Legendre polynomials which can be obtained using the famous Rodrigues' formula.

A.2 Spin raising/lowering differential operators**A.3 Generalized addition of angular momentum formula****A.4 Some useful harmonics**

Bibliography

- [1] R. H. Dicke, *Coherence in Spontaneous Radiation Processes*, [Phys. Rev. **93**, 99 \(1954\)](#).
- [2] Y. B. Zel'dovich, *Generation of Waves by a Rotating Body*, [JETP Lett. **14**, 180 \(1971\)](#), [*Zh. Eksp. Teor. Fiz. Pis'ma Red.* **14**, 270 (1971)].
- [3] Y. B. Zel'dovich, *Amplification of Cylindrical Electromagnetic Waves Reflected from a Rotating Body*, [Sov. Phys. JETP **35**, 1085 \(1972\)](#), [*Zh. Eksp. Teor. Fiz.* **62**, 2076 (1972)].
- [4] O. Klein, *Die Reflexion von Elektronen an einem Potentialsprung nach der relativistischen Dynamik von Dirac*, [Z. Phys. **53**, 157 \(1929\)](#).
- [5] F. Sauter, *Über das Verhalten eines Elektrons im homogenen elektrischen Feld nach der relativistischen Theorie Diracs*, [Z. Phys. **69**, 742 \(1931\)](#).
- [6] C. A. Manogue, *The Klein paradox and superradiance*, [Ann. Phys. **181**, 261 \(1988\)](#).
- [7] J. G. Rosa, *Superradiance in the sky*, [Phys. Rev. **D95**, 064017 \(2017\)](#), [arXiv:1612.01826 \[gr-qc\]](#).
- [8] S. Chandrasekhar, *The Mathematical Theory of Black Holes*, Oxford Classic Texts in the Physical Sciences (Clarendon Press, 1998).
- [9] A. A. Starobinsky, *Amplification of waves reflected from a rotating "black hole"*, [Sov. Phys. JETP **37**, 28 \(1973\)](#), [*Zh. Eksp. Teor. Fiz.* **64**, 48 (1973)].
- [10] W. H. Press and S. A. Teukolsky, *Perturbations of a Rotating Black Hole. II. Dynamical Stability of the Kerr Metric*, [Astrophys. J. **185**, 649 \(1973\)](#).
- [11] S. A. Teukolsky and W. H. Press, *Perturbations of a rotating black hole. III. Interaction of the hole with gravitational and electromagnetic radiation*, [Astrophys. J. **193**, 443 \(1974\)](#).
- [12] S. A. Teukolsky, *Perturbations of a rotating black hole. I. Fundamental equations for gravitational electromagnetic and neutrino field perturbations*, [Astrophys. J. **185**, 635 \(1973\)](#).

- [13] S. A. Teukolsky, *Rotating Black Holes: Separable Wave Equations for Gravitational and Electromagnetic Perturbations*, [Phys. Rev. Lett. **29**, 1114 \(1972\)](#).
- [14] E. Berti, V. Cardoso and M. Casals, *Eigenvalues and eigenfunctions of spin-weighted spheroidal harmonics in four and higher dimensions*, [Phys. Rev. **D73**, 02401 \(2006\)](#), [Erratum: [Phys. Rev. **D73**, 109902 \(2006\)](#)], [arXiv:gr-qc/051111 \[gr-qc\]](#) .
- [15] R. Brito, V. Cardoso and P. Pani, *Superradiance: Energy Extraction, Black-Hole Bombs and Implications for Astrophysics and Particle Physics*, Lecture Notes in Physics, Vol. 906 (Springer International Publishing, 2015) [arXiv:1501.06570](#) .
- [16] R. G. Winter, *Klein Paradox for the Klein-Gordon Equation*, [Am. J. Phys. **27**, 355 \(1959\)](#).