

**Preoperative Cognitive Profile Predictive of Cognitive Decline after
Subthalamic Deep Brain Stimulation in Parkinson's Disease**

Josef Mana¹, Ondrej Bezdicek¹, Filip Ruzicka¹, Anna Fecikova¹, Olga Klempirova¹,
Tomas Nikolai¹, Tereza Uhrova¹, Evzen Ruzicka¹, Dusan Urgosik², and Robert Jech¹

¹Department of Neurology and Centre of Clinical Neuroscience, First Faculty of
Medicine and General University Hospital in Prague, Charles University, Czech
Republic

²Department of stereotactic and radiation neurosurgery, Na Homolce Hospital, Prague,
Czech Republic

Author Note

Josef Mana  <https://orcid.org/0000-0002-7817-3978>

Ondrej Bezdicek  <https://orcid.org/0000-0002-5108-0181>

Robert Jech  <https://orcid.org/0000-0002-9732-8947>

Author roles were classified using the Contributor Role Taxonomy (CRediT; <https://credit.niso.org/>) as follows: *Josef Mana*: Conceptualization, Data curation, Investigation, Formal analysis, Methodology, Software, Visualization, Writing - original draft, Writing - review & editing; *Ondrej Bezdicek*: Conceptualization, Data curation, Investigation, Methodology, Supervision, Writing - original draft, Writing - review & editing; *Filip Ruzicka*: Investigation, Writing - review & editing; *Anna Fecikova*: Investigation; *Olga Klempirova*: Investigation; *Tomas Nikolai*: Investigation, Writing - review & editing; *Tereza Uhrova*: Investigation; *Evzen Ruzicka*: Conceptualization, Funding acquisition, Investigation, Writing - review & editing; *Dusan Urgosik*: Investigation; *Robert Jech*: Conceptualization, Data curation, Funding acquisition, Investigation, Resources, Supervision, Writing - review & editing

Correspondence concerning this article should be addressed to Ondrej Bezdicek, Email: ondrej.bezdicek@gmail.com

Abstract

Cognitive decline represents a severe non-motor symptom of Parkinson's disease (PD) that can significantly reduce benefits of subthalamic deep brain stimulation (STN DBS). Here, we aimed to identify pre-surgery cognitive profile associated with faster post-surgery cognitive decline in STN DBS treated PD patients to characterize patients who could benefit from more monitoring during treatment course. A retrospective observational study of 126 PD patients treated by STN DBS combined with oral dopaminergic therapy followed for 3.54 years on average ($SD = 2.32$) with repeated assessments of cognition was conducted. Pre-surgery cognitive profile was obtained via a comprehensive neuropsychological examination. Data were analyzed using exploratory factor analysis for pre-surgery cognitive profile extraction and Bayesian generalized linear mixed models for description of the longitudinal cognitive outcome. Overall, we observed a mild annual cognitive decline of 0.90 points from a total of 144 points in the Mattis Dementia Rating Scale (95% posterior probability interval (PPI) [-1.19, -0.62]). Pre-surgery executive deficit predicted the rate of post-surgery cognitive decline ($b = -0.39$, 95% PPI [-0.63, -0.15]). The predictive utility of pre-surgery executive deficit resulted from summing small effects of several single test scores. Patients with PD treated with STN DBS experience only mild annual post-surgery cognitive decline. According to our data and models patients with worse long-term cognitive prognosis can be identified via pre-surgery examination of executive functions. Aggregating results from multiple executive tests to estimate cognitive prognosis of PD patients treated with STN DBS is likely superior to examining single test scores.

Keywords: Parkinson's disease, deep brain stimulation, cognition, longitudinal, latent variable analysis

**Preoperative Cognitive Profile Predictive of Cognitive Decline after
Subthalamic Deep Brain Stimulation in Parkinson's Disease**